

**Package leaflet: information for the user**  
**Solifenacin succinate 5 mg Film-coated Tablets**  
**Solifenacin succinate 10 mg Film-coated Tablets**  
Solifenacin succinate

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, or pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Solifenacin succinate film-coated tablets is and what it is used for
2. What you need to know before you take Solifenacin succinate film-coated tablets
3. How to take Solifenacin succinate film-coated tablets
4. Possible side effects
5. How to store Solifenacin succinate film-coated tablets
6. Contents of the pack and other information

**1. What Solifenacin succinate film-coated tablets is and what it is used for**

The active substance of Solifenacin succinate film-coated tablets belongs to the group called anticholinergics. These medicines are used to reduce the activity of an overactive bladder. This enables you to wait longer before having to go to the bathroom and increases the amount of urine that can be held by your bladder.

Solifenacin succinate film-coated tablets is used to treat the symptoms of a condition called overactive bladder. These symptoms include: having a strong, sudden urge to urinate without prior warning, having to urinate frequently or wetting yourself because you could not get to the bathroom in time.

**2. What you need to know before you take Solifenacin succinate film-coated tablets**

**Do not take Solifenacin succinate film-coated tablets**

- if you are allergic to solifenacin succinate or any of the other ingredients of this medicine (listed in section 6).
- if you have inability to pass water or to empty your bladder completely (urinary retention).
- if you have a severe stomach or bowel condition (including toxic megacolon, a complication associated with ulcerative colitis).
- if you suffer from the muscle disease called myasthenia gravis, which can cause an extreme weakness of certain muscles.
- if you suffer from increased pressure in the eyes, with gradual loss of eye sight (glaucoma).
- if you are undergoing kidney dialysis.
- if you have severe liver disease.
- if you suffer from severe kidney disease or moderate liver disease AND at the same time are being treated with medicines that may decrease the removal of Solifenacin succinate film-coated tablets from the body (for example, ketoconazole). Your doctor or pharmacist will have informed you if this is the case.

- Inform your doctor if you have or ever had any of the above mentioned conditions before treatment with Solifenacin succinate film-coated tablets starts.

### **Warnings and precautions**

Talk to your doctor or pharmacist before taking Solifenacin succinate film-coated tablets

- if you have trouble emptying your bladder (bladder obstruction) or have difficulty in passing urine, e.g. a thin urine flow. Risk of accumulation of urine in the bladder is much higher.
- if you have some obstruction of the digestive system (constipation).
- if you are at risk of your digestive system slowing down (stomach and bowel movements). Your doctor will have informed you if this is the case.
- if you suffer from severe kidney disease.
- if you have moderate liver disease.
- if you have a stomach tear (hiatus hernia) or heartburn.
- if you have a nervous disorder (autonomic neuropathy).

### **Children and adolescents**

**Solifenacin succinate film-coated tablets** is not to be used in children or adolescents under 18 years.

Inform your doctor if you have or ever had any of the above mentioned conditions before treatment with Solifenacin succinate film-coated tablets starts.

Before starting Solifenacin succinate film-coated tablets, your doctor will assess whether there are other causes for your need to pass urine frequently, for example heart failure (insufficient pumping power of the heart) or kidney disease. If you have a urinary tract infection, your doctor will prescribe you an antibiotic (a treatment against particular bacterial infections).

### **Other medicines and Solifenacin succinate film-coated tablets**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

It is especially important to inform your doctor if you are taking:

- other anticholinergic medicines, as the effects and side effects of both medications could be increased.
- cholinergics, as they can reduce the effect of Solifenacin succinate film-coated tablets.
- medicines, like metoclopramide and cisapride, which make the digestive system work faster. Solifenacin succinate film-coated tablets can reduce their effect.
- medicines, like ketoconazole, itraconazole (medicines used to treat fungal infections), ritonavir, nelfinavir (medicines used to treat HIV infections) and verapamil, diltiazem (medicines used to treat high blood pressure and heart diseases). These medicines decrease the rate at which Solifenacin succinate film-coated tablets is broken down by the body.
- medicines like rifampicin (medicine used to treat tuberculosis and other bacterial infections) and phenytoin, carbamazepine (medicines used to treat epilepsy). They may increase the rate at which Solifenacin succinate film-coated tablets is broken down by the body.
- medicines such as bisphosphonates, that can cause or exacerbate inflammation of the gullet (oesophagitis).

### **Solifenacin succinate film-coated tablets with food, drink and alcohol**

Solifenacin succinate film-coated tablets can be taken with or without food, depending on your preference.

### **Pregnancy, breast-feeding and fertility**

You should not use Solifenacin succinate film-coated tablets if you are pregnant unless your doctor thinks it is necessary. Do not use Solifenacin succinate film-coated tablets if you are breast-feeding as solifenacin may pass into your breast milk.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

### **Driving and using machines**

Solifenacin succinate film-coated tablets may cause blurred vision and sometimes sleepiness or tiredness. If you suffer from any of these side effects, do not drive or operate machinery.

### **Solifenacin succinate film-coated tablets contains lactose monohydrate**

If you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking this medicinal product.

## **3. How to take Solifenacin succinate film-coated tablets**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is 5 mg per day, unless your doctor has told you to take 10 mg per day.

You should swallow the tablet whole with some liquid, e.g. a glass of water. It can be taken with or without food, according to your preference. Do not crush the tablets.

### **Use in children and adolescents**

Solifenacin succinate film-coated tablets is not to be used in children or adolescents under 18 years.

### **If you take more Solifenacin succinate film-coated tablets than you should**

If you have taken too much Solifenacin succinate film-coated tablets or if a child has accidentally taken Solifenacin succinate film-coated tablets, contact your doctor or pharmacist immediately.

Symptoms of overdose may include: headache, dry mouth, dizziness, drowsiness and blurred vision, perceiving things that are not there (hallucinations), over-excitability, seizures (convulsions), difficulty breathing, elevated heart rate (tachycardia), accumulation of urine in the bladder (urinary retention) and dilated pupils (mydriasis).

### **If you forget to take Solifenacin succinate film-coated tablets**

If you forget to take a dose at the usual time, take it as soon as you remember, unless it is time to take your next dose. Never take more than one dose per day. If you are in doubt, always consult your doctor or pharmacist. Do not take a double dose to make up for a forgotten dose.

### **If you stop taking Solifenacin succinate film-coated tablets**

If you stop taking Solifenacin succinate film-coated tablets, your symptoms of overactive bladder may return or worsen. Always consult your doctor, if you are considering stopping the treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them. If you experience an allergic attack, or a severe skin reaction (e.g. blistering and peeling of the skin), you must inform your doctor or pharmacist immediately.

Angioedema (skin allergy that results in the swelling that occurs in the tissue just below the surface of the skin) with airway obstruction (difficulty in breathing) has been reported in some patients on solifenacin succinate Solifenacin succinate film-coated tablets. If angioedema occurs, solifenacin succinate Solifenacin succinate film-coated tablets should be discontinued immediately and appropriate therapy and/or measures should be taken.

Solifenacin succinate film-coated tablets may cause the following other side effects:

Very common (may affect more than 1 in 10 people)

- dry mouth

Common (may affect up to 1 in 10 people)

- blurred vision
- constipation, nausea, indigestion with symptoms such as abdominal fullness, abdominal pain, burping, nausea, and heartburn (dyspepsia), stomach discomfort

Uncommon (may affect up to 1 in 100 people)

- urinary tract infection, bladder infection
- sleepiness
- impaired sense of taste (dysgeusia)
- dry (irritated) eyes
- dry nasal passages
- reflux disease (gastro-oesophageal reflux)
- dry throat
- dry skin
- difficulty in passing urine
- tiredness
- accumulation of fluid in the lower legs (oedema)

Rare (may affect up to 1 in 1,000 people)

- lodging of a large amount of hardened stool in the large intestine (faecal impaction)
- build up of urine in the bladder due to inability to empty the bladder (urinary retention)
- dizziness, headache
- vomiting
- itching, rash

Very rare (may affect up to 1 in 10,000 people)

- hallucinations, confusion
- allergic rash

Not known (frequency cannot be estimated from the available data)

- decreased appetite, high levels of blood potassium which can cause abnormal heart rhythm
- increased pressure in the eyes
- changes in the electrical activity of the heart (ECG), irregular heartbeat (Torsade de Pointes), feeling your heartbeat, faster heartbeat
- voice disorder
- liver disorder
- muscle weakness
- renal disorder

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance

Earlsfort Terrace  
IRL - Dublin 2  
Tel: +353 1 6764971  
Fax: +353 1 6762517  
Website: [www.hpra.ie](http://www.hpra.ie)  
e-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie)

By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Solifenacin succinate film-coated tablets**

- This medicinal product does not require any special storage conditions.
- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the blister or carton after EXP. The expiry date refers to the last day of that month.
- Do not use this medicine if you notice the pack that is damaged or shows signs of tampering.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Solifenacin succinate film-coated tablets contains**

The active substance is Solifenacin succinate

Each film-coated tablet contains 5 mg solifenacin succinate, equivalent to 3.8 mg solifenacin.  
Each film-coated tablet contains 10 mg solifenacin succinate, equivalent to 7.5 mg solifenacin.

The other ingredients are:

Tablet core: Lactose monohydrate, Maize starch, Hypromellose (3 cps) (E464), Magnesium stearate (E572)

Film-coating: Hypromellose (5cps) (E464), Talc (E553b), Titanium Dioxide (E171), Macrogol 6000 (E1521), Iron oxide yellow (E172) (For 5 mg), Iron oxide red (E172) (For 10 mg)

### **What Solifenacin succinate film-coated tablets looks like and contents of the pack**

Solifenacin succinate 5 mg Film-coated Tablets: Light yellow colored, round, approximately 7.1 mm in diameter, biconvex, film coated tablets, debossed with “EG” on one side and “1” on other side.

Solifenacin succinate 10 mg Film-coated Tablets: Light pink colored, round, approximately 7.1 mm in diameter, biconvex, film coated tablets, debossed with “EG” on one side and “2” on other side.

Solifenacin succinate film-coated tablets are available in PVC/PVdC-Aluminium blister packs of 3, 5, 10, 20, 30, 50, 60, 90,100 and 200 tablets.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

#### **Marketing Authorisation Holder**

#### **United Kingdom**

Accord Healthcare Limited,  
Sage House, 319 Pinner Road,

North Harrow, Middlesex, HA1 4HF,  
United Kingdom

### Ireland

Accord Healthcare Ireland Limited  
Euro House  
Euro Business Park  
Little Island  
Cork T45 K857  
Ireland

### Manufacturer

Accord Healthcare B.V.,  
Winthortlaan 200,  
3526 KV Utrecht,  
The Netherlands

Pharmadox Healthcare Ltd.  
KW20A Kordin Industrial Park,  
Paola, PLA 3000  
Malta

Accord Healthcare Polska Sp.z o.o.,  
ul. Lutomierska 50,95-200 Pabianice, Poland

**This medicinal product is authorised in the Member States of the EEA under the following names:**

| Country | Invented Name  |
|---------|--|
| BG      | Солифенацин Акорд 5 mg/10 mg филмирани таблетки                    |
| CY      | Solifenacin Accord 5mg/10 mg Film-coated Tablets                   |
| CZ      | Solifenacin Accord 5mg/10 mg potahované tablety                    |
| LV      | Solifenacin Accord 5 mg/10 mg apvalkotās tabletes                  |
| LT      | Solifenacin Accord 5 mg/10 mg plēvele dengtos tabletēs             |
| SL      | Solifenacin Accord 5 mg/10 mg filmsko obložene tablete             |
| DK      | Solifenacinsuccinat Accord 5mg/10 mg filmovertrukne tabletter      |
| AT      | Solifenacin Accord 5 mg/10 mg Filmtabletten                        |
| DE      | Solifenacin Accord 5 mg/10 mg Filmtabletten                        |
| FI      | Solifenacin Accord 5 mg/10 mg kalvopäällysteinen tabletti          |
| IE      | Solifenacin succinate 5 mg/10 mg film-coated tablet                |
| IT      | Solifenacina Accord  |
| NL      | Solifenacinesuccinat Accord 5 mg/10 mg, filmomhulde tabletten      |
| NO      | Solifenacin Accord   |
| PL      | Soluro   |
| SE      | Solifenacin Accord 5 mg/10 mg filmdragerad tabletter               |
| UK      | Solifenacin Succinate 5 mg/10 mg Film-coated Tablets               |
| ES      | Solifenacin Accord 5 mg/10 mg comprimidos recubiertos con película |
| FR      | Solifenacin Accord 5 mg/10 mg comprimé pelliculé                   |

**This leaflet was approved in 11/2022.**