

**Package Leaflet: Information for the user**  
**Locoid® Lipocream 0.1% w/w Cream**  
Hydrocortisone butyrate

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet:**

1. What Locoid® Lipocream is and what it is used for
2. What you need to know before you use Locoid® Lipocream
3. How to use Locoid® Lipocream
4. Possible side effects
5. How to store Locoid® Lipocream
6. Contents of the pack and other information

**1. What Locoid® Lipocream is and what it is used for**

The active substance in Locoid Lipocream (hydrocortisone butyrate) is a corticosteroid which when made up as a cream is for use on the skin. This is known as topical application. Topical corticosteroids are ranked as mildly potent, moderately potent, potent and very potent; Locoid Lipocream is classed as potent. Topical corticosteroids are able to reduce the inflammation caused by a variety of skin conditions, and so allow it to get better. They can also affect the cause of the condition, which can lead to its healing.

Locoid Lipocream is indicated in adults, children and infants above 3 months of age. It is used to treat a variety of skin conditions which are not caused by micro-organisms. Among these are eczema and dermatitis (itching) of all types including atopic (childhood) eczema, photodermatitis (itching caused by exposure to light) and dermatitis caused by skin irritants and allergies.

It can also be used in the treatment of psoriasis, a condition resulting in thickening of areas of the skin. It is important that you follow your doctor's instructions particularly in psoriasis, and use it only on small areas on the scalp, hands or feet. Because Locoid Lipocream contains such a high percentage of fats and oils (see section 6 "Contents of the pack and other information"), and because these are dispersed as small droplets in the water in the cream, it is suitable for use on both dry, scaly lesions and moist, weeping lesions. The original symptoms may return if you stop using Locoid Lipocream prematurely.

**2. What you need to know before you use Locoid® Lipocream**

**Do not use Locoid Lipocream if:**

- you are **allergic** (hypersensitive) to hydrocortisone butyrate or any of the other ingredients of Locoid Lipocream. See section 6 “Contents of the pack and other information” for a list of other ingredients.
- your skin disorder is caused by an untreated **bacterial, viral (e.g. cold sores, herpes, chicken pox), fungal (e.g. athlete’s foot), yeast or parasitic (e.g. ringworm) infection**. Locoid Lipocream may worsen or hide these disorders. Your doctor will check these.
- you have skin problems associated with **syphilis (a sexually transmitted disease) or tuberculosis**.
- the **skin around your mouth is inflamed (perioral dermatitis)**. It might be the result of previous treatment with corticosteroids.
- you have **pimples or spots (acne)**.
- you have **redness of the facial skin**.

## Warnings and precautions

**Talk to your doctor before using Locoid Lipocream. This is especially important if you intend to:**

- treat facial skin, thin skin (e.g. skin of your genitals). These areas of your skin are particularly sensitive to corticosteroids. Do not apply to the eyelids in the view of the risk of glaucoma or cataract.
- use Locoid Lipocream at skin folds, under an airtight dressing or on large areas of your skin. If you are using Locoid Lipocream under bandages, it should only be on small areas for a short time, and only on the advice of your doctor.
- treat your child (see further information below under the heading “Children and adolescents”).
- use it for psoriasis. Topical corticosteroids can be effective in psoriasis in the short term. The condition may relapse or significantly worsen on stopping treatment and there is a risk of widespread pustular psoriasis. This is a condition where the psoriasis spreads and becomes very inflamed.

In order to minimise side effects Locoid Lipocream should only be used on small areas of psoriasis (e.g. scalp, hands and feet). If your doctor has prescribed Locoid Lipocream to treat psoriasis, you should let your doctor review your progress regularly as such treatment needs careful supervision.

During treatment:

Contact your doctor if you experience blurred vision or other visual disturbances.

Do not smoke or go near naked flames - risk of severe burns. Fabric (clothing, bedding, dressings, etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.

## Children and adolescents

In infants, Locoid Lipocream should not normally be used for more than seven days. Do not apply Locoid Cream under bandages or dressings in babies or children.

## Other medicines and Locoid Lipocream

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines.

**Pregnancy and breast-feeding**

You should not use Locoid Lipocream if you are pregnant or breast-feeding unless clearly necessary. There may be a small risk of harm to your baby if you use any corticosteroid cream whilst pregnant, so if your doctor does prescribe it for you knowing you are pregnant, you should not use it in large amounts for a long time.

Ask your doctor or pharmacist for advice before taking any medicine.

**Driving and using machines**

Locoid Lipocream is unlikely to affect your ability to drive or use machines.

**Locoid Lipocream contains cetostearyl alcohol, propyl parahydroxybenzoate and benzyl alcohol.**

Locoid Lipocream contains cetostearyl alcohol that may cause local skin reactions (e.g. contact dermatitis), propyl parahydroxybenzoate (E216) that may cause allergic reactions (possibly delayed) and benzyl alcohol that may cause allergic reactions and local skin reactions.

**3. How to use Locoid® Lipocream****Instructions for proper use**

Always use Locoid Lipocream exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. If your skin problem does not improve, you should go back to your doctor.

The usual dose of Locoid Lipocream is one or two times daily. Thickened areas of psoriasis on elbows and knees may be treated with Locoid Lipocream under a dressing, but only if your doctor has told you so.

Apply a thin layer of Locoid Lipocream to the affected skin.

On advice of your doctor you may have to apply Locoid Lipocream under an airtight dressing in order to obtain a better result.

Unless you are using Locoid to treat your hands, always wash your hands after using Locoid Lipocream.

**If you use more Locoid Lipocream than you should**

If you have used too much Locoid Lipocream the side effects (as described in Section 4) can occur. Contact your doctor or pharmacist immediately.

**If you forget to use Locoid Lipocream**

Do not use a double dose to make up for a forgotten dose.

**If you stop using Locoid Lipocream**

The original symptoms may return if you stop using Locoid Lipocream prematurely. If you have any further questions on the use of this product, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

A small number of people may be hypersensitive (allergic) to the constituents of Locoid Lipocream. If your skin condition seems to get worse, the skin becomes red, itchy or irritated, tell your doctor or pharmacist if this becomes troublesome. If Locoid Lipocream is used in moist skin areas (e.g. skin folds) the skin may become thin and damaged. Tell your doctor if this happens.

Locoid Lipocream may cause the following:

**Rare side effects (may affect up to 1 in 1,000 people)**

- Thinning of the skin (atrophy), inflammation of the skin (dermatitis)
- Bruising and stretch marks
- Facial spots (acne)
- Redness of the skin around the mouth (perioral dermatitis)
- Worsening of the skin condition after treatment has stopped
- Discolouration of skin.

**Very rare side effects (may affect up to 1 in 10,000 people)**

- Suppression of the release of the stress hormones.  
The signs and symptoms of stress hormone suppression include weight loss, darkening of the skin and vomiting, irritability, depression, dehydration, low blood pressure, muscle fatigue and weakness as well as muscle or joint pain and diarrhoea.

**Frequency not known** (cannot be estimated from the available data)

- Skin infection
- Allergic reaction (hypersensitivity)
- Blurred vision
- Itching (pruritus)
- Redness of the skin (erythema)
- Rash
- Application site pain.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

**Reporting side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, website: [www.hpra.ie](http://www.hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Locoid® Lipocream**

Keep this medicine out of the sight and reach of children.

Do not store Locoid Lipocream above 25°C. Do not refrigerate or freeze.

Do not use Locoid Lipocream after the expiry date which is stated on the tube or carton after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Locoid Lipocream contains**

- The active substance is hydrocortisone butyrate. Each gram of Locoid Lipocream contains 1 mg of hydrocortisone butyrate.
- The other ingredients are macrogol cetostearyl ether, cetostearyl alcohol, white soft paraffin, light liquid paraffin, anhydrous sodium citrate (E331), anhydrous citric acid (E330), propyl parahydroxybenzoate (E216), benzyl alcohol and purified water.

### **What Locoid Lipocream looks like and contents of the pack**

Locoid Lipocream is a white cream with a high percentage (about 70%) of fats and oils and is supplied in 30 or 100 gram tubes with white polyethylene screw caps.

### **Marketing Authorisation Holder and Manufacturer**

#### **Marketing authorisation holder:**

CHEPLAPHARM Arzneimittel GmbH, Ziegelhof 24, 17489 Greifswald, Germany

#### **Manufacturer:**

Temmler Italia S.r.l., Via Delle Industrie 2, 20061, Carugate (MI), Italy

**This leaflet was last revised in May 2021.**

®Registered Trade Mark