
PALFORZIA® and you

Important safety information for PALFORZIA® ▼ (defatted powder of *Arachis hypogaea L.*, semen (peanuts))

This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information.

Patient safety educational booklet



PALFORZIA® Patient Educational booklet (4-6)_IE Version 1.0_June 2022. PALF-NP-IE-0002.



Indication:

PALFORZIA contains peanut protein from defatted powder of peanut seed. It is a treatment for people who are allergic to peanuts (*Arachis hypogaea L.*).

PALFORZIA is intended for children and adolescents aged 4 through 17 years and those who become adults whilst on treatment.

PALFORZIA works in people with peanut allergy by gradually increasing the body's ability to tolerate small amounts of peanut; this is known as "desensitisation". PALFORZIA can help reduce the severity of allergic reactions after exposure to peanut.

PALFORZIA is not effective against any other nut or food allergies.

Patients who are taking PALFORZIA should continue to avoid eating peanuts.

Reporting side effects

This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects your child may get to their doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via:

https://www.hpra.ie

By reporting side effects, you can help provide more information of the safety of this medicine.

For Caregivers: About this booklet

Now that your child has been prescribed PALFORZIA, this booklet aims to give you and your child important safety information about PALFORZIA treatment. It should be read alongside the patient information leaflet contained in your child's pack of PALFORZIA.

Booklets and videos

There is also a handbook for caregivers and a series of videos for your child to watch that follow the story in this booklet, which you can access by going to:

www.medicines.ie



Note for caregivers

Summary of safety information

Please ensure that you read and understand the following information and have explained it to your child – if you have any questions, don't hesitate to ask your doctor. PALFORZIA is a medicine for treating peanut allergy and, like all medicines, it can have side effects.

This booklet contains important information about how to safely take the medicine, but before we get into that, we need to explain two potential side effects that can occur with PALFORZIA: **Anaphylaxis and Eosinophilic oesophagitis (EoE)**.

To minimise the risk of a reaction, it is important that you and your child know:

- · Symptoms to watch out for
- · When to contact your doctor or emergency services
- The important Do's and Don'ts of treatment.

Anaphylaxis is a type of allergic reaction that may affect different parts of the body at the same time. It can range from mild to severe and it can be life-threatening. Reactions can happen immediately or a few hours after coming into contact with an allergen (a substance that causes an allergy). Anaphylaxis can include a combination of the following symptoms:

- · Trouble breathing
- Throat tightness or feeling of fulness in the throat
- Trouble swallowing or speaking
- · Changes in the voice
- Dizziness or fainting or feeling of impending doom
- · Severe stomach cramps or pain, vomiting, or diarrhoea
- · Severe flushing or itching of the skin

If your child experiences any combination of these symptoms and you suspect anaphylaxis,, follow your child's emergency anaphylaxis plan, use your child's adrenaline autoinjectors following the instructions on the devices and call emergency services.

If you or your child have questions or concerns about using your child's adrenaline autoinjectors talk to your doctor who can explain how and when to use the devices.

Eosinophilic oesophagitis (EoE) is a disease where a type of white blood cells (called eosinophils) build up in the tube that connects the mouth to the stomach (oesophagus). It has been reported by some people with peanut allergy receiving oral immunotherapy.

Symptoms of eosinophilic oesophagitis can include:

- · Trouble swallowing
- · Food stuck in throat
- · Burning in chest, mouth, or throat
- Regurgitation
- · Loss of appetite

If your child experiences any of these symptoms, please contact your doctor who will carefully monitor your child for stomach problems. If they suspect EoE, the PALFORZIA treatment will be stopped.

Other possible side effects:

Throat tightness, vomiting, stomach discomfort or ache, feeling sick, allergic reactions (mild, moderate or severe), cough, sneezing, throat irritation, tingling of mouth or itching in and around the mouth, hives, itchy skin, difficulty breathing, wheezing.

Dos and Don'ts

To minimise your child's risk of side effects, please make sure your child sticks to the following Do's and Don'ts of PALFORZIA treatment.

V Dos



Take your PALFORZIA every day



Take your PALFORZIA with food



Take your
PALFORZIA at
roughly the same
time of day (with
your evening
meal is best)



Carry your adrenaline autoinjectors with you at all times

X Don'ts



DO NOT eat peanuts



DO NOT take your PALFORZIA if you feel sick



DO NOT take your PALFORZIA within 2 hours of bedtime



DO NOT swallow the capsule or breathe in the powder





DO NOT run about a lot or take a hot shower or bath right before or for 3 hours after taking your PALFORZIA


Contents


- Page 7 Section 1: Arlo and PALFORZIA
- Page 15 Section 2: Arlo takes PALFORZIA for the first time
- Page 20 Section 3: Arlo's second day taking PALFORZIA
- Page 25 Section 4: Arlo gets a tummy ache

1 Arlo and PALFORZIA









"Hello Arlo," said the doctor.

"There are some very important things to remember now that you are taking PALFORZIA."

"You must remember to never eat peanuts and to always keep your adrenaline with you. Your adrenaline autoinjectors will help you if you feel very unwell because of peanuts."


"Thank you doctor!" said Arlo.



"That's great!" said the doctor.

"Now let's talk about how you should take your PALFORZIA."







"You need to take your PALFORZIA once a day, every day" said the doctor.


"Yes doctor, I will remember to take my PALFORZIA every day," replied Arlo.



"And you need to take it with some food that you aren't allergic to. With your evening meal is best," said the doctor.

"Yes doctor, I will make sure my PALFORZIA is mixed in some food," replied Arlo.






"Now, Arlo, I need to tell you something else about your PALFORZIA," said the doctor.

"Some people feel sick after they take the medicine. Some people feel itchy. Some feel sneezy. And others feel wheezy."


"Some people find it hard to swallow. Or get a pain in their chest or a tummy ache. And some need an injection of their adrenaline autoinjector to make them feel better. Remember to always carry 2 adrenaline autoinjectors."



2

Arlo takes PALFORZIA for the first time

Day




"Because you've never had PALFORZIA before, today you need to eat it 5 times in a row during this visit," said the doctor. "It's a powder. I'll put it in your food so you can eat it. Remember to let me or your parents know if you don't feel well."



17

So Arlo ate his PALFORZIA and food, and then had a little rest. Then he ate some more, and had another rest. And then again, and again, and again.

And when all of the PALFORZIA had gone and he'd sat at the doctor's for an hour, Arlo was ready to go home.



"You've taken all of your PALFORZIA for today. Well done," said the doctor.

"But remember to come back tomorrow so I can check how you feel."

"Yes doctor, I will," replied Arlo.

"Great," said the doctor. "You will need to keep coming back to see me again and again, so I can help you take your PALFORZIA. Do you promise to do that?" "Yes, I promise!" said Arlo.



Arlo's second day taking PALFORZIA




The next day Arlo went back to see the doctor.

"Hello again Arlo," said the doctor. "How are you feeling today?"

"I feel great!" replied Arlo.

"That's good news!" said the doctor. "Today I will give you your PALFORZIA once more before you can go home. Then, from tomorrow, you'll need to take the PALFORZIA once a day at roughly the same time at home with food. Taking it with your evening meal might be best... I know how you like to eat!"







"Remember, if you've been running around or playing, you need to wait a little while until you've calmed down before you take your PALFORZIA."




"And remember that you shouldn't run around or have a hot bath or shower right before or for 3 hours after you take your PALFORZIA," said the doctor.

"One last thing Arlo. Here is a special card with your name on it. Always carry this around with you in your pocket. It tells people that you have a peanut allergy, that you are taking PALFORZIA and that you have adrenaline autoinjectors. You should show it to a grown-up if you have to go to the doctors or hospital."

"Yes I will - then people will know that peanuts make me sick," replied Arlo.

Name:		Phone:		
Address:		Date of birth:		
		Phone:		
Emergency conta	<u> </u>	Phone:		
- augency cont	ct #2:	d powder of		
		civing PALFORZIA® (defatted powder of at peanut allergy, Adverse reactions (included to peanut allergy, Adverse reactions to the severe and may require emergency tree carry with them at all times.	ing atment	
This person is P	ANUT-ALLERGIC and services (peanuts), to treat	iving PALFORZIA® (delated) at peanut allergy. Adverse reactions (includ) at peanut allergy. Adverse reactions (inc		
Arachistiyas	with this imitture see should o	carry Willi the doce of ADRENALINE		
with ADRENAL	at this person is having ANA	carry Will during the a dose of ADRENALINE PHYLAXIS, give a dose of ADRENALINE ne autoinjector, and contact EMERGENCY mptoms of ANAPHYLAXIS include: abdom motors of ANAPHYLAXIS include: abdom thoat tightness/closure, shortness of breat and weak pulse.	_{linal}	
If you suspect according to the				
SERVICES on Spain itching, h	79 immediatery. yes, rash, nausea, vomiting, the hing, dizziness, passing out, PALFOR	ne autoinjector, and over the include: abuun mptoms of ANAPHYLAXIS include: abuun mptoms of breat hroat tightness/closure, shortness of breat and weak pulse. ZIA® Patient Card_IE version 1.0_June 2022. PALF-N	P-IE-0001.	
wheezing, cou	hing, dizziriess, P	ZIA® Patient Card_IE version 1.0_2		



Section 3
Arlo's second day taking PALFORZIA

4

Arlo gets a tummy ache





Every 2 weeks, Arlo went back to the see the doctor. The doctor asked how Arlo was feeling and if everything was okay, he was given a little bit more PALFORZIA.

One time when he was taking his PALFORZIA, Arlo had a tummy ache. He told his parents and they went to the doctor to see if everything was okay.

The doctor asked: "Have you been taking you PALFORZIA once a day?"


"Yes, every day!" replied Arlo.


"And how did your PALFORZIA make you feel?" asked the doctor.

"Well," said Arlo, with a frown.
"At first I felt great. But today I have had a tummy ache. It hurt and made me cry. So I told my parents, like you asked me to."

"I'm sorry to hear that, Arlo," replied the doctor. "But I'm glad you told your parents - and me too. I will give you less PALFORZIA until I can be sure that you feel better."

The doctor gave Arlo just a little bit of PALFORZIA and asked him to come back again in 2 weeks.






29


After about 6 months, Arlo had reached the Maintenance dose of PALFORZIA.















Now he doesn't see the doctor as much but he still remembers that he has to do these important things:



 Mix his PALFORZIA into a little bit of food and take it once a day, every day



 Never eat peanuts or food with peanuts in it



Always carry
 his adrenaline
 autoinjectors in case
 he needs to use them



Contact information

Aimmune Therapeutics Ireland Ltd.
Block B
The Crescent Building
Northwood, Santry, Dublin 9
Dublin
D09 C6X8
Republic of Ireland

Phone: +353 1893 4957

Email: info.ie@aimmune.com

Medical information and safety reporting

Email: medinfo-eu@aimmune.com

Tel: +353 (0) 15827964

