

Package leaflet: Information for the user
Symbicort[®] Turbohaler[®] 100 micrograms/6 micrograms/inhalation,
inhalation powder
Budesonide/formoterol fumarate dihydrate

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Symbicort Turbohaler is and what it is used for
2. What you need to know before you use Symbicort Turbohaler
3. How to use Symbicort Turbohaler
4. Possible side effects
5. How to store Symbicort Turbohaler
6. Contents of the pack and other information

1. What Symbicort Turbohaler is and what it is used for

Symbicort Turbohaler is an inhaler that is used to treat asthma in adults, adolescents and children aged 6 years and older. It contains two different medicines: budesonide and formoterol fumarate dihydrate.

- Budesonide belongs to a group of medicines called ‘corticosteroids’. It works by reducing and preventing swelling and inflammation in your lungs.
- Formoterol fumarate dihydrate belongs to a group of medicines called ‘long-acting beta₂ adrenoceptor agonists’ or ‘bronchodilators’. It works by relaxing the muscles in your airways. This helps you to breathe more easily.

This medicine is not suitable for people with severe asthma. It can be prescribed for asthma in two different ways.

a) Some people are prescribed two asthma inhalers: Symbicort Turbohaler and a separate ‘reliever inhaler’.

- They use Symbicort Turbohaler every day. This helps to prevent asthma symptoms from happening.
- They use their ‘reliever inhaler’ when they get asthma symptoms, to make it easier to breathe again.

b) Some people are prescribed Symbicort Turbohaler as their only asthma inhaler.

- They use Symbicort Turbohaler every day. This helps to prevent asthma symptoms from happening.
- They also use Symbicort Turbohaler when they need extra doses for relief of asthma symptoms, to make it easier to breathe again. They do not need a separate inhaler for this.

2. What you need to know before you use Symbicort Turbohaler

Do not use Symbicort Turbohaler:

- if you are allergic to budesonide, formoterol or the other ingredient of this medicine (listed in section 6), which is lactose (which contains small amounts of milk protein).

Warnings and precautions

Talk to your doctor or pharmacist before using Symbicort Turbohaler if:

- You are diabetic.
- You have a lung infection.
- You have high blood pressure or you have ever had a heart problem (including an uneven heart beat, a very fast pulse, narrowing of the arteries or heart failure).
- You have problems with your thyroid or adrenal glands.
- You have low levels of potassium in your blood.
- You have severe liver problems.

Contact your doctor if you experience blurred vision or other visual disturbances.

Other medicines and Symbicort Turbohaler

Tell your doctor or pharmacist if you are using, have recently used or might use any other medicines.

In particular, tell your doctor or pharmacist if you are using any of the following medicines:

- Beta-blocker medicines (such as atenolol or propranolol for high blood pressure), including eyedrops (such as timolol for glaucoma).
- Medicines for a fast or uneven heart beat (such as quinidine).
- Medicines like digoxin, often used to treat heart failure.
- Diuretics, also known as 'water tablets' (such as furosemide). These are used to treat high blood pressure.
- Steroid medicines that you take by mouth (such as prednisolone).
- Xanthine medicines (such as theophylline or aminophylline). These are often used to treat asthma.
- Other bronchodilators (such as salbutamol).
- Tricyclic anti-depressants (such as amitriptyline) and the anti-depressant nefazodone.
- Phenothiazine medicines (such as chlorpromazine and prochlorperazine).
- Medicines called 'HIV-protease inhibitors' (such as ritonavir) to treat HIV infection.
- Medicines to treat infections (such as ketoconazole, itraconazole, voriconazole, posaconazole, clarithromycin and telithromycin).
- Medicines for Parkinson's disease (such as leva-dopa).
- Medicines for thyroid problems (such as levo-thyroxine).

If any of the above applies to you, or if you are not sure, talk to your doctor or pharmacist before using Symbicort Turbohaler.

Also tell your doctor or pharmacist if you are going to have a general anaesthetic for an operation or for dental work.

Pregnancy, breast-feeding and fertility

- If you are pregnant, or planning to get pregnant, talk to your doctor before using Symbicort Turbohaler - do not use Symbicort Turbohaler unless your doctor tells you to.

- If you get pregnant while using Symbicort Turbohaler, do not stop using Symbicort Turbohaler but talk to your doctor immediately.
- If you are breast-feeding, talk to your doctor before using Symbicort Turbohaler.

Driving and using machines

Symbicort Turbohaler has no or negligible effect on your ability to drive or to use tools or machines.

Symbicort Turbohaler contains lactose

Symbicort Turbohaler contains lactose, which is a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars, talk to your doctor before using this medicine. The amount of lactose in this medicine does not normally cause problems in people who are lactose intolerant.

The excipient lactose contains small amounts of milk proteins, which may cause an allergic reaction.

3. How to use Symbicort Turbohaler

- Always use this medicine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.
- It is important to use Symbicort Turbohaler every day, even if you have no asthma symptoms at the time.
- Your doctor will want to regularly check your asthma symptoms.

If you have been taking steroid tablets for your asthma or COPD, your doctor may reduce the number of tablets that you take, once you start to use Symbicort Turbohaler. If you have been taking oral steroid tablets for a long time, your doctor may want you to have blood tests from time to time. When reducing oral steroid tablets, you may feel generally unwell even though your chest symptoms may be improving. You might experience symptoms such as a stuffy or runny nose, weakness or joint or muscle pain and rash (eczema). If any of these symptoms bother you, or if symptoms such as headache, tiredness, nausea (feeling sick) or vomiting (being sick) occur, please contact your doctor immediately. You may need to take other medication if you develop allergic or arthritic symptoms. You should speak to your doctor if you are concerned as to whether you should continue to use Symbicort Turbohaler.

Your doctor may consider adding steroid tablets to your usual treatment during periods of stress (for example, when you have a chest infection or before an operation).

Important information about your asthma symptoms

If you feel you are getting breathless or wheezy while using Symbicort Turbohaler, you should continue to use Symbicort Turbohaler but go to see your doctor as soon as possible, as you may need additional treatment.

Contact your doctor immediately if:

- Your breathing is getting worse or you often wake up at night with asthma.
- Your chest starts to feel tight in the morning or your chest tightness lasts longer than usual.

These signs could mean that your asthma is not being properly controlled and you may need different or additional treatment immediately.

Asthma

Symbicort Turbohaler can be prescribed for asthma in two different ways. The amount of Symbicort Turbohaler to use and when to use it depends on how it has been prescribed for you.

- a) If you have been prescribed Symbicort Turbohaler and a separate reliever inhaler, read the section called 'a) Using Symbicort Turbohaler and a separate reliever inhaler'.
- b) If you have been prescribed Symbicort Turbohaler as your only inhaler, read the section called 'b) Using Symbicort Turbohaler as your only asthma inhaler'.

a) Using Symbicort Turbohaler and a separate reliever inhaler

Use your Symbicort Turbohaler every day. This helps to prevent asthma symptoms from happening.

Adults (18 years and above)

- The usual dose is 1 or 2 inhalations, twice a day.
- Your doctor may increase this to 4 inhalations, twice a day.
- If your symptoms are well controlled, your doctor may ask you to take your medicine once a day.

Adolescents (12 to 17 years)

- The usual dose is 1 or 2 inhalations, twice a day.
- If your symptoms are well controlled, your doctor may ask you to take your medicine once a day.

Children (6 to 11 years)

- The usual dose is 2 inhalations, twice a day.

Symbicort Turbohaler is not recommended to be used in children who are younger than 6 years.

Your doctor (or asthma nurse) will help you to manage your asthma. They will adjust the dose of this medicine to the lowest dose that controls your asthma. However, do not adjust the dose without talking to your doctor (or asthma nurse) first.

Use your separate 'reliever inhaler' to treat asthma symptoms when they happen.

Always keep your 'reliever inhaler' with you to use when you need it. Do not use Symbicort Turbohaler to treat asthma symptoms - use your reliever inhaler.

b) Using Symbicort Turbohaler as your only asthma inhaler

Only use Symbicort Turbohaler in this way if your doctor has told you to and if you are aged 12 or above.

Use your Symbicort Turbohaler every day. This helps to prevent asthma symptoms from happening. You can take:

- 1 inhalation in the morning **and** 1 inhalation in the evening
- or**
- 2 inhalations in the morning
- or**

- 2 inhalations in the evening.

Also use Symbicort Turbohaler as a ‘reliever inhaler’ to treat asthma symptoms when they happen.

- If you get asthma symptoms, take 1 inhalation and wait a few minutes.
- If you do not feel better, take another inhalation.
- Do not take more than 6 inhalations at a single time.

Always keep your Symbicort Turbohaler with you, so you can use it when you need it.

A total daily dose of more than 8 inhalations is not normally needed. However, your doctor may allow you to take up to 12 inhalations a day for a limited period.

If you regularly need to use 8 or more inhalations a day, make an appointment to see your doctor or nurse. They may need to change your treatment.

Do not use more than 12 inhalations in total in 24 hours.

If you are doing exercise and you get asthma symptoms, use Symbicort Turbohaler as described here. However, do not use Symbicort Turbohaler just before exercise to stop asthma symptoms from happening.

Preparing your new Symbicort Turbohaler

Before using your **new** Symbicort Turbohaler **for the first time**, you need to prepare it for use as follows:

- Unscrew the cover and lift it off. You may hear a rattling sound.
- Hold your Turbohaler upright with the red grip at the bottom.
- Turn the red grip as far as it will go in one direction. Then turn it as far as it will go in the other direction (it does not matter which way you turn it first). You should hear a click sound. It does not matter whether the click comes on the first or the second twist.
- Do this again, turning the red grip in both directions.
- Your Turbohaler is now loaded and ready for use.



How to take an inhalation

Every time you need to take an inhalation, follow the instructions below.

1. Unscrew the cover and lift it off. You may hear a rattling sound.
2. **Hold your Turbohaler upright** with the red grip at the bottom.
3. Do not hold the mouthpiece when you load your Turbohaler. To load your Turbohaler with a dose, turn the red grip as far as it will go in one direction. Then turn it as far as it will go in the other direction (it does not matter which way you turn it first). You should hear a click sound. It does not matter whether the click comes on the first or the second twist. Your Turbohaler is now loaded and ready to use. Only load your Turbohaler when you need to use it.
4. Hold your Turbohaler away from your mouth. Breathe out gently (as far as is comfortable). Do not breathe out through your Turbohaler.



5. Place the mouthpiece gently between your teeth. Close your lips. Breathe in as deeply and as hard as you can through your mouth. Do not chew or bite on the mouthpiece.
6. **Remove your Turbohaler from your mouth. Then breathe out gently.** The amount of medicine that is inhaled is very small. This means you may not be able to taste it after inhalation. If you have followed the instructions, you can still be confident that you have inhaled the dose and the medicine is now in your lungs.
7. If you are to take a second inhalation, repeat steps 2 to 6.
8. Replace the cover tightly after use.
9. Rinse your mouth with water after your daily morning and/or evening doses, and spit it out.



Do not try to remove or twist the mouthpiece. It is fixed to your Turbohaler and must not be taken off. Do not use your Turbohaler if it has been damaged or if the mouthpiece has come apart from your Turbohaler.

As with all inhalers, caregivers should ensure that children prescribed Symbicort Turbohaler use correct inhalation technique, as described above.

Cleaning your Turbohaler

Wipe the outside of the mouthpiece once a week with a dry tissue. Do not use water or liquids.

When to start using a new inhaler

- The dose indicator tells you how many doses (inhalations) are left in your Symbicort Turbohaler Inhaler, starting with either 60 or 120 doses when it is full.
- The dose indicator is marked in intervals of 10 doses. Therefore, it does not show every dose.
- When you first see a red mark at the edge of the indicator window, there are approximately 20 doses left. For the last 10 doses, the background of the dose indicator is red. When the '0' on the red background has reached the middle of the window, you must start using your new Symbicort Turbohaler Inhaler.



Note:

- The grip will still twist and 'click' even when your Symbicort Turbohaler Inhaler is empty.
- The sound that you hear as you shake your Symbicort Turbohaler Inhaler is produced by a drying agent and not the medicine. Therefore, the sound does not tell you how much medicine is left in your Symbicort Turbohaler Inhaler.
- If you load your Symbicort Turbohaler Inhaler more than once by mistake before taking your dose, you will still only receive one dose. However, the dose indicator will register all the loaded doses.

If you use more Symbicort Turbohaler than you should

It is important that you take your dose as stated on the pharmacist's label or as advised by your doctor. You should not exceed your prescribed dose without seeking medical advice.

The most common symptoms that may occur if you use more Symbicort Turbohaler than you should are trembling, headache or a rapid heart beat.

If you forget to use Symbicort Turbohaler

- If you forget to take a dose, take it as soon as you remember. However, if it is nearly time for your next dose, skip the missed dose.
- Do **not** take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If either of the following happen to you, stop using Symbicort Turbohaler and talk to your doctor immediately:

- Swelling of your face, particularly around your mouth (tongue and/or throat and/or difficulty swallowing) or hives together with difficulty breathing (angioedema) and/or sudden feeling of faintness. This may mean that you are having an allergic reaction. This happens rarely, affecting less than 1 in 1,000 people.
- Sudden acute wheezing or shortness of breath immediately after using your inhaler. **If either of these symptoms occur, stop using your Symbicort Turbohaler straightaway and use your 'reliever' inhaler. Contact your doctor immediately as you may need to have your treatment changed.** This happens very rarely, affecting less than 1 in 10,000 people.

Other possible side effects:

Common (may affect up to 1 in 10 people)

- Palpitations (awareness of your heart beating), trembling or shaking. If these effects occur, they are usually mild and usually disappear as you continue to use Symbicort Turbohaler.
- Thrush (a fungal infection) in the mouth. This is less likely if you rinse your mouth out with water after using your Symbicort Turbohaler.
- Mild sore throat, coughing and a hoarse voice.
- Headache.

Uncommon (may affect up to 1 in 100 people)

- Feeling restless, nervous or agitated.
- Disturbed sleep.
- Feeling dizzy.
- Nausea (feeling sick).
- Fast heart beat.
- Bruising of the skin.
- Muscle cramps.
- Blurred vision.

Rare (may affect up to 1 in 1,000 people)

- Rash, itching.

- Bronchospasm (tightening of the muscles in the airways which causes wheezing). If the wheezing comes on suddenly after using Symbicort Turbohaler stop using Symbicort Turbohaler and talk to your doctor immediately.
- Low levels of potassium in your blood.
- Uneven heart beat.

Very rare (may affect up to 1 in 10,000 people)

- Depression.
- Changes in behaviour, especially in children.
- Chest pain or tightness in the chest (angina pectoris).
- An increase in the amount of sugar (glucose) in your blood.
- Taste changes, such as an unpleasant taste in the mouth.
- Changes in your blood pressure.

Inhaled corticosteroids can affect the normal production of steroid hormones in your body, particularly if you use high doses for a long time. The effects include:

- changes in bone mineral density (thinning of the bones)
- cataract (clouding of the lens in the eye)
- glaucoma (increased pressure in the eye)
- a slowing of the rate of growth of children and adolescents
- an effect on the adrenal gland (a small gland next to the kidney).

These effects are much less likely to happen with inhaled corticosteroids than with corticosteroid tablets.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

Ireland

HPRA Pharmacovigilance

Website: www.hpra.ie

Malta

ADR Reporting

Website: www.medicinesauthority.gov.mt/adrportal

5. How to store Symbicort Turbohaler

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton or on the label of your inhaler after EXP. The expiry date refers to the last day of that month.
- This medicine does not require any special storage conditions.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Symbicort 100/6 Turbohaler contains

The active substances are budesonide and formoterol fumarate dihydrate. Each inhaled dose contains 80 micrograms of budesonide and 4.5 micrograms of formoterol fumarate dihydrate, this corresponds to a metered dose of 100 micrograms of budesonide and 6 micrograms of formoterol fumarate dihydrate.

The other ingredient is lactose monohydrate (which contains milk proteins).

What Symbicort 100/6 Turbohaler looks like and contents of the pack

Symbicort 100/6 Turbohaler is an inhaler containing your medicine. The inhalation powder is white in colour. Each Turbohaler contains either 60 or 120 doses and has a white body with a red turning grip. The turning grip contains a Braille code with the number 6 for identification, to differ from other AstraZeneca inhaled products.

Symbicort 100/6 Turbohaler is available in packs of 1, 2, 3, 10 or 18 Turbohalers containing 60 or 120 doses.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

The Marketing Authorisation for Symbicort 100/6 Turbohaler in Ireland and Malta is held by AstraZeneca AB, SE-151 85 Södertälje, Sweden.

Symbicort 100/6 Turbohaler is manufactured by AstraZeneca AB, Forskargatan 18, SE-151 36 Södertälje, Sweden.

This medicinal product is authorised in the Member States of the EEA under the following names:

Country	Trade name and strength
Austria	Symbicort mite Turbohaler 80 µg/4.5 µg/inhalation
Belgium	Symbicort mite Turbohaler 80 µg/4.5 µg/inhalation
Bulgaria	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Croatia	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Cyprus	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Czech Republic	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Denmark	Symbicort mite Turbuhaler 80 µg/4.5 µg/inhalation
Estonia	Symbicort Turbuhaler 80 µg/4.5 µg
Finland	Symbicort Turbuhaler mite 80 µg/4.5 µg/inhalation
France	Symbicort Turbuhaler 100 µg/6 µg/inhalation
Germany	Symbicort mite Turbohaler 80 µg/4.5 µg/inhalation
Greece	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Hungary	Symbicort mite Turbuhaler 80 µg/4.5 µg/inhalation
Iceland	Symbicort mite Turbuhaler 80 µg/4.5 µg/inhalation
Ireland	Symbicort Turbohaler 100 µg/6 µg/inhalation
Italy	Symbicort mite 80 µg/4.5 µg/inhalation
Latvia	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Lithuania	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Luxembourg	Symbicort mite Turbohaler 80 µg/4.5 µg/inhalation

Malta	Symbicort Turbohaler 100 µg/6 µg/inhalation
Netherlands	Symbicort Turbuhaler 100 µg/6 µg/inhalation
Norway	Symbicort mite Turbuhaler 80 µg/4.5 µg/inhalation
Poland	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Portugal	Symbicort Turbohaler 80 µg/4.5 µg/inhalation
Romania	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Slovakia	Symbicort Turbuhaler 100 µg/6 µg/inhalation
Slovenia	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Spain	Symbicort Turbuhaler 80 µg/4.5 µg/inhalation
Sweden	Symbicort mite Turbuhaler 80 µg/4.5 µg/inhalation
United Kingdom	Symbicort Turbohaler 100 µg/6 µg/inhalation

This leaflet was last revised in October 2023.

© AstraZeneca 2023

Symbicort and Turbohaler are trade marks of the AstraZeneca group of companies.

RSP 23 0015