

LIPITOR® 10mg, 20 mg, 40mg, 80mg Film-coated Tablets

(Atorvastatin)

PACKAGE LEAFLET: INFORMATION FOR THE USER

LIPITOR 10 mg film-coated tablets
LIPITOR 20 mg film-coated tablets
LIPITOR 40 mg film-coated tablets
LIPITOR 80 mg film-coated tablets

Atorvastatin

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist

In this leaflet:

1. What Lipitor is and what it is used for
2. Before you take Lipitor
3. How to take Lipitor
4. Possible side effects
5. How to store Lipitor
6. Further information

1. WHAT LIPITOR IS AND WHAT IT IS USED FOR

Lipitor belongs to a group of medicines known as statins, which are lipid (fat) regulating medicines.

Lipitor is used to lower lipids known as cholesterol and triglycerides in the blood when a low fat diet and life style changes on their own have failed. If you are at an increased risk of heart disease, Lipitor can also be used to reduce such risk even if your cholesterol levels are normal. You should maintain a standard cholesterol lowering diet during treatment.

2. BEFORE YOU TAKE LIPITOR

Do not take Lipitor

- if you are hypersensitive (allergic) to Lipitor or to any similar medicines used to lower blood lipids or to any of the other ingredients of the medicine – see Section 6 for details.
- if you have or have ever had a disease which affects the liver
- if you have had any unexplained abnormal blood tests for liver function
- if you are a woman able to have children and not using reliable contraception
- if you are pregnant or trying to become pregnant
- if you are breast-feeding.

Take special care with Lipitor

The following are reasons why Lipitor may not be suitable for you:

- if you have had a previous stroke with bleeding into the brain, or have small pockets of fluid in the brain from previous strokes
- if you have kidney problems
- if you have an under-active thyroid gland (hypothyroidism)
- if you have had repeated or unexplained muscle aches or pains, a personal history or family history of muscle problems
- if you have had previous muscular problems during treatment with other lipid-lowering medicines (e.g. other '-statin' or '-fibrate' medicines)
- if you regularly drink a large amount of alcohol
- if you have a history of liver disease
- if you are older than 70 years.

Check with your doctor or pharmacist before taking Lipitor

- if you have severe respiratory failure.
- If any of these apply to you, your doctor will need to carry out a blood test before and possibly during your Lipitor treatment to predict your risk of muscle related side effects. The risk of muscle related side effects e.g. rhabdomyolysis is known to increase when certain medicines are taken at the same time (see Section 2 "Taking other medicines").

Taking other medicines

There are some medicines that may change the effect of Lipitor or their effect may be changed by Lipitor. This type of interaction could make one or both of the medicines less effective. Alternatively it could increase the risk or severity of side-effects, including the important muscle wasting condition known as rhabdomyolysis described in Section 4:

- Medicines used to alter the way your immune system works, e.g. ciclosporin
- Certain antibiotics or antifungal medicines, e.g. erythromycin, clarithromycin, telithromycin, ketoconazole, itraconazole, voriconazole, fluconazole, posaconazole, rifampin, fusidic acid
- Other medicines to regulate lipid levels, e.g. gemfibrozil, other fibrates, colestipol

- Some calcium channel blockers used for angina or high blood pressure, e.g. amlodipine, diltiazem.; medicines to regulate your heart rhythm e.g. digoxin, verapamil, amiodarone
- Medicines used in the treatment of HIV e.g. ritonavir, lopinavir, atazanavir, indinavir, darunavir, etc.
- Other medicines known to interact with Lipitor include ezetimibe (which lowers cholesterol), warfarin (which reduces blood clotting), oral contraceptives, stiripentol (an anti-convulsant for epilepsy), cimetidine (used for heartburn and peptic ulcers), phenazone (a painkiller) and antacids (indigestion products containing aluminium or magnesium)
- Medicines obtained without a prescription: St John's Wort

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Taking Lipitor with food and drink

See Section 3 for instructions on how to take Lipitor. Please note the following:

Grapefruit juice

Do not take more than one or two small glasses of grapefruit juice per day because large quantities of grapefruit juice can change the effects of Lipitor.

Alcohol

Avoid drinking too much alcohol while taking this medicine. See Section 2 "Take special care with Lipitor" for details

Pregnancy and breast-feeding

Do not take Lipitor if you are pregnant, or if you are trying to become pregnant.

Do not take Lipitor if you are able to become pregnant unless you use reliable contraceptive measures.

Do not take Lipitor if you are breast-feeding.

The safety of Lipitor during pregnancy and breast-feeding has not yet been proven. Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Normally this medicine does not affect your ability to drive or operate machines. However, do not drive if this medicine affects your ability to drive. Do not use any tools or machines if your ability to use them is affected by this medicine.

Important information about some of the ingredients of Lipitor

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. HOW TO TAKE LIPITOR

Before starting treatment, your doctor will place you on a low-cholesterol diet, which you should maintain also during therapy with Lipitor.

The usual starting dose of Lipitor is 10 mg once a day in adults and children aged 10 years or older. This may be increased if necessary by your doctor until you are taking the amount you need. Your doctor will adapt the dose at intervals of 4 weeks or more. The maximum dose of Lipitor is 80 mg once daily for adults and 20 mg once daily for children.

Lipitor tablets should be swallowed whole with a drink of water, and can be taken at any time of day, with or without food. However, try to take your tablet at the same time every day.

Always take Lipitor exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

The duration of treatment with Lipitor is determined by your doctor.

Please ask your doctor if you think that the effect of Lipitor is too strong or too weak.

If you take more Lipitor than you should

If you accidentally take too many Lipitor tablets (more than your usual daily dose), contact your doctor or nearest hospital for advice.

If you forget to take Lipitor

If you forget to take a dose, just take your next scheduled dose at the correct time. Do not take a double dose to make up for a forgotten dose.

If you stop taking Lipitor

If you have any further questions on the use of this medicine or wish to stop your treatment, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Lipitor can cause side effects, although not everybody gets them.

If you experience any of the following serious side effects, stop taking your tablets and tell your doctor immediately or go to the nearest hospital accident

and emergency department.

Rare: affects 1 to 10 users in 10,000:

- Serious allergic reaction which causes swelling of the face, tongue and throat that can cause great difficulty in breathing.
- Serious illness with severe peeling and swelling of the skin, blistering of the skin, mouth, eyes genitals and fever. Skin rash with pink-red blotches especially on palms of hands or soles of feet which may blister.
- Muscle weakness, tenderness or pain and particularly, if at the same time, you feel unwell or have a high temperature it may be caused by an abnormal muscle breakdown which can be life-threatening and lead to kidney problems.

Very rare: affect less than 1 user in 10,000:

- If you experience problems with unexpected or unusual bleeding or bruising, this may be suggestive of a liver complaint. You should consult your doctor as soon as possible.

Other possible side effects with Lipitor:

Common side effects (affects 1 to 10 users in 100) include:

- inflammation of the nasal passages, pain in the throat, nose bleed
- allergic reactions
- increases in blood sugar levels (if you have diabetes continue careful monitoring of your blood sugar levels), increase in blood creatine kinase
- headache
- nausea, constipation, wind, indigestion, diarrhoea
- joint pain, muscle pain and back pain
- blood test results that show your liver function can become abnormal

Uncommon side effects (affects 1 to 10 users in 1000) include:

- anorexia (loss of appetite), weight gain, decreases in blood sugar levels (if you have diabetes you should continue careful monitoring of your blood sugar levels)
- having nightmares, insomnia
- dizziness, numbness or tingling in the fingers and toes, reductions of sensation to pain or touch, change in sense of taste, loss of memory
- blurred vision
- ringing in the ears and/or head
- vomiting, belching, abdominal pain upper and lower, pancreatitis (inflammation of the pancreas leading to stomach pain)
- hepatitis (liver inflammation)
- rash, skin rash and itching, hives, hair loss
- neck pain, muscle fatigue
- fatigue, feeling unwell, weakness, chest pain, swelling especially in the ankles (oedema), raised temperature
- urine tests that are positive for white blood cells

Rare side effects (affects 1 to 10 users in 10,000) include:

- visual disturbance
- unexpected bleeding or bruising
- cholestasis (yellowing of the skin and whites of the eyes)
- tendon injury

Very rare side effects (affects less than 1 user in 10,000) include:

- an allergic reaction - symptoms may include sudden wheezing and chest pain or tightness, swelling of the eyelids, face, lips, mouth, tongue or throat, difficulty breathing, collapse
- hearing loss
- gynecomastia (breast enlargement in men and women).

Possible side effects reported with some statins (medicines of the same type):

- Sexual difficulties
- Depression
- Breathing problems including persistent cough and/or shortness of breath or fever

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

5. HOW TO STORE LIPITOR

Keep out of the reach and sight of children.

This medicine does not require any special storage conditions.

Do not use Lipitor after the expiry date which is stated on the blister and outer packaging after {EXP}. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Lipitor contains

The active substance is atorvastatin.

- Each Lipitor 10 mg Film-Coated Tablets contains 10 mg atorvastatin (as atorvastatin calcium trihydrate).
- Each Lipitor 20 mg Film-Coated Tablets contains 20 mg atorvastatin (as atorvastatin calcium trihydrate).
- Each Lipitor 40 mg Film-Coated Tablets contains 40 mg atorvastatin (as atorvastatin calcium trihydrate).
- Each Lipitor 80 mg Film-Coated Tablets contains 80 mg atorvastatin (as atorvastatin calcium trihydrate).

The other ingredients of Lipitor are:

calcium carbonate, microcrystalline cellulose, lactose monohydrate, croscarmellose sodium, polysorbate 80 (E433), hydroxypropyl cellulose (E463) and magnesium stearate (E572).

The coating of Lipitor 10mg, 20mg, 40mg, and 80mg sourced from the UK contains hypromellose, macrogol 8000, titanium dioxide (E171), talc, simeticone emulsion (containing simeticone, stearate emulsifiers (polyethylene glycol sorbitan tristearate, polyethoxylate stearate, glycerides) thickeners (methylcellulose, xanthan gum) benzoic acid, sorbic acid, water)

The coating of Lipitor 10mg, 20mg, 40mg, and 80mg sourced from Spain contains hypromellose, macrogol 8000, titanium dioxide (E171), talc, simeticone emulsion (containing simeticone, stearate emulsifiers, thickeners, benzoic acid, sorbic acid)

The coating of Lipitor 10mg, 20 mg and 40 mg sourced from Italy contains: hypromellose, macrogol 8000, titanium dioxide (E171), talc, simeticone, stearate emulsifiers, sorbic acid and candelilla wax.

What Lipitor looks like and contents of the pack

Lipitor 10 mg Film-Coated Tablets are white, round, debossed '10' on one side and 'ATV' on the other side.

Lipitor 20 mg Film-Coated Tablets are white, round, debossed '20' on one side and 'ATV' on the other side.

Lipitor 40 mg Film-Coated Tablets are white, round, debossed '40' on one side and 'ATV' on the other side.

Lipitor 80 mg Film-Coated Tablets are white, round, debossed '80' on one side and 'ATV' on the other side.

The 10mg, 20mg, 40mg strengths of Lipitor are supplied in blister packs of 28 or 30 tablets in overlabelled cartons.

The 80mg strength is supplied in blister packs of 28 tablets in overlabelled cartons

This medicine is available as 10 mg, 20 mg, 40 mg and 80 mg film-coated tablets.

Manufacturer and Authorisation Holder

Lipitor tablets are manufactured by Pfizer Manufacturing Deutschland GmbH, Mooswaldallee 1, D-79090 Freiburg, Germany or Godecke GmbH, Mooswaldallee 1, D-79090 Freiburg, Germany.

Procured from within the EU by the parallel product authorisation holder: Primecrown 2010 Ltd, Northolt Trading Estate, Belvue Road, Northolt, Middlesex UB5 5QS UK and repackaged by Primecrown Ltd, Northolt Trading Estate, Belvue Road, Northolt, Middlesex UB5 5QS UK

Distributed by Merit Pharmaceuticals, Swords, Co. Dublin

This leaflet was revised Nov 2011

Lipitor is a registered trademark of Pfizer Pharmaceuticals Pottery Road, Dun Laoghaire, Co. Dublin, Ireland

This medicinal product is authorised in the Member States of the EEA under the following names:

Austria, Bulgaria, Czech Republic, Estonia, Germany, Hungary, Latvia, Lithuania, Poland, Romania, Slovakia, Slovenia	Sortis
Belgium, Cyprus, Finland, Greece, Ireland, Italy, Luxembourg, Malta, Netherlands, Norway, Sweden, UK	Lipitor
Denmark, Greece, Iceland, Portugal, Spain	Zarator
Finland	Orbeos
France	Tahor
Germany	Atorvastatin Pfizer, Liprimar
Greece	Edovin
Hungary	Obradon
Italy	Torvast, Totalip, Xarator
Portugal	Atorvastatina Parke-Davis, Texzor
Spain	Cardyl, Atorvastatina Nostrum, Atorvastatina Pharmacia, Prevencor