

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Cipramil® 20mg Film-coated Tablets Citalopram (as hydrobromide)

Your medicine is available using the name Cipramil 20mg Film-coated Tablets but will be referred to as Cipramil throughout this leaflet.

#### Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

#### What is in this leaflet

- What Cipramil is and what it is used for
- What you need to know before you take Cipramil
- How to take Cipramil
- Possible side effects
- How to store Cipramil
- Contents of the pack and other information

### 1. What Cipramil is and what it is used for

#### How does Cipramil work

Cipramil belongs to a group of antidepressants called Selective Serotonin Reuptake Inhibitors (SSRIs). These medicines act on the serotonin-system in the brain by increasing the serotonin level. Disturbances in the serotonin-system are considered an important factor in the development of depression and related diseases.

#### What is Cipramil used for

Cipramil contains citalopram and is used for the treatment of depression and when you feel better, to help prevent these symptoms recurring.

Further, Cipramil is used for long-term treatment to prevent the occurrence of new depressive episodes in patients who have recurrent depression.

Cipramil is also beneficial in relieving symptoms in patients prone to panic attacks.

Ask your doctor if you have any questions about why Cipramil has been prescribed for you.

### 2. What you need to know before you take Cipramil

#### Do not take Cipramil:

- if you are allergic to citalopram or any of the other ingredients of this medicine (listed in section 6)
- if you take other medicines which belong to a group called monoamine oxidase inhibitors (MAOIs). MAOIs include medicines such as phenelzine, iproniazid, isocarboxazid, nialamide, tranylcypromine, selegiline (used in the treatment of Parkinson's disease), moclobemide (used in the treatment of depression) and linezolid (an antibiotic)
- if you are born with or have had an episode of abnormal heart rhythm (seen at ECG; an examination to evaluate how the heart is functioning)
- if you take medicines for heart rhythm problems or that may affect the heart's rhythm (see "Other medicines and Cipramil")

Even if you have finished taking MAOIs, you will need to wait 2 weeks before you start getting your Cipramil treatment.

After stopping Cipramil you must allow 1 week before taking any MAOI.

#### Warnings and precautions

Please tell your doctor if you have any other condition or illness, as your doctor may need to take this into consideration. In particular, tell your doctor:

- if you have episodes of mania or panic disorder
- if you suffer from impaired liver or kidney function. Your doctor may need to adjust your dosage
- if you have diabetes. Treatment with Cipramil may alter glycaemic control. Insulin and/or oral hypoglycaemic dosage may need to be adjusted.
- if you have epilepsy. Treatment with Cipramil should be stopped if seizures occur or if there is an increase in the seizure frequency (see also section 4 "Possible side effects")
- if you have some kind of bleeding disorder
- if you have a decreased level of sodium in the blood
- if you are receiving electroconvulsive treatment
- if you suffer or have suffered from heart problems or have recently had a heart attack
- if you have a low resting heart-rate and/or you know that you may have salt depletion as a result of prolonged severe diarrhoea and vomiting (being sick) or usage of diuretics (water tablets)
- if you experience a fast or irregular heartbeat, fainting, collapse or dizziness on standing up which may indicate abnormal functioning of the heart rate
- if you have a problem with dilatation of the pupil of the eye (mydriasis) or if you have or have previously had eye problems, Such as certain kinds of glaucoma (increased pressure in the eye).

Please consult your doctor, even if these statements were applicable to you at any time in the past.

#### Please note

Some patients with manic-depressive illness may enter into a manic phase. This is characterized by unusual and rapidly changing ideas, inappropriate happiness and excessive physical activity. If you experience this, contact your doctor.

Symptoms such as restlessness or difficulty to sit or stand still (akathisia) can also occur during the first weeks of the treatment. Tell your doctor immediately if you experience these symptoms.

#### Thoughts of suicide and worsening of your depression or anxiety disorder

If you are depressed and/or have anxiety disorders you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this:

- if you have previously had thoughts about killing or harming yourself.
- if you are a **young adult**. Information from clinical trials has shown an increased risk of suicidal behaviour in young adults (less than 25 years old) with psychiatric conditions who were treated with an antidepressant

If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away. You may find it helpful to tell a relative or close friend** that you are depressed or have an anxiety disorder, and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

#### Children and adolescents

Cipramil should normally not be used for children and adolescents under 18 years. Also, you should know that patients under 18 have an increased risk of side-effects such as suicide attempts, suicidal thoughts and hostility (predominately aggression, oppositional behaviour and anger) when they take this class of medicines. Despite this, your doctor may prescribe Cipramil for patients under 18 because he/she decides that this is in their best interest. If your doctor has prescribed Cipramil for a patient under 18 and you want to discuss this, please go back to your doctor. You should inform your doctor if any symptoms listed above develop or worsen when patients under 18 are taking Cipramil. Also, the long term safety effects concerning growth, maturation and cognitive and behavioural development of Cipramil in this age group have not yet been demonstrated.

#### Special information relating to your disease

As with other medicines used to treat depression or related diseases, the improvement is not achieved immediately. After the start of Cipramil treatment it may take several weeks before you experience any improvement. In the treatment of panic disorder it usually takes 2-4 weeks before any improvement is seen.

In the beginning of the treatment certain patients may experience increased anxiety, which will disappear during the continued treatment. Therefore, it is very important that you follow exactly your doctor's orders and do not stop the treatment or change the dose without consulting your doctor.

Occasionally, the symptoms of depression or panic disorder may include thoughts of suicide or self-harm. It is possible that these symptoms continue or get worse until the full antidepressant effect of the medicine becomes apparent. This is more likely to occur if you are a young adult, i.e. under 25 years of age and you have not used antidepressive medicines before.

Sometimes you may be unaware of the above-mentioned symptoms and therefore you may find it helpful to ask a friend or relative to help you to observe the possible signs of change in your behaviour.

Tell your doctor immediately or contact the nearest hospital if you have distressing thoughts or experiences or if any of the above-mentioned symptoms occurs during the treatment.

#### Other medicines and Cipramil

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Some medicinal products may affect the action of another and this can sometimes cause serious adverse reactions.

Tell your doctor if you are taking any of the following medicines:

- "Non-selective monoamine oxidase inhibitors (MAOIs)", containing phenelzine, iproniazid, isocarboxazid, nialamide, and tranylcypromine as active substance. If you have taken any of these medicines you will need to wait 14 days before you start taking Cipramil. After stopping Cipramil you must allow 7 days before taking any of these medicines.
- "Reversible, selective MAO-A inhibitors", containing moclobemide (used to treat depression)
- The antibiotic linezolid
- Lithium (used in the prophylaxis and treatment of manic-depressive disorder) and tryptophan.
- Imipramine and desipramine (both used to treat depression)
- "Irreversible MAO-B inhibitors", containing selegiline (used to treat Parkinson's disease); these increase the risk of side effects. The dose of selegiline must not exceed 10mg per day.
- Metoprolol (used for high blood pressure and/or heart disease); the blood levels of metoprolol are increased. Dose adjustment may be required.
- Sumatriptan and similar medicines (used to treat migraine) and tramadol (used against severe pain); these increase the risk of side effects; if you get any unusual symptoms when using this combination you should see your doctor.
- Cimetidine, omeprazole and esomeprazole (used to treat stomach ulcers) and fluvoxamine (antidepressant), blood levels of Cipramil may be increased and dose adjustment may be required.
- Drugs known to affect the platelet function (e.g. some antipsychotic drugs, acetylsalicylic acid (used as pain killers), non-steroidal anti-inflammatory drugs (used for arthritis)); slightly increased risk of bleeding abnormalities.
- St John's Wort (*Hypericum perforatum*) (a herbal remedy used to treat depression) - concomitant intake with Cipramil may increase the risk of side effects.
- Mefloquin (used to treat malaria), bupropion (used to treat depression) and tramadol (used to treat severe pain) due to a possible risk of a lowered threshold for seizures
- Neuroleptics (medicines to treat schizophrenia, psychosis) and antidepressants (SSRIs) due to a possible risk of a lowered threshold for seizures
- Medicines that decrease blood levels of potassium or magnesium as these conditions increase the risk of life-threatening heart rhythm disorder

Do not take Cipramil if you take medicines for heart rhythm problems or medicines that may affect the heart's rhythm, e.g. Class IA and III antiarrhythmics, antipsychotics (e.g. fentiazine derivatives, pimozide, haloperidol), tricyclic antidepressants, certain antimicrobial agents (e.g. sparfloxacin, moxifloxacin, erythromycin IV, pentamidine, anti-malarial treatment particularly halofantrine), certain antihistamines (astemizole, mizolastine). If you have any further questions about this you should speak to your doctor.

#### Cipramil with food, drink and alcohol

Cipramil can be taken with or without food (see section 3 "How to take Cipramil").

Cipramil has been shown not to increase the effects of alcohol. Nevertheless, it is recommended not to drink alcohol during treatment with Cipramil.

#### Pregnancy, breast-feeding and fertility

Inform your doctor if you are pregnant or planning to become pregnant. Pregnant women should not usually take Cipramil nor should mothers breast-feed their babies while taking this medicine, unless you and your doctor have discussed the risks and benefits involved.

If you take Cipramil during the last 3 months of your pregnancy and until the date of birth you should be aware that the following effects may be seen in your newborn: trouble with breathing, blue-ish skin, fits, body temperature changes, feeding difficulties, vomiting, low blood sugar, stiff or floppy muscles, vivid reflexes, tremor, jitteriness, irritability, lethargy, constant crying, sleepiness and sleeping difficulties. If your newborn baby has any of these symptoms, please contact your doctor immediately.

Make sure your midwife and/or doctor know you are on Cipramil. When taken during pregnancy, particularly in the last 3 months of pregnancy, medicines like Cipramil may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the newborn (PPHN), making the baby breathe faster and appear blue-ish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby you should contact your midwife and/or doctor immediately.

Citalopram has been shown to reduce the quality of sperm in animal studies. Theoretically, this could affect fertility, but impact on human fertility has not been observed as yet.

#### Driving and using machines

Cipramil generally does not cause drowsiness; however, if you feel dizzy or sleepy when you start to take this medicine, do not drive or work any tools or machinery until these effects wear off.

#### Important information about some of the ingredients of Cipramil

This medicinal product contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### 3. How to take Cipramil

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

#### Adults

##### Depression

The usual dose is 20mg daily. The dose may be increased by your doctor to a maximum of 40mg daily.

##### Panic disorder

The starting dose is 10mg daily for the first week before increasing the dose to 20-30mg daily. The dose may be increased by your doctor to a maximum of 40mg daily.

#### Elderly patients (above 65 years of age)

The starting dose should be decreased to half of the recommended dose, e.g. 10-20mg daily. Elderly patients should not usually receive more than 20mg daily.

#### Patients with special risks

Patients with liver complaints should not receive more than 20mg daily.

#### Children and adolescents (<18 years)

Cipramil should not be given to children or adolescents. For further information please see section 2 "What you need to know before you take Cipramil".

#### How and when to take Cipramil

Cipramil is taken every day as a single daily dose. Cipramil can be taken any time of the day with or without food. Swallow the tablets with a drink of water. Do not chew them (they have a bitter taste).

#### Duration of treatment

Like other medicines for depression and panic disorder these tablets may take a few weeks before you feel any improvement. Continue to take Cipramil even if it takes some time before you feel any improvement in your condition.

Never change the dose of the medicine without talking to your doctor first.

The duration of treatment is individual, usually at least 6 months. Continue to take the tablets for as long as your doctor recommends. Do not stop taking them even if you begin to feel better, unless you are told to do so by your doctor. The underlying illness may persist for a long time and if you stop your treatment too soon your symptoms may return.

Patients who have recurrent depression benefit from continued treatment, sometimes for several years, to prevent the occurrence of new depressive episodes.

#### If you take more Cipramil than you should

If you think that you or anyone else may have taken too much Cipramil contact your doctor or nearest hospital emergency department immediately. Do this even if there are no signs of discomfort or poisoning. Take the Cipramil box/container with you if you go to a doctor or hospital.

Some of the signs of an overdose could be life-threatening irregular heart beat, convulsion, change in heart rhythm, drowsiness, coma, vomiting, tremor, decreased blood pressure, increased blood pressure, nausea (feeling sick), serotonin syndrome (see section 4), agitation, dizziness, dilated pupils of the eye, sweating, blue-ish skin, hyperventilation.

#### If you forget to take Cipramil

Do not take a double dose to make up for a forgotten dose. If you forget to take a dose, and you remember before you go to bed, take it straight away. Carry on as usual the next day. If you only remember during the night, or the next day, leave out the missed dose and carry on as usual.

### **If you stop taking Cipramil**

Do not stop taking Cipramil until your doctor tells you to do so. When you have completed your course of treatment, it is generally advised that the dose of Cipramil is gradually reduced over a number of weeks.

If you stop taking Cipramil too quickly, you may experience discontinuation symptoms. Most people find that the symptoms are mild and go away on their own within two weeks. However, in some patients they may be severe in intensity or they may be prolonged (2-3 months or more). If you get severe discontinuation symptoms when you stop taking Cipramil, please contact your doctor.

Discontinuation symptoms include: dizziness, feelings like pins and needles, sleep disturbances (vivid dreams, nightmares, inability to sleep), feeling anxious, headaches, feeling sick (nausea), vomiting, sweating, feeling restless or agitated, tremor, feeling confused or disorientated, feeling emotional or irritable, diarrhoea (loose stools), visual disturbances, fluttering or pounding heartbeat (palpitations). When you have completed your course of treatment it is therefore advised that the dose of Cipramil is gradually reduced over a couple of weeks rather than stopped abruptly.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

***If you experience any of the following symptoms contact your doctor or go to the hospital straight away:***

#### Rare (may affect up to 1 in 1,000 people):

- Hyponatraemia: abnormally low blood levels of sodium which can cause tiredness, confusion, and muscle twitching

#### Not known (frequency cannot be estimated from the available data):

- High fever, agitation, confusion, trembling and abrupt contractions of muscles; these may be signs of a rare condition called serotonin syndrome which has been reported with the combined use of antidepressants.
- Swelling of skin, tongue, lips, or face, feeling dizzy or having difficulties breathing or swallowing (serious allergic reaction)
- Unusual bleeds, including gastrointestinal bleeds
- Fast, irregular heart beat, fainting which could be symptoms of a life-threatening condition known as Torsades de Pointes

The majority of the side effects listed below are mild. Please be aware that many of the effects may also be symptoms of your illness and therefore will improve when you start to get better.

#### Very common (may affect more than 1 in 10 people):

Sleepiness  
Difficulty in sleeping  
Increased sweating  
Dry mouth (as dry mouth may increase the risk of tooth decay it is advisable to brush your teeth more often)  
Nausea (feeling sick)  
Headache

#### Common (may affect up to 1 in 10 people):

Decreased appetite  
Agitation  
Decreased sexual drive  
Anxiety  
Nervousness  
Confusional state  
Abnormal dreams  
Tremor  
Tingling or numbness in the hands or feet  
Dizziness  
Disturbance in attention  
Ringing in the ears (tinnitus)  
Yawning  
Diarrhoea  
Vomiting  
Constipation  
Itching  
Pain in muscle and joints  
Men may experience problems with ejaculation and erection  
Women may experience failure to achieve an orgasm  
Fatigue  
Fever  
Prickling of the skin  
Decreased weight

#### Uncommon (may affect up to 1 in 100 people):

Purpura: red or purple patches/spots (smaller than 1cm) of the skin caused by bleeding underneath  
Increased appetite  
Aggression  
Depersonalisation (feeling detached from yourself)  
Hallucination  
Mania (feeling excessively elated, impulsive, irritable, or irrational)  
Fainting  
Enlarged pupils  
Fast heart beat  
Slow heart beat  
Nettle rash  
Loss of hair  
Rash  
Light sensitivity  
Difficulties urinating  
Excessive menstrual bleeding  
Swelling of the arms or legs  
Increased weight

#### Rare (may affect up to 1 in 1,000 people):

Convulsions  
Involuntary movements  
Taste disturbance  
Bleeding  
Hepatitis

#### Not known (frequency cannot be estimated from the available data):

Thoughts of harming yourself or thoughts of killing yourself, see also section "Warnings and precautions"  
Reduction in blood platelets, which increases risk of bleeding or bruising  
Hypersensitivity (rash)  
Disruption to the hormones affecting urine production  
Hypokalaemia: low blood levels of potassium which can cause muscle weakness, twitching or abnormal heart rhythm  
Panic attack  
Grinding one's teeth  
Restlessness  
Unusual muscle movements or stiffness  
Akathisia (involuntary movements of the muscles)  
Visual disturbance  
Low blood pressure  
Nosebleed  
Ecchymosis: blue or purplish patches (bigger than 1cm) of the skin caused by bleeding underneath  
Sudden swelling of skin or mucosa  
Painful erections  
Flow of milk in men and in women that are not nursing  
Irregular menstrual period  
Abnormal liver function test  
An increased risk of bone fractures has been observed in patients taking these types of medicines  
Alteration of the heart rhythm (called "prolongation of QT interval", seen on ECG, a trace of the electrical activity of the heart).

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine. Reports may be made by following the links to the online reporting option accessible from the IMB homepage, or by completing the downloadable report form also accessible from the IMB website, which may be completed manually and submitted to the IMB via freepost, to the following address:

FREEPOST  
Pharmacovigilance Section  
Irish Medicines Board  
Kevin O'Malley House  
Earlsfort Centre Earlsfort Terrace  
Dublin 2  
Tel: +353 1 6764971  
Fax: +353 1 6762517  
Website: [www.imb.ie](http://www.imb.ie)  
e-mail: [imbpharmacovigilance@imb.ie](mailto:imbpharmacovigilance@imb.ie)

### **5. How to store Cipramil**

Keep out of the sight and reach of children  
Do not store above 25°C.  
Do not use Cipramil after the expiry date (EXP) which is printed on the carton and blister label.  
If your doctor tells you to stop taking your medicine you can return any left-over medicine to the pharmacist (for safe disposal), unless the doctor tells you to keep it at home.  
If your tablets appear to be discoloured, damaged or show any other signs of deterioration, please return them to your pharmacist who will advise you.  
Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

### **6. Contents of the pack and other information**

#### **What Cipramil contains**

Each film-coated Cipramil tablet contains 20mg citalopram (as hydrobromide) as the active ingredient.

The other ingredients in the tablets are: maize starch, lactose monohydrate, copovidone, magnesium stearate, glycerol, microcrystalline cellulose, croscarmellose sodium, hypromellose, macrogol and titanium dioxide (E171).

#### **What Cipramil looks like and contents of the pack**

Cipramil are white, oval tablets, deeply scored on one face with the letters 'C' and 'N' on either side of the score and plain on the reverse.

Cipramil is available in blister packs containing 28 tablets.

#### **Manufacturer**

Cipramil is manufactured by: H. Lundbeck A/S, Ottiliavej 9, DK-2500 Copenhagen, Denmark.

Procured from within the EU by the Parallel Product Authorisation holder:  
Imbat Ltd, Unit L2, North Ring Business Park, Santry, Dublin 9.

Repackaged by: Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, South Yorkshire, DN3 1QR, UK.

Distributed by: Eurodrug Ltd, Santry, Dublin 9.

PPA No: 1151/49/1

Leaflet revision and issue date (ref): 15.01.14

Cipramil® is a registered trademark of H. Lundbeck A/S.

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Cipramil® 20mg Film-coated Tablets Citalopram (as hydrobromide)

Your medicine is available using the name Cipramil 20mg Film-coated Tablets but will be referred to as Cipramil throughout this leaflet.

#### Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

#### What is in this leaflet

- What Cipramil is and what it is used for
- What you need to know before you take Cipramil
- How to take Cipramil
- Possible side effects
- How to store Cipramil
- Contents of the pack and other information

### 1. What Cipramil is and what it is used for

#### How does Cipramil work

Cipramil belongs to a group of antidepressants called Selective Serotonin Reuptake Inhibitors (SSRIs). These medicines act on the serotonin-system in the brain by increasing the serotonin level. Disturbances in the serotonin-system are considered an important factor in the development of depression and related diseases.

#### What is Cipramil used for

Cipramil contains citalopram and is used for the treatment of depression and when you feel better, to help prevent these symptoms recurring.

Further, Cipramil is used for long-term treatment to prevent the occurrence of new depressive episodes in patients who have recurrent depression.

Cipramil is also beneficial in relieving symptoms in patients prone to panic attacks.

Ask your doctor if you have any questions about why Cipramil has been prescribed for you.

### 2. What you need to know before you take Cipramil

#### Do not take Cipramil:

- if you are allergic to citalopram or any of the other ingredients of this medicine (listed in section 6)
- if you take other medicines which belong to a group called monoamine oxidase inhibitors (MAOIs). MAOIs include medicines such as phenelzine, iproniazid, isocarboxazid, nialamide, tranylcypromine, selegiline (used in the treatment of Parkinson's disease), moclobemide (used in the treatment of depression) and linezolid (an antibiotic)
- if you are born with or have had an episode of abnormal heart rhythm (seen at ECG; an examination to evaluate how the heart is functioning)
- if you take medicines for heart rhythm problems or that may affect the heart's rhythm (see "Other medicines and Cipramil")

Even if you have finished taking MAOIs, you will need to wait 2 weeks before you start getting your Cipramil treatment.

After stopping Cipramil you must allow 1 week before taking any MAOI.

#### Warnings and precautions

Please tell your doctor if you have any other condition or illness, as your doctor may need to take this into consideration. In particular, tell your doctor:

- if you have episodes of mania or panic disorder
- if you suffer from impaired liver or kidney function. Your doctor may need to adjust your dosage
- if you have diabetes. Treatment with Cipramil may alter glycaemic control. Insulin and/or oral hypoglycaemic dosage may need to be adjusted.
- if you have epilepsy. Treatment with Cipramil should be stopped if seizures occur or if there is an increase in the seizure frequency (see also section 4 "Possible side effects")
- if you have some kind of bleeding disorder
- if you have a decreased level of sodium in the blood
- if you are receiving electroconvulsive treatment
- if you suffer or have suffered from heart problems or have recently had a heart attack
- if you have a low resting heart-rate and/or you know that you may have salt depletion as a result of prolonged severe diarrhoea and vomiting (being sick) or usage of diuretics (water tablets)
- if you experience a fast or irregular heartbeat, fainting, collapse or dizziness on standing up which may indicate abnormal functioning of the heart rate
- if you have a problem with dilatation of the pupil of the eye (mydriasis) or if you have or have previously had eye problems, Such as certain kinds of glaucoma (increased pressure in the eye).

Please consult your doctor, even if these statements were applicable to you at any time in the past.

#### Please note

Some patients with manic-depressive illness may enter into a manic phase. This is characterized by unusual and rapidly changing ideas, inappropriate happiness and excessive physical activity. If you experience this, contact your doctor.

Symptoms such as restlessness or difficulty to sit or stand still (akathisia) can also occur during the first weeks of the treatment. Tell your doctor immediately if you experience these symptoms.

#### Thoughts of suicide and worsening of your depression or anxiety disorder

If you are depressed and/or have anxiety disorders you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this:

- if you have previously had thoughts about killing or harming yourself.
- if you are a **young adult**. Information from clinical trials has shown an increased risk of suicidal behaviour in young adults (less than 25 years old) with psychiatric conditions who were treated with an antidepressant

If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away. You may find it helpful to tell a relative or close friend** that you are depressed or have an anxiety disorder, and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

#### Children and adolescents

Cipramil should normally not be used for children and adolescents under 18 years. Also, you should know that patients under 18 have an increased risk of side-effects such as suicide attempts, suicidal thoughts and hostility (predominately aggression, oppositional behaviour and anger) when they take this class of medicines. Despite this, your doctor may prescribe Cipramil for patients under 18 because he/she decides that this is in their best interest. If your doctor has prescribed Cipramil for a patient under 18 and you want to discuss this, please go back to your doctor. You should inform your doctor if any symptoms listed above develop or worsen when patients under 18 are taking Cipramil. Also, the long term safety effects concerning growth, maturation and cognitive and behavioural development of Cipramil in this age group have not yet been demonstrated.

#### Special information relating to your disease

As with other medicines used to treat depression or related diseases, the improvement is not achieved immediately. After the start of Cipramil treatment it may take several weeks before you experience any improvement. In the treatment of panic disorder it usually takes 2-4 weeks before any improvement is seen.

In the beginning of the treatment certain patients may experience increased anxiety, which will disappear during the continued treatment. Therefore, it is very important that you follow exactly your doctor's orders and do not stop the treatment or change the dose without consulting your doctor.

Occasionally, the symptoms of depression or panic disorder may include thoughts of suicide or self-harm. It is possible that these symptoms continue or get worse until the full antidepressant effect of the medicine becomes apparent. This is more likely to occur if you are a young adult, i.e. under 25 years of age and you have not used antidepressive medicines before.

Sometimes you may be unaware of the above-mentioned symptoms and therefore you may find it helpful to ask a friend or relative to help you to observe the possible signs of change in your behaviour.

Tell your doctor immediately or contact the nearest hospital if you have distressing thoughts or experiences or if any of the above-mentioned symptoms occurs during the treatment.

#### Other medicines and Cipramil

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Some medicinal products may affect the action of another and this can sometimes cause serious adverse reactions.

Tell your doctor if you are taking any of the following medicines:

- "Non-selective monoamine oxidase inhibitors (MAOIs)", containing phenelzine, iproniazid, isocarboxazid, nialamide, and tranylcypromine as active substance. If you have taken any of these medicines you will need to wait 14 days before you start taking Cipramil. After stopping Cipramil you must allow 7 days before taking any of these medicines.
- "Reversible, selective MAO-A inhibitors", containing moclobemide (used to treat depression)
- The antibiotic linezolid
- Lithium (used in the prophylaxis and treatment of manic-depressive disorder) and tryptophan.
- Imipramine and desipramine (both used to treat depression)
- "Irreversible MAO-B inhibitors", containing selegiline (used to treat Parkinson's disease); these increase the risk of side effects. The dose of selegiline must not exceed 10mg per day.
- Metoprolol (used for high blood pressure and/or heart disease); the blood levels of metoprolol are increased. Dose adjustment may be required.
- Sumatriptan and similar medicines (used to treat migraine) and tramadol (used against severe pain); these increase the risk of side effects; if you get any unusual symptoms when using this combination you should see your doctor.
- Cimetidine, omeprazole and esomeprazole (used to treat stomach ulcers) and fluvoxamine (antidepressant), blood levels of Cipramil may be increased and dose adjustment may be required.
- Drugs known to affect the platelet function (e.g. some antipsychotic drugs, acetylsalicylic acid (used as pain killers), non-steroidal anti-inflammatory drugs (used for arthritis)); slightly increased risk of bleeding abnormalities.
- St John's Wort (*Hypericum perforatum*) (a herbal remedy used to treat depression) - concomitant intake with Cipramil may increase the risk of side effects.
- Mefloquin (used to treat malaria), bupropion (used to treat depression) and tramadol (used to treat severe pain) due to a possible risk of a lowered threshold for seizures
- Neuroleptics (medicines to treat schizophrenia, psychosis) and antidepressants (SSRIs) due to a possible risk of a lowered threshold for seizures
- Medicines that decrease blood levels of potassium or magnesium as these conditions increase the risk of life-threatening heart rhythm disorder

Do not take Cipramil if you take medicines for heart rhythm problems or medicines that may affect the heart's rhythm, e.g. Class IA and III antiarrhythmics, antipsychotics (e.g. fenflazine derivatives, pimozide, haloperidol), tricyclic antidepressants, certain antimicrobial agents (e.g. sparfloxacin, moxifloxacin, erythromycin IV, pentamidine, anti-malarial treatment particularly halofantrine), certain antihistamines (astemizole, mizolastine). If you have any further questions about this you should speak to your doctor.

#### Cipramil with food, drink and alcohol

Cipramil can be taken with or without food (see section 3 "How to take Cipramil").

Cipramil has been shown not to increase the effects of alcohol. Nevertheless, it is recommended not to drink alcohol during treatment with Cipramil.

#### Pregnancy, breast-feeding and fertility

Inform your doctor if you are pregnant or planning to become pregnant. Pregnant women should not usually take Cipramil nor should mothers breast-feed their babies while taking this medicine, unless you and your doctor have discussed the risks and benefits involved.

If you take Cipramil during the last 3 months of your pregnancy and until the date of birth you should be aware that the following effects may be seen in your newborn: trouble with breathing, blue-ish skin, fits, body temperature changes, feeding difficulties, vomiting, low blood sugar, stiff or floppy muscles, vivid reflexes, tremor, jitteriness, irritability, lethargy, constant crying, sleepiness and sleeping difficulties. If your newborn baby has any of these symptoms, please contact your doctor immediately.

Make sure your midwife and/or doctor know you are on Cipramil. When taken during pregnancy, particularly in the last 3 months of pregnancy, medicines like Cipramil may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the newborn (PPHN), making the baby breathe faster and appear blue-ish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby you should contact your midwife and/or doctor immediately.

Citalopram has been shown to reduce the quality of sperm in animal studies. Theoretically, this could affect fertility, but impact on human fertility has not been observed as yet.

#### Driving and using machines

Cipramil generally does not cause drowsiness; however, if you feel dizzy or sleepy when you start to take this medicine, do not drive or work any tools or machinery until these effects wear off.

#### Important information about some of the ingredients of Cipramil

This medicinal product contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

### 3. How to take Cipramil

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

#### Adults

##### Depression

The usual dose is 20mg daily. The dose may be increased by your doctor to a maximum of 40mg daily.

##### Panic disorder

The starting dose is 10mg daily for the first week before increasing the dose to 20-30mg daily. The dose may be increased by your doctor to a maximum of 40mg daily.

#### Elderly patients (above 65 years of age)

The starting dose should be decreased to half of the recommended dose, e.g. 10-20mg daily.

Elderly patients should not usually receive more than 20mg daily.

#### Patients with special risks

Patients with liver complaints should not receive more than 20mg daily.

#### Children and adolescents (<18 years)

Cipramil should not be given to children or adolescents.

For further information please see section 2 "What you need to know before you take Cipramil".

#### How and when to take Cipramil

Cipramil is taken every day as a single daily dose. Cipramil can be taken any time of the day with or without food. Swallow the tablets with a drink of water. Do not chew them (they have a bitter taste).

#### Duration of treatment

Like other medicines for depression and panic disorder these tablets may take a few weeks before you feel any improvement. Continue to take Cipramil even if it takes some time before you feel any improvement in your condition.

Never change the dose of the medicine without talking to your doctor first.

The duration of treatment is individual, usually at least 6 months. Continue to take the tablets for as long as your doctor recommends. Do not stop taking them even if you begin to feel better, unless you are told to do so by your doctor. The underlying illness may persist for a long time and if you stop your treatment too soon your symptoms may return.

Patients who have recurrent depression benefit from continued treatment, sometimes for several years, to prevent the occurrence of new depressive episodes.

#### If you take more Cipramil than you should

If you think that you or anyone else may have taken too much Cipramil contact your doctor or nearest hospital emergency department immediately. Do this even if there are no signs of discomfort or poisoning. Take the Cipramil box/container with you if you go to a doctor or hospital.

Some of the signs of an overdose could be life-threatening irregular heart beat, convulsion, change in heart rhythm, drowsiness, coma, vomiting, tremor, decreased blood pressure, increased blood pressure, nausea (feeling sick), serotonin syndrome (see section 4), agitation, dizziness, dilated pupils of the eye, sweating, blue-ish skin, hyperventilation.

#### If you forget to take Cipramil

Do not take a double dose to make up for a forgotten dose. If you forget to take a dose, and you remember before you go to bed, take it straight away. Carry on as usual the next day. If you only remember during the night, or the next day, leave out the missed dose and carry on as usual.

### **If you stop taking Cipramil**

Do not stop taking Cipramil until your doctor tells you to do so. When you have completed your course of treatment, it is generally advised that the dose of Cipramil is gradually reduced over a number of weeks.

If you stop taking Cipramil too quickly, you may experience discontinuation symptoms. Most people find that the symptoms are mild and go away on their own within two weeks. However, in some patients they may be severe in intensity or they may be prolonged (2-3 months or more). If you get severe discontinuation symptoms when you stop taking Cipramil, please contact your doctor.

Discontinuation symptoms include: dizziness, feelings like pins and needles, sleep disturbances (vivid dreams, nightmares, inability to sleep), feeling anxious, headaches, feeling sick (nausea), vomiting, sweating, feeling restless or agitated, tremor, feeling confused or disorientated, feeling emotional or irritable, diarrhoea (loose stools), visual disturbances, fluttering or pounding heartbeat (palpitations). When you have completed your course of treatment it is therefore advised that the dose of Cipramil is gradually reduced over a couple of weeks rather than stopped abruptly.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

***If you experience any of the following symptoms contact your doctor or go to the hospital straight away:***

#### Rare (may affect up to 1 in 1,000 people):

- Hyponatraemia: abnormally low blood levels of sodium which can cause tiredness, confusion, and muscle twitching

#### Not known (frequency cannot be estimated from the available data):

- High fever, agitation, confusion, trembling and abrupt contractions of muscles; these may be signs of a rare condition called serotonin syndrome which has been reported with the combined use of antidepressants.
- Swelling of skin, tongue, lips, or face, feeling dizzy or having difficulties breathing or swallowing (serious allergic reaction)
- Unusual bleeds, including gastrointestinal bleeds
- Fast, irregular heart beat, fainting which could be symptoms of a life-threatening condition known as Torsades de Pointes

The majority of the side effects listed below are mild. Please be aware that many of the effects may also be symptoms of your illness and therefore will improve when you start to get better.

#### Very common (may affect more than 1 in 10 people):

Sleepiness  
Difficulty in sleeping  
Increased sweating  
Dry mouth (as dry mouth may increase the risk of tooth decay it is advisable to brush your teeth more often)  
Nausea (feeling sick)  
Headache

#### Common (may affect up to 1 in 10 people):

Decreased appetite  
Agitation  
Decreased sexual drive  
Anxiety  
Nervousness  
Confusional state  
Abnormal dreams  
Tremor  
Tingling or numbness in the hands or feet  
Dizziness  
Disturbance in attention  
Ringing in the ears (tinnitus)  
Yawning  
Diarrhoea  
Vomiting  
Constipation  
Itching  
Pain in muscle and joints  
Men may experience problems with ejaculation and erection  
Women may experience failure to achieve an orgasm  
Fatigue  
Fever  
Prickling of the skin  
Decreased weight

#### Uncommon (may affect up to 1 in 100 people):

Purpura: red or purple patches/spots (smaller than 1cm) of the skin caused by bleeding underneath  
Increased appetite  
Aggression  
Depersonalisation (feeling detached from yourself)  
Hallucination  
Mania (feeling excessively elated, impulsive, irritable, or irrational)  
Fainting  
Enlarged pupils  
Fast heart beat  
Slow heart beat  
Nettle rash  
Loss of hair  
Rash  
Light sensitivity  
Difficulties urinating  
Excessive menstrual bleeding  
Swelling of the arms or legs  
Increased weight

#### Rare (may affect up to 1 in 1,000 people):

Convulsions  
Involuntary movements  
Taste disturbance  
Bleeding  
Hepatitis

#### Not known (frequency cannot be estimated from the available data):

Thoughts of harming yourself or thoughts of killing yourself, see also section "Warnings and precautions"  
Reduction in blood platelets, which increases risk of bleeding or bruising  
Hypersensitivity (rash)  
Disruption to the hormones affecting urine production  
Hypokalaemia: low blood levels of potassium which can cause muscle weakness, twitching or abnormal heart rhythm  
Panic attack  
Grinding one's teeth  
Restlessness  
Unusual muscle movements or stiffness  
Akathisia (involuntary movements of the muscles)  
Visual disturbance  
Low blood pressure  
Nosebleed  
Ecchymosis: blue or purplish patches (bigger than 1cm) of the skin caused by bleeding underneath  
Sudden swelling of skin or mucosa  
Painful erections  
Flow of milk in men and in women that are not nursing  
Irregular menstrual period  
Abnormal liver function test  
An increased risk of bone fractures has been observed in patients taking these types of medicines  
Alteration of the heart rhythm (called "prolongation of QT interval", seen on ECG, a trace of the electrical activity of the heart).

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine. Reports may be made by following the links to the online reporting option accessible from the IMB homepage, or by completing the downloadable report form also accessible from the IMB website, which may be completed manually and submitted to the IMB via freepost, to the following address:

FREEPOST  
Pharmacovigilance Section  
Irish Medicines Board  
Kevin O'Malley House  
Earlsfort Centre Earlsfort Terrace  
Dublin 2  
Tel: +353 1 6764971  
Fax: +353 1 6762517  
Website: [www.imb.ie](http://www.imb.ie)  
e-mail: [imbpharmacovigilance@imb.ie](mailto:imbpharmacovigilance@imb.ie)

### **5. How to store Cipramil**

Keep out of the sight and reach of children  
Do not store above 25°C.  
Do not use Cipramil after the expiry date (EXP) which is printed on the carton and blister label.  
If your doctor tells you to stop taking your medicine you can return any left-over medicine to the pharmacist (for safe disposal), unless the doctor tells you to keep it at home.  
If your tablets appear to be discoloured, damaged or show any other signs of deterioration, please return them to your pharmacist who will advise you.  
Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

### **6. Contents of the pack and other information**

#### **What Cipramil contains**

Each film-coated tablet contains 20mg citalopram (as hydrobromide) as the active ingredient.

The other ingredients in the tablets are maize starch, lactose monohydrate, copovidone, magnesium stearate, glycerol, microcrystalline cellulose, croscarmellose sodium, hypromellose, macrogol and titanium dioxide (E171).

#### **What Cipramil looks like and contents of the pack**

Cipramil are white, oval tablets, deeply scored on one face with the letters 'C' and 'N' on either side of the score and plain on the reverse.

Cipramil is available in blister packs containing 28 tablets.

#### **Manufacturer**

Cipramil is manufactured by: H. Lundbeck A/S, Ottiliavej 9, DK-2500 Copenhagen, Denmark.

Procured from within the EU by the Parallel Product Authorisation holder:  
Imbat Ltd, Unit L2, North Ring Business Park, Santry, Dublin 9.

Repackaged by: Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, South Yorkshire, DN3 1QR, UK.

Distributed by: Eurodrug Ltd, Santry, Dublin 9.

PPA No: 1151/49/1

Leaflet revision and issue date (ref): 15.01.14

Cipramil® is a registered trademark of H. Lundbeck A/S.

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Cipramil® 20mg Film-coated Tablets Citalopram (as hydrobromide)

Your medicine is available using the name Cipramil 20mg Film-coated Tablets but will be referred to as Cipramil throughout this leaflet.

#### Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

#### What is in this leaflet

- What Cipramil is and what it is used for
- What you need to know before you take Cipramil
- How to take Cipramil
- Possible side effects
- How to store Cipramil
- Contents of the pack and other information

#### 1. What Cipramil is and what it is used for

##### How does Cipramil work

Cipramil belongs to a group of antidepressants called Selective Serotonin Reuptake Inhibitors (SSRIs). These medicines act on the serotonin-system in the brain by increasing the serotonin level. Disturbances in the serotonin-system are considered an important factor in the development of depression and related diseases.

##### What is Cipramil used for

Cipramil contains citalopram and is used for the treatment of depression and when you feel better, to help prevent these symptoms recurring.

Further, Cipramil is used for long-term treatment to prevent the occurrence of new depressive episodes in patients who have recurrent depression.

Cipramil is also beneficial in relieving symptoms in patients prone to panic attacks.

Ask your doctor if you have any questions about why Cipramil has been prescribed for you.

#### 2. What you need to know before you take Cipramil

##### Do not take Cipramil:

- if you are allergic to citalopram or any of the other ingredients of this medicine (listed in section 6)
- if you take other medicines which belong to a group called monoamine oxidase inhibitors (MAOIs). MAOIs include medicines such as phenelzine, iproniazid, isocarboxazid, nialamide, tranylcypromine, selegiline (used in the treatment of Parkinson's disease), moclobemide (used in the treatment of depression) and linezolid (an antibiotic)
- if you are born with or have had an episode of abnormal heart rhythm (seen at ECG; an examination to evaluate how the heart is functioning)
- if you take medicines for heart rhythm problems or that may affect the heart's rhythm (see "Other medicines and Cipramil")

Even if you have finished taking MAOIs, you will need to wait 2 weeks before you start getting your Cipramil treatment.

After stopping Cipramil you must allow 1 week before taking any MAOI.

##### Warnings and precautions

Please tell your doctor if you have any other condition or illness, as your doctor may need to take this into consideration. In particular, tell your doctor:

- if you have episodes of mania or panic disorder
- if you suffer from impaired liver or kidney function. Your doctor may need to adjust your dosage
- if you have diabetes. Treatment with Cipramil may alter glycaemic control. Insulin and/or oral hypoglycaemic dosage may need to be adjusted.
- if you have epilepsy. Treatment with Cipramil should be stopped if seizures occur or if there is an increase in the seizure frequency (see also section 4 "Possible side effects")
- if you have some kind of bleeding disorder
- if you have a decreased level of sodium in the blood
- if you are receiving electroconvulsive treatment
- if you suffer or have suffered from heart problems or have recently had a heart attack
- if you have a low resting heart-rate and/or you know that you may have salt depletion as a result of prolonged severe diarrhoea and vomiting (being sick) or usage of diuretics (water tablets)
- if you experience a fast or irregular heartbeat, fainting, collapse or dizziness on standing up which may indicate abnormal functioning of the heart rate
- if you have a problem with dilatation of the pupil of the eye (mydriasis) or if you have or have previously had eye problems, Such as certain kinds of glaucoma (increased pressure in the eye).

Please consult your doctor, even if these statements were applicable to you at any time in the past.

##### Please note

Some patients with manic-depressive illness may enter into a manic phase. This is characterized by unusual and rapidly changing ideas, inappropriate happiness and excessive physical activity. If you experience this, contact your doctor.

Symptoms such as restlessness or difficulty to sit or stand still (akathisia) can also occur during the first weeks of the treatment. Tell your doctor immediately if you experience these symptoms.

#### Thoughts of suicide and worsening of your depression or anxiety disorder

If you are depressed and/or have anxiety disorders you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this:

- if you have previously had thoughts about killing or harming yourself.
- if you are a **young adult**. Information from clinical trials has shown an increased risk of suicidal behaviour in young adults (less than 25 years old) with psychiatric conditions who were treated with an antidepressant

If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away. You may find it helpful to tell a relative or close friend** that you are depressed or have an anxiety disorder, and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

#### Children and adolescents

Cipramil should normally not be used for children and adolescents under 18 years. Also, you should know that patients under 18 have an increased risk of side-effects such as suicide attempts, suicidal thoughts and hostility (predominately aggression, oppositional behaviour and anger) when they take this class of medicines. Despite this, your doctor may prescribe Cipramil for patients under 18 because he/she decides that this is in their best interest. If your doctor has prescribed Cipramil for a patient under 18 and you want to discuss this, please go back to your doctor. You should inform your doctor if any symptoms listed above develop or worsen when patients under 18 are taking Cipramil. Also, the long term safety effects concerning growth, maturation and cognitive and behavioural development of Cipramil in this age group have not yet been demonstrated.

#### Special information relating to your disease

As with other medicines used to treat depression or related diseases, the improvement is not achieved immediately. After the start of Cipramil treatment it may take several weeks before you experience any improvement. In the treatment of panic disorder it usually takes 2-4 weeks before any improvement is seen.

In the beginning of the treatment certain patients may experience increased anxiety, which will disappear during the continued treatment. Therefore, it is very important that you follow exactly your doctor's orders and do not stop the treatment or change the dose without consulting your doctor.

Occasionally, the symptoms of depression or panic disorder may include thoughts of suicide or self-harm. It is possible that these symptoms continue or get worse until the full antidepressant effect of the medicine becomes apparent. This is more likely to occur if you are a young adult, i.e. under 25 years of age and you have not used antidepressive medicines before.

Sometimes you may be unaware of the above-mentioned symptoms and therefore you may find it helpful to ask a friend or relative to help you to observe the possible signs of change in your behaviour.

Tell your doctor immediately or contact the nearest hospital if you have distressing thoughts or experiences or if any of the above-mentioned symptoms occurs during the treatment.

#### Other medicines and Cipramil

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Some medicinal products may affect the action of another and this can sometimes cause serious adverse reactions.

Tell your doctor if you are taking any of the following medicines:

- "Non-selective monoamine oxidase inhibitors (MAOIs)", containing phenelzine, iproniazid, isocarboxazid, nialamide, and tranylcypromine as active substance. If you have taken any of these medicines you will need to wait 14 days before you start taking Cipramil. After stopping Cipramil you must allow 7 days before taking any of these medicines.
- "Reversible, selective MAO-A inhibitors", containing moclobemide (used to treat depression)
- The antibiotic linezolid
- Lithium (used in the prophylaxis and treatment of manic-depressive disorder) and tryptophan.
- Imipramine and desipramine (both used to treat depression)
- "Irreversible MAO-B inhibitors", containing selegiline (used to treat Parkinson's disease); these increase the risk of side effects. The dose of selegiline must not exceed 10mg per day.
- Metoprolol (used for high blood pressure and/or heart disease); the blood levels of metoprolol are increased. Dose adjustment may be required.
- Sumatriptan and similar medicines (used to treat migraine) and tramadol (used against severe pain); these increase the risk of side effects; if you get any unusual symptoms when using this combination you should see your doctor.
- Cimetidine, omeprazole and esomeprazole (used to treat stomach ulcers) and fluvoxamine (antidepressant), blood levels of Cipramil may be increased and dose adjustment may be required.
- Drugs known to affect the platelet function (e.g. some antipsychotic drugs, acetylsalicylic acid (used as pain killers), non-steroidal anti-inflammatory drugs (used for arthritis)); slightly increased risk of bleeding abnormalities.
- St John's Wort (*Hypericum perforatum*) (a herbal remedy used to treat depression) - concomitant intake with Cipramil may increase the risk of side effects.
- Mefloquin (used to treat malaria), bupropion (used to treat depression) and tramadol (used to treat severe pain) due to a possible risk of a lowered threshold for seizures
- Neuroleptics (medicines to treat schizophrenia, psychosis) and antidepressants (SSRIs) due to a possible risk of a lowered threshold for seizures
- Medicines that decrease blood levels of potassium or magnesium as these conditions increase the risk of life-threatening heart rhythm disorder

Do not take Cipramil if you take medicines for heart rhythm problems or medicines that may affect the heart's rhythm, e.g. Class IA and III antiarrhythmics, antipsychotics (e.g. fenflazine derivatives, pimozide, haloperidol), tricyclic antidepressants, certain antimicrobial agents (e.g. sparfloxacin, moxifloxacin, erythromycin IV, pentamidine, anti-malarial treatment particularly halofantrine), certain antihistamines (astemizole, mizolastine). If you have any further questions about this you should speak to your doctor.

#### Cipramil with food, drink and alcohol

Cipramil can be taken with or without food (see section 3 "How to take Cipramil").

Cipramil has been shown not to increase the effects of alcohol. Nevertheless, it is recommended not to drink alcohol during treatment with Cipramil.

#### Pregnancy, breast-feeding and fertility

Inform your doctor if you are pregnant or planning to become pregnant. Pregnant women should not usually take Cipramil nor should mothers breast-feed their babies while taking this medicine, unless you and your doctor have discussed the risks and benefits involved.

If you take Cipramil during the last 3 months of your pregnancy and until the date of birth you should be aware that the following effects may be seen in your newborn: trouble with breathing, blue-ish skin, fits, body temperature changes, feeding difficulties, vomiting, low blood sugar, stiff or floppy muscles, vivid reflexes, tremor, jitteriness, irritability, lethargy, constant crying, sleepiness and sleeping difficulties. If your newborn baby has any of these symptoms, please contact your doctor immediately.

Make sure your midwife and/or doctor know you are on Cipramil. When taken during pregnancy, particularly in the last 3 months of pregnancy, medicines like Cipramil may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the newborn (PPHN), making the baby breathe faster and appear blue-ish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby you should contact your midwife and/or doctor immediately.

Citalopram has been shown to reduce the quality of sperm in animal studies. Theoretically, this could affect fertility, but impact on human fertility has not been observed as yet.

#### Driving and using machines

Cipramil generally does not cause drowsiness; however, if you feel dizzy or sleepy when you start to take this medicine, do not drive or work any tools or machinery until these effects wear off.

#### Important information about some of the ingredients of Cipramil

This medicinal product contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

#### 3. How to take Cipramil

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

##### Adults

###### Depression

The usual dose is 20mg daily. The dose may be increased by your doctor to a maximum of 40mg daily.

###### Panic disorder

The starting dose is 10mg daily for the first week before increasing the dose to 20-30mg daily. The dose may be increased by your doctor to a maximum of 40mg daily.

##### Elderly patients (above 65 years of age)

The starting dose should be decreased to half of the recommended dose, e.g. 10-20mg daily.

Elderly patients should not usually receive more than 20mg daily.

##### Patients with special risks

Patients with liver complaints should not receive more than 20mg daily.

##### Children and adolescents (<18 years)

Cipramil should not be given to children or adolescents. For further information please see section 2 "What you need to know before you take Cipramil".

#### How and when to take Cipramil

Cipramil is taken every day as a single daily dose. Cipramil can be taken any time of the day with or without food. Swallow the tablets with a drink of water. Do not chew them (they have a bitter taste).

#### Duration of treatment

Like other medicines for depression and panic disorder these tablets may take a few weeks before you feel any improvement. Continue to take Cipramil even if it takes some time before you feel any improvement in your condition.

Never change the dose of the medicine without talking to your doctor first.

The duration of treatment is individual, usually at least 6 months. Continue to take the tablets for as long as your doctor recommends. Do not stop taking them even if you begin to feel better, unless you are told to do so by your doctor. The underlying illness may persist for a long time and if you stop your treatment too soon your symptoms may return.

Patients who have recurrent depression benefit from continued treatment, sometimes for several years, to prevent the occurrence of new depressive episodes.

#### If you take more Cipramil than you should

If you think that you or anyone else may have taken too much Cipramil contact your doctor or nearest hospital emergency department immediately. Do this even if there are no signs of discomfort or poisoning. Take the Cipramil box/container with you if you go to a doctor or hospital.

Some of the signs of an overdose could be life-threatening irregular heart beat, convulsion, change in heart rhythm, drowsiness, coma, vomiting, tremor, decreased blood pressure, increased blood pressure, nausea (feeling sick), serotonin syndrome (see section 4), agitation, dizziness, dilated pupils of the eye, sweating, blue-ish skin, hyperventilation.

#### If you forget to take Cipramil

Do not take a double dose to make up for a forgotten dose. If you forget to take a dose, and you remember before you go to bed, take it straight away. Carry on as usual the next day. If you only remember during the night, or the next day, leave out the missed dose and carry on as usual.

#### **If you stop taking Cipramil**

Do not stop taking Cipramil until your doctor tells you to do so. When you have completed your course of treatment, it is generally advised that the dose of Cipramil is gradually reduced over a number of weeks.

If you stop taking Cipramil too quickly, you may experience discontinuation symptoms. Most people find that the symptoms are mild and go away on their own within two weeks. However, in some patients they may be severe in intensity or they may be prolonged (2-3 months or more). If you get severe discontinuation symptoms when you stop taking Cipramil, please contact your doctor.

Discontinuation symptoms include: dizziness, feelings like pins and needles, sleep disturbances (vivid dreams, nightmares, inability to sleep), feeling anxious, headaches, feeling sick (nausea), vomiting, sweating, feeling restless or agitated, tremor, feeling confused or disorientated, feeling emotional or irritable, diarrhoea (loose stools), visual disturbances, fluttering or pounding heartbeat (palpitations). When you have completed your course of treatment it is therefore advised that the dose of Cipramil is gradually reduced over a couple of weeks rather than stopped abruptly.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

#### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

***If you experience any of the following symptoms contact your doctor or go to the hospital straight away:***

##### Rare (may affect up to 1 in 1,000 people):

- Hyponatraemia: abnormally low blood levels of sodium which can cause tiredness, confusion, and muscle twitching

##### Not known (frequency cannot be estimated from the available data):

- High fever, agitation, confusion, trembling and abrupt contractions of muscles: these may be signs of a rare condition called serotonin syndrome which has been reported with the combined use of antidepressants.
- Swelling of skin, tongue, lips, or face, feeling dizzy or having difficulties breathing or swallowing (serious allergic reaction)
- Unusual bleeds, including gastrointestinal bleeds
- Fast, irregular heart beat, fainting which could be symptoms of a life-threatening condition known as Torsades de Pointes

The majority of the side effects listed below are mild. Please be aware that many of the effects may also be symptoms of your illness and therefore will improve when you start to get better.

##### Very common (may affect more than 1 in 10 people):

Sleepiness  
Difficulty in sleeping  
Increased sweating  
Dry mouth (as dry mouth may increase the risk of tooth decay it is advisable to brush your teeth more often)  
Nausea (feeling sick)  
Headache

##### Common (may affect up to 1 in 10 people):

Decreased appetite  
Agitation  
Decreased sexual drive  
Anxiety  
Nervousness  
Confusional state  
Abnormal dreams  
Tremor  
Tingling or numbness in the hands or feet  
Dizziness  
Disturbance in attention  
Ringing in the ears (tinnitus)  
Yawning  
Diarrhoea  
Vomiting  
Constipation  
Itching  
Pain in muscle and joints  
Men may experience problems with ejaculation and erection  
Women may experience failure to achieve an orgasm  
Fatigue  
Fever  
Prickling of the skin  
Decreased weight

##### Uncommon (may affect up to 1 in 100 people):

Purpura: red or purple patches/spots (smaller than 1cm) of the skin caused by bleeding underneath  
Increased appetite  
Aggression  
Depersonalisation (feeling detached from yourself)  
Hallucination  
Mania (feeling excessively elated, impulsive, irritable, or irrational)  
Fainting  
Enlarged pupils  
Fast heart beat  
Slow heart beat  
Nettle rash  
Loss of hair  
Rash  
Light sensitivity  
Difficulties urinating  
Excessive menstrual bleeding  
Swelling of the arms or legs  
Increased weight

##### Rare (may affect up to 1 in 1,000 people):

Convulsions  
Involuntary movements  
Taste disturbance  
Bleeding  
Hepatitis

##### Not known (frequency cannot be estimated from the available data):

Thoughts of harming yourself or thoughts of killing yourself, see also section "Warnings and precautions"  
Reduction in blood platelets, which increases risk of bleeding or bruising  
Hypersensitivity (rash)  
Disruption to the hormones affecting urine production  
Hypokalaemia: low blood levels of potassium which can cause muscle weakness, twitching or abnormal heart rhythm  
Panic attack  
Grinding one's teeth  
Restlessness  
Unusual muscle movements or stiffness  
Akathisia (involuntary movements of the muscles)  
Visual disturbance  
Low blood pressure  
Nosebleed  
Ecchymosis: blue or purplish patches (bigger than 1cm) of the skin caused by bleeding underneath  
Sudden swelling of skin or mucosa  
Painful erections  
Flow of milk in men and in women that are not nursing  
Irregular menstrual period  
Abnormal liver function test  
An increased risk of bone fractures has been observed in patients taking these types of medicines  
Alteration of the heart rhythm (called "prolongation of QT interval", seen on ECG, a trace of the electrical activity of the heart).

##### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine. Reports may be made by following the links to the online reporting option accessible from the IMB homepage, or by completing the downloadable report form also accessible from the IMB website, which may be completed manually and submitted to the IMB via freepost, to the following address:

FREEPOST  
Pharmacovigilance Section  
Irish Medicines Board  
Kevin O'Malley House  
Earlsfort Centre Earlsfort Terrace  
Dublin 2  
Tel: +353 1 6764971  
Fax: +353 1 6762517  
Website: [www.imb.ie](http://www.imb.ie)  
e-mail: [imbpharmacovigilance@imb.ie](mailto:imbpharmacovigilance@imb.ie)

#### **5. How to store Cipramil**

Keep out of the sight and reach of children  
Do not store above 25°C.  
Do not use Cipramil after the expiry date (EXP) which is printed on the carton and blister label.  
If your doctor tells you to stop taking your medicine you can return any left-over medicine to the pharmacist (for safe disposal), unless the doctor tells you to keep it at home.  
If your tablets appear to be discoloured, damaged or show any other signs of deterioration, please return them to your pharmacist who will advise you.  
Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

#### **6. Contents of the pack and other information**

##### **What Cipramil contains**

Each film-coated tablet contains 20mg citalopram (as hydrobromide) as the active ingredient.

The other ingredients in the tablets are: maize starch, lactose monohydrate, copovidone, magnesium stearate, glycerol, microcrystalline cellulose, croscarmellose sodium, hypromellose, macrogol and titanium dioxide (E171).

##### **What Cipramil looks like and contents of the pack**

Cipramil are white, oval tablets, deeply scored on one face with the letters 'C' and 'N' on either side of the score and plain on the reverse.

Cipramil is available in blister packs containing 28 tablets.

##### **Manufacturer**

Cipramil is manufactured by: H. Lundbeck A/S, Ottiliavej 9, DK-2500 Copenhagen, Denmark.

Procured from within the EU by the Parallel Product Authorisation holder:  
Imbat Ltd, Unit L2, North Ring Business Park, Santry, Dublin 9.

Repackaged by: Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, South Yorkshire, DN3 1QR, UK.

Distributed by: Eurodrug Ltd, Santry, Dublin 9.

PPA No: 1151/49/1

Leaflet revision and issue date (ref): 15.01.14

Cipramil® is a registered trademark of H. Lundbeck A/S.



## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Cipramil® 20mg Film-coated Tablets

Citalopram (as hydrobromide)

Your medicine is available using the name Cipramil 20mg Film-coated Tablets but will be referred to as Cipramil throughout this leaflet.

### Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

### What is in this leaflet

- What Cipramil is and what it is used for
- What you need to know before you take Cipramil
- How to take Cipramil
- Possible side effects
- How to store Cipramil
- Contents of the pack and other information

## 1. What Cipramil is and what it is used for

### How does Cipramil work

Cipramil belongs to a group of antidepressants called Selective Serotonin Reuptake Inhibitors (SSRIs). These medicines act on the serotonin-system in the brain by increasing the serotonin level. Disturbances in the serotonin-system are considered an important factor in the development of depression and related diseases.

### What is Cipramil used for

Cipramil contains citalopram and is used for the treatment of depression and when you feel better, to help prevent these symptoms recurring.

Further, Cipramil is used for long-term treatment to prevent the occurrence of new depressive episodes in patients who have recurrent depression.

Cipramil is also beneficial in relieving symptoms in patients prone to panic attacks.

Ask your doctor if you have any questions about why Cipramil has been prescribed for you.

## 2. What you need to know before you take Cipramil

### Do not take Cipramil:

- if you are allergic to citalopram or any of the other ingredients of this medicine (listed in section 6)
- if you take other medicines which belong to a group called monoamine oxidase inhibitors (MAOIs). MAOIs include medicines such as phenelzine, iproniazid, isocarboxazid, nialamide, tranylcypromine, selegiline (used in the treatment of Parkinson's disease), moclobemide (used in the treatment of depression) and linezolid (an antibiotic)
- if you are born with or have had an episode of abnormal heart rhythm (seen at ECG; an examination to evaluate how the heart is functioning)
- if you take medicines for heart rhythm problems or that may affect the heart's rhythm (see "Other medicines and Cipramil")

Even if you have finished taking MAOIs, you will need to wait 2 weeks before you start getting your Cipramil treatment.

After stopping Cipramil you must allow 1 week before taking any MAOI.

### Warnings and precautions

Please tell your doctor if you have any other condition or illness, as your doctor may need to take this into consideration. In particular, tell your doctor:

- if you have episodes of mania or panic disorder
- if you suffer from impaired liver or kidney function. Your doctor may need to adjust your dosage
- if you have diabetes. Treatment with Cipramil may alter glycaemic control. Insulin and/or oral hypoglycaemic dosage may need to be adjusted.
- if you have epilepsy. Treatment with Cipramil should be stopped if seizures occur or if there is an increase in the seizure frequency (see also section 4 "Possible side effects")
- if you have some kind of bleeding disorder
- if you have a decreased level of sodium in the blood
- if you are receiving electroconvulsive treatment
- if you suffer or have suffered from heart problems or have recently had a heart attack
- if you have a low resting heart-rate and/or you know that you may have salt depletion as a result of prolonged severe diarrhoea and vomiting (being sick) or usage of diuretics (water tablets)
- if you experience a fast or irregular heartbeat, fainting, collapse or dizziness on standing up which may indicate abnormal functioning of the heart rate
- if you have a problem with dilatation of the pupil of the eye (mydriasis) or if you have or have previously had eye problems. Such as certain kinds of glaucoma (increased pressure in the eye).

Please consult your doctor, even if these statements were applicable to you at any time in the past.

### Please note

Some patients with manic-depressive illness may enter into a manic phase. This is characterized by unusual and rapidly changing ideas, inappropriate happiness and excessive physical activity. If you experience this, contact your doctor.

Symptoms such as restlessness or difficulty to sit or stand still (akathisia) can also occur during the first weeks of the treatment. Tell your doctor immediately if you experience these symptoms.

### Thoughts of suicide and worsening of your depression or anxiety disorder

If you are depressed and/or have anxiety disorders you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this:

- if you have previously had thoughts about killing or harming yourself.
- if you are a **young adult**. Information from clinical trials has shown an increased risk of suicidal behaviour in young adults (less than 25 years old) with psychiatric conditions who were treated with an antidepressant

If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away. You may find it helpful to tell a relative or close friend** that you are depressed or have an anxiety disorder, and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

### Children and adolescents

Cipramil should normally not be used for children and adolescents under 18 years. Also, you should know that patients under 18 have an increased risk of side-effects such as suicide attempts, suicidal thoughts and hostility (predominately aggression, oppositional behaviour and anger) when they take this class of medicines. Despite this, your doctor may prescribe Cipramil for patients under 18 because he/she decides that this is in their best interest. If your doctor has prescribed Cipramil for a patient under 18 and you want to discuss this, please go back to your doctor. You should inform your doctor if any symptoms listed above develop or worsen when patients under 18 are taking Cipramil. Also, the long term safety effects concerning growth, maturation and cognitive and behavioural development of Cipramil in this age group have not yet been demonstrated.

### Special information relating to your disease

As with other medicines used to treat depression or related diseases, the improvement is not achieved immediately. After the start of Cipramil treatment it may take several weeks before you experience any improvement. In the treatment of panic disorder it usually takes 2-4 weeks before any improvement is seen.

In the beginning of the treatment certain patients may experience increased anxiety, which will disappear during the continued treatment. Therefore, it is very important that you follow exactly your doctor's orders and do not stop the treatment or change the dose without consulting your doctor.

Occasionally, the symptoms of depression or panic disorder may include thoughts of suicide or self-harm. It is possible that these symptoms continue or get worse until the full antidepressant effect of the medicine becomes apparent. This is more likely to occur if you are a young adult, i.e. under 25 years of age and you have not used antidepressive medicines before.

Sometimes you may be unaware of the above-mentioned symptoms and therefore you may find it helpful to ask a friend or relative to help you to observe the possible signs of change in your behaviour.

Tell your doctor immediately or contact the nearest hospital if you have distressing thoughts or experiences or if any of the above-mentioned symptoms occurs during the treatment.

### Other medicines and Cipramil

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Some medicinal products may affect the action of another and this can sometimes cause serious adverse reactions.

Tell your doctor if you are taking any of the following medicines:

- "Non-selective monoamine oxidase inhibitors (MAOIs)", containing phenelzine, iproniazid, isocarboxazid, nialamide, and tranylcypromine as active substance. If you have taken any of these medicines you will need to wait 14 days before you start taking Cipramil. After stopping Cipramil you must allow 7 days before taking any of these medicines.
- "Reversible, selective MAO-A inhibitors", containing moclobemide (used to treat depression)
- The antibiotic linezolid
- Lithium (used in the prophylaxis and treatment of manic-depressive disorder) and tryptophan.
- Imipramine and desipramine (both used to treat depression)
- "Irreversible MAO-B inhibitors", containing selegiline (used to treat Parkinson's disease); these increase the risk of side effects. The dose of selegiline must not exceed 10mg per day.
- Metoprolol (used for high blood pressure and/or heart disease); the blood levels of metoprolol are increased. Dose adjustment may be required.
- Sumatriptan and similar medicines (used to treat migraine) and tramadol (used against severe pain); these increase the risk of side effects; if you get any unusual symptoms when using this combination you should see your doctor.
- Cimetidine, omeprazole and esomeprazole (used to treat stomach ulcers) and fluvoxamine (antidepressant), blood levels of Cipramil may be increased and dose adjustment may be required.
- Drugs known to affect the platelet function (e.g. some antipsychotic drugs, acetylsalicylic acid (used as pain killers), non-steroidal anti-inflammatory drugs (used for arthritis)); slightly increased risk of bleeding abnormalities.
- St John's Wort (*Hypericum perforatum*) (a herbal remedy used to treat depression) – concomitant intake with Cipramil may increase the risk of side effects.
- Mefloquin (used to treat malaria), bupropion (used to treat depression) and tramadol (used to treat severe pain) due to a possible risk of a lowered threshold for seizures
- Neuroleptics (medicines to treat schizophrenia, psychosis) and antidepressants (SSRIs) due to a possible risk of a lowered threshold for seizures
- Medicines that decrease blood levels of potassium or magnesium as these conditions increase the risk of life-threatening heart rhythm disorder

Do not take Cipramil if you take medicines for heart rhythm problems or medicines that may affect the heart's rhythm, e.g. Class IA and III antiarrhythmics, antipsychotics (e.g. fentiazine derivatives, pimozide, haloperidol), tricyclic antidepressants, certain antimicrobial agents (e.g. sparfloxacin, moxifloxacin, erythromycin IV, pentamidine, anti-malarial treatment particularly halofantrine), certain antihistamines (astemizole, mizolastine). If you have any further questions about this you should speak to your doctor.

### Cipramil with food, drink and alcohol

Cipramil can be taken with or without food (see section 3 "How to take Cipramil").

Cipramil has been shown not to increase the effects of alcohol. Nevertheless, it is recommended not to drink alcohol during treatment with Cipramil.

### Pregnancy, breast-feeding and fertility

Inform your doctor if you are pregnant or planning to become pregnant. Pregnant women should not usually take Cipramil nor should mothers breast-feed their babies while taking this medicine, unless you and your doctor have discussed the risks and benefits involved.

If you take Cipramil during the last 3 months of your pregnancy and until the date of birth you should be aware that the following effects may be seen in your newborn: trouble with breathing, blue-ish skin, fits, body temperature changes, feeding difficulties, vomiting, low blood sugar, stiff or floppy muscles, vivid reflexes, tremor, jitteriness, irritability, lethargy, constant crying, sleepiness and sleeping difficulties. If your newborn baby has any of these symptoms, please contact your doctor immediately.

Make sure your midwife and/or doctor know you are on Cipramil. When taken during pregnancy, particularly in the last 3 months of pregnancy, medicines like Cipramil may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the newborn (PPHN), making the baby breathe faster and appear blue-ish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby you should contact your midwife and/or doctor immediately.

Citalopram has been shown to reduce the quality of sperm in animal studies. Theoretically, this could affect fertility, but impact on human fertility has not been observed as yet.

### Driving and using machines

Cipramil generally does not cause drowsiness; however, if you feel dizzy or sleepy when you start to take this medicine, do not drive or work any tools or machinery until these effects wear off.

### Important information about some of the ingredients of Cipramil

This medicinal product contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

## 3. How to take Cipramil

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

### Adults

#### Depression

The usual dose is 20mg daily. The dose may be increased by your doctor to a maximum of 40mg daily.

#### Panic disorder

The starting dose is 10mg daily for the first week before increasing the dose to 20-30mg daily. The dose may be increased by your doctor to a maximum of 40mg daily.

### Elderly patients (above 65 years of age)

The starting dose should be decreased to half of the recommended dose, e.g. 10-20mg daily.

Elderly patients should not usually receive more than 20mg daily.

### Patients with special risks

Patients with liver complaints should not receive more than 20mg daily.

### Children and adolescents (<18 years)

Cipramil should not be given to children or adolescents. For further information please see section 2 "What you need to know before you take Cipramil".

### How and when to take Cipramil

Cipramil is taken every day as a single daily dose. Cipramil can be taken any time of the day with or without food. Swallow the tablets with a drink of water. Do not chew them (they have a bitter taste).

### Duration of treatment

Like other medicines for depression and panic disorder these tablets may take a few weeks before you feel any improvement. Continue to take Cipramil even if it takes some time before you feel any improvement in your condition.

Never change the dose of the medicine without talking to your doctor first.

The duration of treatment is individual, usually at least 6 months. Continue to take the tablets for as long as your doctor recommends. Do not stop taking them even if you begin to feel better, unless you are told to do so by your doctor. The underlying illness may persist for a long time and if you stop your treatment too soon your symptoms may return.

Patients who have recurrent depression benefit from continued treatment, sometimes for several years, to prevent the occurrence of new depressive episodes.

### If you take more Cipramil than you should

If you think that you or anyone else may have taken too much Cipramil contact your doctor or nearest hospital emergency department immediately. Do this even if there are no signs of discomfort or poisoning. Take the Cipramil box/container with you if you go to a doctor or hospital.

Some of the signs of an overdose could be life-threatening irregular heart beat, convulsion, change in heart rhythm, drowsiness, coma, vomiting, tremor, decreased blood pressure, increased blood pressure, nausea (feeling sick), serotonin syndrome (see section 4), agitation, dizziness, dilated pupils of the eye, sweating, blue-ish skin, hyperventilation.

### If you forget to take Cipramil

Do not take a double dose to make up for a forgotten dose. If you forget to take a dose, and you remember before you go to bed, take it straight away. Carry on as usual the next day. If you only remember during the night, or the next day, leave out the missed dose and carry on as usual.

### **If you stop taking Cipramil**

Do not stop taking Cipramil until your doctor tells you to do so. When you have completed your course of treatment, it is generally advised that the dose of Cipramil is gradually reduced over a number of weeks.

If you stop taking Cipramil too quickly, you may experience discontinuation symptoms. Most people find that the symptoms are mild and go away on their own within two weeks. However, in some patients they may be severe in intensity or they may be prolonged (2-3 months or more). If you get severe discontinuation symptoms when you stop taking Cipramil, please contact your doctor.

Discontinuation symptoms include: dizziness, feelings like pins and needles, sleep disturbances (vivid dreams, nightmares, inability to sleep), feeling anxious, headaches, feeling sick (nausea), vomiting, sweating, feeling restless or agitated, tremor, feeling confused or disorientated, feeling emotional or irritable, diarrhoea (loose stools), visual disturbances, fluttering or pounding heartbeat (palpitations). When you have completed your course of treatment it is therefore advised that the dose of Cipramil is gradually reduced over a couple of weeks rather than stopped abruptly.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

***If you experience any of the following symptoms contact your doctor or go to the hospital straight away:***

#### Rare (may affect up to 1 in 1,000 people):

- Hyponatraemia: abnormally low blood levels of sodium which can cause tiredness, confusion, and muscle twitching

#### Not known (frequency cannot be estimated from the available data):

- High fever, agitation, confusion, trembling and abrupt contractions of muscles; these may be signs of a rare condition called serotonin syndrome which has been reported with the combined use of antidepressants.
- Swelling of skin, tongue, lips, or face, feeling dizzy or having difficulties breathing or swallowing (serious allergic reaction)
- Unusual bleeds, including gastrointestinal bleeds
- Fast, irregular heart beat, fainting which could be symptoms of a life-threatening condition known as Torsades de Pointes

The majority of the side effects listed below are mild. Please be aware that many of the effects may also be symptoms of your illness and therefore will improve when you start to get better.

#### Very common (may affect more than 1 in 10 people):

Sleepiness  
Difficulty in sleeping  
Increased sweating  
Dry mouth (as dry mouth may increase the risk of tooth decay it is advisable to brush your teeth more often)  
Nausea (feeling sick)  
Headache

#### Common (may affect up to 1 in 10 people):

Decreased appetite  
Agitation  
Decreased sexual drive  
Anxiety  
Nervousness  
Confusional state  
Abnormal dreams  
Tremor  
Tingling or numbness in the hands or feet  
Dizziness  
Disturbance in attention  
Ringing in the ears (tinnitus)  
Yawning  
Diarrhoea  
Vomiting  
Constipation  
Itching  
Pain in muscle and joints  
Men may experience problems with ejaculation and erection  
Women may experience failure to achieve an orgasm  
Fatigue  
Fever  
Prickling of the skin  
Decreased weight

#### Uncommon (may affect up to 1 in 100 people):

Purpura: red or purple patches/spots (smaller than 1cm) of the skin caused by bleeding underneath  
Increased appetite  
Aggression  
Depersonalisation (feeling detached from yourself)  
Hallucination  
Mania (feeling excessively elated, impulsive, irritable, or irrational)  
Fainting  
Enlarged pupils  
Fast heart beat  
Slow heart beat  
Nettle rash  
Loss of hair  
Rash  
Light sensitivity  
Difficulties urinating  
Excessive menstrual bleeding  
Swelling of the arms or legs  
Increased weight

#### Rare (may affect up to 1 in 1,000 people):

Convulsions  
Involuntary movements  
Taste disturbance  
Bleeding  
Hepatitis

#### Not known (frequency cannot be estimated from the available data):

Thoughts of harming yourself or thoughts of killing yourself, see also section "Warnings and precautions"  
Reduction in blood platelets, which increases risk of bleeding or bruising  
Hypersensitivity (rash)  
Disruption to the hormones affecting urine production  
Hypokalaemia: low blood levels of potassium which can cause muscle weakness, twitching or abnormal heart rhythm  
Panic attack  
Grinding one's teeth  
Restlessness  
Unusual muscle movements or stiffness  
Akathisia (involuntary movements of the muscles)  
Visual disturbance  
Low blood pressure  
Nosebleed  
Ecchymosis: blue or purplish patches (bigger than 1cm) of the skin caused by bleeding underneath  
Sudden swelling of skin or mucosa  
Painful erections  
Flow of milk in men and in women that are not nursing  
Irregular menstrual period  
Abnormal liver function test  
An increased risk of bone fractures has been observed in patients taking these types of medicines  
Alteration of the heart rhythm (called "prolongation of QT interval", seen on ECG, a trace of the electrical activity of the heart).

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine. Reports may be made by following the links to the online reporting option accessible from the IMB homepage, or by completing the downloadable report form also accessible from the IMB website, which may be completed manually and submitted to the IMB via freepost, to the following address:

FREEPOST  
Pharmacovigilance Section  
Irish Medicines Board  
Kevin O'Malley House  
Earlsfort Centre Earlsfort Terrace  
Dublin 2  
Tel: +353 1 6764971  
Fax: +353 1 6762517  
Website: [www.imb.ie](http://www.imb.ie)  
e-mail: [imbpharmacovigilance@imb.ie](mailto:imbpharmacovigilance@imb.ie)

### **5. How to store Cipramil**

Keep out of the sight and reach of children.  
Do not store above 25°C.  
Do not use Cipramil after the expiry date, which is stated on the label or carton. The expiry date refers to the last day of that month.  
Unused tablets or tablets older than the expiry date should be returned to a pharmacy for safe disposal.  
If your tablets appear to be discoloured, damaged or show any other signs of deterioration, please return to your pharmacist who will advise you.  
Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

### **6. Contents of the pack and other information**

#### **What Cipramil contains**

The active substance is citalopram (as hydrobromide).  
Each film-coated tablet contains 20mg citalopram (as citalopram hydrobromide).

The other ingredients in the tablets are: maize starch, lactose monohydrate, microcrystalline cellulose, crospovidone, glycerol 85%, croscarmellose sodium, magnesium stearate, hypromellose 5, macrogol 400 and titanium dioxide (E171).

#### **What Cipramil looks like and contents of the pack**

Cipramil are white, oval tablets, deeply scored on one face with the letters 'C' and 'N' on either side of the score and plain on the reverse.

Cipramil is available in blister packs of 28 tablets.

#### **Manufacturer**

Cipramil is manufactured by: H. Lundbeck A/S, Ottiliavej 9, DK-2500 Copenhagen, Denmark.

Procured from within the EU by the Parallel Product Authorisation holder: Imbat Ltd, Unit L2, North Ring Business Park, Santry, Dublin 9.

Repackaged by: Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, South Yorkshire, DN3 1QR, UK.

Distributed by: Eurodrug Ltd, Santry, Dublin 9.

PPA No: 1151/49/1

Leaflet revision and issue date (ref): 15.01.14

Cipramil® is a registered trademark of H. Lundbeck A/S.