

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### NEBILET® 5mg tablets nebivolol (as hydrochloride)

#### Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet:

1. What NEBILET is and what it is used for
2. What you need to know before you take NEBILET
3. How to take NEBILET
4. Possible side effects
5. How to store NEBILET
6. Contents of the pack and other information

#### 1. What NEBILET is and what it is used for

NEBILET contains nebivolol, a cardiovascular drug belonging to the group of selective beta-blocking agents (i.e. with a selective action on the cardiovascular system). It prevents increased heart rate, controls heart pumping strength. It also exerts a dilating action on blood vessels, which contributes as well to lower blood pressure. It is used to treat raised blood pressure (hypertension). NEBILET is also used to treat mild and moderate chronic heart failure in patients aged 70 or over, in addition to other therapies.

#### 2. What you need to know before you take NEBILET

##### Do not take NEBILET

- if you are allergic to nebivolol or any of the other ingredients of this medicine (listed in section 6).
- if you have one or more of the following disorders:
  - low blood pressure
  - serious circulation problems in the arms or legs
  - very slow heartbeat (less than 60 beats per minute)
  - certain other serious heart rhythm problems (e.g. 2<sup>nd</sup> and 3<sup>rd</sup> degree atrioventricular block, heart conduction disorders).
  - heart failure, which has just occurred or which has recently become worse, or you are receiving treatment for circulatory shock due to acute heart failure by intravenous drip feed to help your heart work
  - asthma or wheezing (now or in the past)
  - untreated phaeochromocytoma, a tumour located on top of the kidneys (in the adrenal glands)
  - liver function disorder
  - a metabolic disorder (metabolic acidosis), for example, diabetic ketoacidosis.

#### Warnings and precautions

Talk to your doctor or pharmacist before taking NEBILET

Inform your doctor if you have or develop one of the following problems:

- abnormally slow heartbeat
- a type of chest pain due to spontaneously occurring heart cramp called Prinzmetal angina
- untreated chronic heart failure
- 1st degree heart block (a kind of light heart conduction disorder that affects heart rhythm)
- poor circulation in the arms or legs, e.g. Raynaud's disease or syndrome, cramp-like pains when walking
- prolonged breathing problems
- diabetes: This medicine has no effect on blood sugar, but it could conceal the warning signs of a low sugar level (e.g. palpitations, fast heartbeat)
- overactive thyroid gland: This medicine may mask the signs of an abnormally fast heart rate due to this condition
- allergy: This medicine may intensify your reaction to pollen or other substances you are allergic to
- psoriasis (a skin disease - scaly pink patches) or if you have ever had psoriasis
- if you have to have surgery, always inform your anaesthetist that you are on NEBILET before being anaesthetised.

If you have serious kidney problems do not take NEBILET for heart failure and tell your doctor.

You will be regularly monitored at the beginning of your treatment for chronic heart failure by an experienced physician (see section 3).

This treatment should not be stopped abruptly unless clearly indicated and evaluated by your doctor (see section 3).

#### Children and adolescents

Because of the lack of data on the use of the product in children and adolescents, NEBILET is **not** recommended for use in them.

#### Other medicines and NEBILET

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Always tell your doctor if you are using or receiving any of the following medicines in addition to NEBILET:

- Medicines for controlling the blood pressure or medicines for heart problems (such as amiodarone, amlodipine, cibenzoline, clonidine, digoxin, diltiazem, disopyramide, felodipine, flecainide, guanfacin, hydroquinidine, lacidipine, lidocaine, methyl dopa, mexiletine, moxonidine, nicardipine, nifedipine, nimodipine, nitrendipine, propafenone, quinidine, rilmenidine, verapamil).
- Sedatives and therapies for psychosis (a mental illness) e.g. barbiturates (also used for epilepsy), phenothiazine (also used for vomiting and nausea) and thioridazine.
- Medicines for depression e.g. amitriptyline, paroxetine, fluoxetine.
- Medicines used for anaesthesia during an operation.
- Medicines for asthma, blocked nose or certain eye disorders such as glaucoma (increased pressure in the eye) or dilation (widening) of the pupil.
- Baclofen (an antispasmodic drug); Amifostine (a protective medicine used during cancer treatment)

All these drugs as well as nebivolol may influence the blood pressure and/or heart function.

- Medicines for treating excessive stomach acid or ulcers (antacid drug), e.g. cimetidine: you should take NEBILET during a meal and the antacid drug between meals.

#### NEBILET with food and drink

Please refer to section 3.

#### Pregnancy and breast-feeding

NEBILET should not be used during pregnancy, unless clearly necessary.

It is not recommended for use while breast-feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

#### Driving and using machines

This medicine may cause dizziness or fatigue. If affected, **do not** drive or operate machinery.

#### Important information about some of the ingredients of NEBILET

This product contains **lactose**. If you have been told by your doctor that you have an intolerance to some sugars, **contact your doctor before** taking this medicine.

### 3. How to take NEBILET

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

NEBILET may be taken before, during or after the meal, but, alternatively, you can take it independently of meals. The tablet is best taken with some water.

#### Treatment of raised blood pressure (hypertension)

- The usual dose is 1 tablet per day. The dose should be taken preferably at the same time of the day.
- Elderly patients and patients with a kidney disorder will usually start with ½ (half) tablet daily.
- The therapeutic effect on blood pressure becomes evident after 1-2 weeks of treatment. Occasionally, the optimal effect is reached only after 4 weeks.

#### Treatment of chronic heart failure

- Your treatment will be started and closely supervised by an experienced physician.
- Your doctor will start your treatment with ¼ (quarter) tablet per day. This may be increased after 1-2 weeks to ½ (half) tablet per day, then to 1 tablet per day and then to 2 tablets per day until the correct dose is reached for you. Your doctor will prescribe the dose that is right for you at each step and you should closely follow his/her instructions.
- The maximum recommended dose is 2 tablets (10mg) a day.
- You will need to be under the close supervision for 2 hours by an experienced physician when you start treatment and every time your dose is increased.
- Your doctor may reduce your dose if necessary.

You should **not stop treatment abruptly** as this can make your heart failure worse.


- Patients with serious kidney problems should not take this medicine.
- Take your medicine once daily, preferably at about the same time of day.

If you have been told by your doctor to take ¼ (quarter) or ½ (half) tablet daily, please refer to the instructions on how to break NEBILET 5mg cross-scored tablets.

- Place the tablets onto a flat, hard surface (e.g. a table or worktop), with the cross score facing up.

- Break the tablet by pushing it with the index fingers of both hands placed along one breakmark (Diagrams 1 and 2).
- Tablet quarters are obtained by breaking the halves in the same way (Diagrams 3 and 4).

**Diagrams 1 and 2:** Easy breaking of the Nebivolol 5mg cross-scored tablet in half.



**Diagrams 3 and 4:** Easy breaking of half of the Nebivolol 5mg cross-scored tablet into quarters.



- Your doctor may decide to combine NEBILET tablets with other medicines to treat your condition.
- Do not use in children or adolescents.

### If you take more NEBILET than you should

If you accidentally take an overdose of this medicine, tell your doctor or pharmacist **immediately**. The most frequent symptoms and signs of a NEBILET overdose are very slow heart beat (bradycardia), low blood pressure with possible fainting (hypotension), breathlessness such as in asthma (bronchospasm), and acute heart failure. You can take activated charcoal (which is available at your pharmacy) while you wait for the arrival of the doctor.

### If you forget to take NEBILET

If you forget a dose of NEBILET, but remember a little later on that you should have taken it, take that day's dose as usual. However, if a long delay has occurred (e.g. several hours), so that the next due dose is near, skip the forgotten dose and take the next, scheduled, **normal dose** at the usual time. Do not take a double dose. Repeated skipping, however, should be avoided.

### If you stop taking NEBILET

You should always consult with your doctor before stopping NEBILET treatment, whether you are taking it for high blood pressure or chronic heart failure.

You should not stop NEBILET treatment abruptly as this can temporarily make your heart failure worse. If it is necessary to stop NEBILET treatment for chronic heart failure, the daily dose should be decreased gradually, by halving the dose, at weekly intervals. If you have any further questions on the use of this product, ask your doctor or pharmacist.

## 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

When NEBILET is **used for the treatment of raised blood pressure**, the possible side effects are:

### Common side effects

**(may affect up to 1 in 10 people):**

- headache
- dizziness
- tiredness
- an unusual itching or tingling feeling

- diarrhoea
- constipation
- nausea
- shortness of breath
- swollen hands or feet.

### Uncommon side effects

**(may affect up to 1 in 100 people):**

- slow heartbeat or other heart complaints
- low blood pressure
- cramp-like leg pains on walking
- abnormal vision
- impotence
- feelings of depression
- digestive difficulties (dyspepsia), gas in stomach or bowel, vomiting
- skin rash, itchiness
- breathlessness such as in asthma, due to sudden cramps in the muscles around the airways (bronchospasm)
- nightmares.

### Very rare side effects

**(may affect up to 1 in 10,000 people):**

- fainting
- worsening of psoriasis (a skin disease - scaly pink patches).

The following side effects have been reported only in some isolated cases during NEBILET treatment:

- whole-body allergic reactions, with generalised skin eruption (hypersensitivity reactions);
- rapid-onset swelling, especially around the lips, eyes, or of the tongue with possible sudden difficulty breathing (angioedema);
- kind of skin rash notable for pale red, raised, itchy bumps of allergic or non allergic causes (urticaria).

In a clinical study for **chronic heart failure**, the following side effects were seen:

### Very common side effects

**(may affect more than 1 in 10 people):**

- slow heart beat
- dizziness.

### Common side effects

**(may affect up to 1 in 10 people):**

- worsening of heart failure
- low blood pressure (such as feeling faint when getting up quickly)
- inability to tolerate this medicine
- a kind of light heart conduction disorder that affects heart rhythm (1st degree AV-block)
- swelling of the lower limbs (such as swollen ankles).

### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via  
 HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2  
 Tel: +353 1 6764971 Fax: +353 1 6762517  
 Website: www.hpra.ie E-mail: medsafety@hpra.ie

By reporting side effects, you can help provide more information on the safety of this medicine.

## 5. How to store NEBILET

Keep out of the sight and reach of children.

This medicinal product does not require any special storage precautions.

Do not take NEBILET after the expiry date (Exp) which is stated on the carton and blister pack. The expiry date refers to the last day of that month.

NEBILET should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment. If your tablets appear to be discoloured, damaged or show any other signs of deterioration, please return them to your pharmacist who will advise you further.

## 6. Contents of the pack and other information

### What NEBILET contains

The active substance is nebivolol.

Each tablet contains 5mg nebivolol (as nebivolol hydrochloride): 2.5mg of d-nebivolol and 2.5mg of l-nebivolol.

The other ingredients are: lactose monohydrate, polysorbate 80 (E433), hypromellose (E464), maize starch, croscarmellose sodium (E468), microcrystalline cellulose (E460), colloidal anhydrous silica (E551), magnesium stearate (E572).

### What NEBILET looks like and contents of the pack

NEBILET tablets are white, round, cross-scored on one side and plain on the other.

NEBILET is available in blister packs of 28 tablets.

Manufactured by: Berlin-Chemie AG, Glienicker Weg 125, 12489 Berlin, Germany or Menarini-Von Heyden GmbH, Leipziger Strasse 7-13, 01097 - Dresden, Germany or Qualiphar N.V., Rijksweg 9, 2880 Bornem, Belgium.

Procured from within the EU by the PPA holder: Imbat Ltd, Unit L2, North Ring Business Park, Santry, Dublin 9.

Repackaged by: Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, South Yorkshire, DN3 1QR, UK.

Distributed by: Eurodrug Ltd, Santry, Dublin 9.

PPA No: 1151/83/1

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Austria: Nomexor	Belgium: Hypoloc
Cyprus: Lobivon	Finland: Nebilet
France: Hypoloc	Germany: Hypoloc
Greece: Hypoloc	Ireland: Nebilet
Italy: Lobivon	Iceland: Nebilet
Luxembourg: Hypoloc	Portugal: Hypoloc
Spain: Silostar	The Netherlands: Hypoloc
United Kingdom: Nebilet	

Leaflet revision & issue date (ref): 13.05.15

NEBILET® is a registered trademark of JOHNSON & JOHNSON.

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### NEBILET® 5mg tablets nebivolol (as hydrochloride)

#### Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet:

1. What NEBILET is and what it is used for
2. What you need to know before you take NEBILET
3. How to take NEBILET
4. Possible side effects
5. How to store NEBILET
6. Contents of the pack and other information

#### 1. What NEBILET is and what it is used for

NEBILET contains nebivolol, a cardiovascular drug belonging to the group of selective beta-blocking agents (i.e. with a selective action on the cardiovascular system). It prevents increased heart rate, controls heart pumping strength. It also exerts a dilating action on blood vessels, which contributes as well to lower blood pressure. It is used to treat raised blood pressure (hypertension). NEBILET is also used to treat mild and moderate chronic heart failure in patients aged 70 or over, in addition to other therapies.

#### 2. What you need to know before you take NEBILET

##### Do not take NEBILET

- if you are allergic to nebivolol or any of the other ingredients of this medicine (listed in section 6).
- if you have one or more of the following disorders:
  - low blood pressure
  - serious circulation problems in the arms or legs
  - very slow heartbeat (less than 60 beats per minute)
  - certain other serious heart rhythm problems (e.g. 2<sup>nd</sup> and 3<sup>rd</sup> degree atrioventricular block, heart conduction disorders).
  - heart failure, which has just occurred or which has recently become worse, or you are receiving treatment for circulatory shock due to acute heart failure by intravenous drip feed to help your heart work
  - asthma or wheezing (now or in the past)
  - untreated phaeochromocytoma, a tumour located on top of the kidneys (in the adrenal glands)
  - liver function disorder
  - a metabolic disorder (metabolic acidosis), for example, diabetic ketoacidosis.

#### Warnings and precautions

Talk to your doctor or pharmacist before taking NEBILET

Inform your doctor if you have or develop one of the following problems:

- abnormally slow heartbeat
- a type of chest pain due to spontaneously occurring heart cramp called Prinzmetal angina
- untreated chronic heart failure
- 1st degree heart block (a kind of light heart conduction disorder that affects heart rhythm)
- poor circulation in the arms or legs, e.g. Raynaud's disease or syndrome, cramp-like pains when walking
- prolonged breathing problems
- diabetes: This medicine has no effect on blood sugar, but it could conceal the warning signs of a low sugar level (e.g. palpitations, fast heartbeat)
- overactive thyroid gland: This medicine may mask the signs of an abnormally fast heart rate due to this condition
- allergy: This medicine may intensify your reaction to pollen or other substances you are allergic to
- psoriasis (a skin disease - scaly pink patches) or if you have ever had psoriasis
- if you have to have surgery, always inform your anaesthetist that you are on NEBILET before being anaesthetised.

If you have serious kidney problems do not take NEBILET for heart failure and tell your doctor.

You will be regularly monitored at the beginning of your treatment for chronic heart failure by an experienced physician (see section 3).

This treatment should not be stopped abruptly unless clearly indicated and evaluated by your doctor (see section 3).

#### Children and adolescents

Because of the lack of data on the use of the product in children and adolescents, NEBILET is **not** recommended for use in them.

#### Other medicines and NEBILET

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Always tell your doctor if you are using or receiving any of the following medicines in addition to NEBILET:

- Medicines for controlling the blood pressure or medicines for heart problems (such as amiodarone, amlodipine, cibenzoline, clonidine, digoxin, diltiazem, disopyramide, felodipine, flecainide, guanfacin, hydroquinidine, lacidipine, lidocaine, methyl dopa, mexiletine, moxonidine, nicardipine, nifedipine, nimodipine, nitrendipine, propafenone, quinidine, rilmenidine, verapamil).
- Sedatives and therapies for psychosis (a mental illness) e.g. barbiturates (also used for epilepsy), phenothiazine (also used for vomiting and nausea) and thioridazine.
- Medicines for depression e.g. amitriptyline, paroxetine, fluoxetine.
- Medicines used for anaesthesia during an operation.
- Medicines for asthma, blocked nose or certain eye disorders such as glaucoma (increased pressure in the eye) or dilation (widening) of the pupil.
- Baclofen (an antispasmodic drug); Amifostine (a protective medicine used during cancer treatment)

All these drugs as well as nebivolol may influence the blood pressure and/or heart function.

- Medicines for treating excessive stomach acid or ulcers (antacid drug), e.g. cimetidine: you should take NEBILET during a meal and the antacid drug between meals.

#### NEBILET with food and drink

Please refer to section 3.

#### Pregnancy and breast-feeding

NEBILET should not be used during pregnancy, unless clearly necessary.

It is not recommended for use while breast-feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

#### Driving and using machines

This medicine may cause dizziness or fatigue. If affected, **do not** drive or operate machinery.

#### Important information about some of the ingredients of NEBILET

This product contains **lactose**. If you have been told by your doctor that you have an intolerance to some sugars, **contact your doctor before** taking this medicine.

### 3. How to take NEBILET

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

NEBILET may be taken before, during or after the meal, but, alternatively, you can take it independently of meals. The tablet is best taken with some water.

#### Treatment of raised blood pressure (hypertension)

- The usual dose is 1 tablet per day. The dose should be taken preferably at the same time of the day.
- Elderly patients and patients with a kidney disorder will usually start with ½ (half) tablet daily.
- The therapeutic effect on blood pressure becomes evident after 1-2 weeks of treatment. Occasionally, the optimal effect is reached only after 4 weeks.

#### Treatment of chronic heart failure

- Your treatment will be started and closely supervised by an experienced physician.
- Your doctor will start your treatment with ¼ (quarter) tablet per day. This may be increased after 1-2 weeks to ½ (half) tablet per day, then to 1 tablet per day and then to 2 tablets per day until the correct dose is reached for you. Your doctor will prescribe the dose that is right for you at each step and you should closely follow his/her instructions.
- The maximum recommended dose is 2 tablets (10mg) a day.
- You will need to be under the close supervision for 2 hours by an experienced physician when you start treatment and every time your dose is increased.
- Your doctor may reduce your dose if necessary.

You should **not stop treatment abruptly** as this can make your heart failure worse.


- Patients with serious kidney problems should not take this medicine.
- Take your medicine once daily, preferably at about the same time of day.

If you have been told by your doctor to take ¼ (quarter) or ½ (half) tablet daily, please refer to the instructions on how to break NEBILET 5mg cross-scored tablets.

- Place the tablets onto a flat, hard surface (e.g. a table or worktop), with the cross score facing up.

- Break the tablet by pushing it with the index fingers of both hands placed along one breakmark (Diagrams 1 and 2).
- Tablet quarters are obtained by breaking the halves in the same way (Diagrams 3 and 4).

**Diagrams 1 and 2:** Easy breaking of the Nebivolol 5mg cross-scored tablet in half.



**Diagrams 3 and 4:** Easy breaking of half of the Nebivolol 5mg cross-scored tablet into quarters.



- Your doctor may decide to combine NEBILET tablets with other medicines to treat your condition.
- Do not use in children or adolescents.

#### If you take more NEBILET than you should

If you accidentally take an overdose of this medicine, tell your doctor or pharmacist **immediately**. The most frequent symptoms and signs of a NEBILET overdose are very slow heart beat (bradycardia), low blood pressure with possible fainting (hypotension), breathlessness such as in asthma (bronchospasm), and acute heart failure. You can take activated charcoal (which is available at your pharmacy) while you wait for the arrival of the doctor.

#### If you forget to take NEBILET

If you forget a dose of NEBILET, but remember a little later on that you should have taken it, take that day's dose as usual. However, if a long delay has occurred (e.g. several hours), so that the next due dose is near, skip the forgotten dose and take the next, scheduled, **normal dose** at the usual time. Do not take a double dose. Repeated skipping, however, should be avoided.

#### If you stop taking NEBILET

You should always consult with your doctor before stopping NEBILET treatment, whether you are taking it for high blood pressure or chronic heart failure.

You should not stop NEBILET treatment abruptly as this can temporarily make your heart failure worse. If it is necessary to stop NEBILET treatment for chronic heart failure, the daily dose should be decreased gradually, by halving the dose, at weekly intervals. If you have any further questions on the use of this product, ask your doctor or pharmacist.

### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

When NEBILET is **used for the treatment of raised blood pressure**, the possible side effects are:

#### Common side effects

(may affect up to 1 in 10 people):

- headache
- dizziness
- tiredness
- an unusual itching or tingling feeling

- diarrhoea
- constipation
- nausea
- shortness of breath
- swollen hands or feet.

#### Uncommon side effects

(may affect up to 1 in 100 people):

- slow heartbeat or other heart complaints
- low blood pressure
- cramp-like leg pains on walking
- abnormal vision
- impotence
- feelings of depression
- digestive difficulties (dyspepsia), gas in stomach or bowel, vomiting
- skin rash, itchiness
- breathlessness such as in asthma, due to sudden cramps in the muscles around the airways (bronchospasm)
- nightmares.

#### Very rare side effects

(may affect up to 1 in 10,000 people):

- fainting
- worsening of psoriasis (a skin disease - scaly pink patches).

The following side effects have been reported only in some isolated cases during NEBILET treatment:

- whole-body allergic reactions, with generalised skin eruption (hypersensitivity reactions);
- rapid-onset swelling, especially around the lips, eyes, or of the tongue with possible sudden difficulty breathing (angioedema);
- kind of skin rash notable for pale red, raised, itchy bumps of allergic or non allergic causes (urticaria).

In a clinical study for **chronic heart failure**, the following side effects were seen:

#### Very common side effects

(may affect more than 1 in 10 people):

- slow heart beat
- dizziness.

#### Common side effects

(may affect up to 1 in 10 people):

- worsening of heart failure
- low blood pressure (such as feeling faint when getting up quickly)
- inability to tolerate this medicine
- a kind of light heart conduction disorder that affects heart rhythm (1st degree AV-block)
- swelling of the lower limbs (such as swollen ankles).

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRAs Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2  
Tel: +353 1 6764971 Fax: +353 1 6762517  
Website: www.hpra.ie E-mail: medsafety@hpra.ie

By reporting side effects, you can help provide more information on the safety of this medicine.

### 5. How to store NEBILET

Keep out of the sight and reach of children.

This medicinal product does not require any special storage precautions.

Do not take NEBILET after the expiry date (EXP) which is stated on the carton and blister pack. The expiry date usually refers to the last day of that month.

NEBILET should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment. If your tablets appear to be discoloured, damaged or show any other signs of deterioration, please return them to your pharmacist who will advise you further.

### 6. Contents of the pack and other information

#### What NEBILET contains

The active substance is nebivolol.

Each tablet contains 5mg nebivolol (as nebivolol hydrochloride): 2.5mg of d-nebivolol and 2.5mg of l-nebivolol.

The other ingredients are: lactose monohydrate, polysorbate 80, hypromellose, maize starch, croscarmellose sodium, microcrystalline cellulose, colloidal anhydrous silica, magnesium stearate.

#### What NEBILET looks like and contents of the pack

NEBILET tablets are white, round, cross-scored on one side and plain on the other.

NEBILET is available in blister packs of 28 tablets.

Manufactured by: Berlin-Chemie AG, Glienicke Weg 125, 12489 Berlin, Germany.

Procured from within the EU by the PPA holder: Imbat Ltd, Unit L2, North Ring Business Park, Santry, Dublin 9.

Repackaged by: Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, South Yorkshire, DN3 1QR, UK.

Distributed by: Eurodrug Ltd, Santry, Dublin 9.

PPA No: 1151/83/1

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Austria: Nomexor	Belgium: Hypoloc
Cyprus: Lobivon	Finland: Nebilet
France: Hypoloc	Germany: Hypoloc
Greece: Hypoloc	Ireland: Nebilet
Italy: Lobivon	Iceland: Nebilet
Luxembourg: Hypoloc	Portugal: Hypoloc
Spain: Silostar	The Netherlands: Hypoloc
United Kingdom: Nebilet	

Leaflet revision & issue date (ref): 13.05.15

NEBILET® is a registered trademark of JOHNSON & JOHNSON.

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### NEBILET® 5mg tablets nebivolol (as hydrochloride)

#### Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet:

1. What NEBILET is and what it is used for
2. What you need to know before you take NEBILET
3. How to take NEBILET
4. Possible side effects
5. How to store NEBILET
6. Contents of the pack and other information

#### 1. What NEBILET is and what it is used for

NEBILET contains nebivolol, a cardiovascular drug belonging to the group of selective beta-blocking agents (i.e. with a selective action on the cardiovascular system). It prevents increased heart rate, controls heart pumping strength. It also exerts a dilating action on blood vessels, which contributes as well to lower blood pressure. It is used to treat raised blood pressure (hypertension). NEBILET is also used to treat mild and moderate chronic heart failure in patients aged 70 or over, in addition to other therapies.

#### 2. What you need to know before you take NEBILET

##### Do not take NEBILET

- if you are allergic to nebivolol or any of the other ingredients of this medicine (listed in section 6).
- if you have one or more of the following disorders:
  - low blood pressure
  - serious circulation problems in the arms or legs
  - very slow heartbeat (less than 60 beats per minute)
  - certain other serious heart rhythm problems (e.g. 2<sup>nd</sup> and 3<sup>rd</sup> degree atrioventricular block, heart conduction disorders).
  - heart failure, which has just occurred or which has recently become worse, or you are receiving treatment for circulatory shock due to acute heart failure by intravenous drip feed to help your heart work
  - asthma or wheezing (now or in the past)
  - untreated phaeochromocytoma, a tumour located on top of the kidneys (in the adrenal glands)
  - liver function disorder
  - a metabolic disorder (metabolic acidosis), for example, diabetic ketoacidosis.

#### Warnings and precautions

Talk to your doctor or pharmacist before taking NEBILET

Inform your doctor if you have or develop one of the following problems:

- abnormally slow heartbeat
- a type of chest pain due to spontaneously occurring heart cramp called Prinzmetal angina
- untreated chronic heart failure
- 1st degree heart block (a kind of light heart conduction disorder that affects heart rhythm)
- poor circulation in the arms or legs, e.g. Raynaud's disease or syndrome, cramp-like pains when walking
- prolonged breathing problems
- diabetes: This medicine has no effect on blood sugar, but it could conceal the warning signs of a low sugar level (e.g. palpitations, fast heartbeat)
- overactive thyroid gland: This medicine may mask the signs of an abnormally fast heart rate due to this condition
- allergy: This medicine may intensify your reaction to pollen or other substances you are allergic to
- psoriasis (a skin disease - scaly pink patches) or if you have ever had psoriasis
- if you have to have surgery, always inform your anaesthetist that you are on NEBILET before being anaesthetised.

If you have serious kidney problems do not take NEBILET for heart failure and tell your doctor.

You will be regularly monitored at the beginning of your treatment for chronic heart failure by an experienced physician (see section 3).

This treatment should not be stopped abruptly unless clearly indicated and evaluated by your doctor (see section 3).

#### Children and adolescents

Because of the lack of data on the use of the product in children and adolescents, NEBILET is **not** recommended for use in them.

#### Other medicines and NEBILET

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Always tell your doctor if you are using or receiving any of the following medicines in addition to NEBILET:

- Medicines for controlling the blood pressure or medicines for heart problems (such as amiodarone, amlodipine, cibenzoline, clonidine, digoxin, diltiazem, disopyramide, felodipine, flecainide, guanfacin, hydroquinidine, lacidipine, lidocaine, methyl dopa, mexiletine, moxonidine, nicardipine, nifedipine, nimodipine, nitrendipine, propafenone, quinidine, rilmenidine, verapamil).
- Sedatives and therapies for psychosis (a mental illness) e.g. barbiturates (also used for epilepsy), phenothiazine (also used for vomiting and nausea) and thioridazine.
- Medicines for depression e.g. amitriptyline, paroxetine, fluoxetine.
- Medicines used for anaesthesia during an operation.
- Medicines for asthma, blocked nose or certain eye disorders such as glaucoma (increased pressure in the eye) or dilation (widening) of the pupil.
- Baclofen (an antispasmodic drug); Amifostine (a protective medicine used during cancer treatment)

All these drugs as well as nebivolol may influence the blood pressure and/or heart function.

- Medicines for treating excessive stomach acid or ulcers (antacid drug), e.g. cimetidine: you should take NEBILET during a meal and the antacid drug between meals.

#### NEBILET with food and drink

Please refer to section 3.

#### Pregnancy and breast-feeding

NEBILET should not be used during pregnancy, unless clearly necessary.

It is not recommended for use while breast-feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

#### Driving and using machines

This medicine may cause dizziness or fatigue. If affected, **do not** drive or operate machinery.

#### Important information about some of the ingredients of NEBILET

This product contains **lactose**. If you have been told by your doctor that you have an intolerance to some sugars, **contact your doctor before** taking this medicine.

### 3. How to take NEBILET

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

NEBILET may be taken before, during or after the meal, but, alternatively, you can take it independently of meals. The tablet is best taken with some water.

#### Treatment of raised blood pressure (hypertension)

- The usual dose is 1 tablet per day. The dose should be taken preferably at the same time of the day.
- Elderly patients and patients with a kidney disorder will usually start with ½ (half) tablet daily.
- The therapeutic effect on blood pressure becomes evident after 1-2 weeks of treatment. Occasionally, the optimal effect is reached only after 4 weeks.

#### Treatment of chronic heart failure

- Your treatment will be started and closely supervised by an experienced physician.
- Your doctor will start your treatment with ¼ (quarter) tablet per day. This may be increased after 1-2 weeks to ½ (half) tablet per day, then to 1 tablet per day and then to 2 tablets per day until the correct dose is reached for you. Your doctor will prescribe the dose that is right for you at each step and you should closely follow his/her instructions.
- The maximum recommended dose is 2 tablets (10mg) a day.
- You will need to be under the close supervision for 2 hours by an experienced physician when you start treatment and every time your dose is increased.
- Your doctor may reduce your dose if necessary.

You should **not stop treatment abruptly** as this can make your heart failure worse.


- Patients with serious kidney problems should not take this medicine.
- Take your medicine once daily, preferably at about the same time of day.

If you have been told by your doctor to take ¼ (quarter) or ½ (half) tablet daily, please refer to the instructions on how to break NEBILET 5mg cross-scored tablets.

- Place the tablets onto a flat, hard surface (e.g. a table or worktop), with the cross score facing up.

- Break the tablet by pushing it with the index fingers of both hands placed along one breakmark (Diagrams 1 and 2).
- Tablet quarters are obtained by breaking the halves in the same way (Diagrams 3 and 4).

**Diagrams 1 and 2:** Easy breaking of the Nebivolol 5mg cross-scored tablet in half.



**Diagrams 3 and 4:** Easy breaking of half of the Nebivolol 5mg cross-scored tablet into quarters.



- Your doctor may decide to combine NEBILET tablets with other medicines to treat your condition.
- Do not use in children or adolescents.

#### If you take more NEBILET than you should

If you accidentally take an overdose of this medicine, tell your doctor or pharmacist **immediately**. The most frequent symptoms and signs of a NEBILET overdose are very slow heart beat (bradycardia), low blood pressure with possible fainting (hypotension), breathlessness such as in asthma (bronchospasm), and acute heart failure. You can take activated charcoal (which is available at your pharmacy) while you wait for the arrival of the doctor.

#### If you forget to take NEBILET

If you forget a dose of NEBILET, but remember a little later on that you should have taken it, take that day's dose as usual. However, if a long delay has occurred (e.g. several hours), so that the next due dose is near, skip the forgotten dose and take the next, scheduled, **normal dose** at the usual time. Do not take a double dose. Repeated skipping, however, should be avoided.

#### If you stop taking NEBILET

You should always consult with your doctor before stopping NEBILET treatment, whether you are taking it for high blood pressure or chronic heart failure.

You should not stop NEBILET treatment abruptly as this can temporarily make your heart failure worse. If it is necessary to stop NEBILET treatment for chronic heart failure, the daily dose should be decreased gradually, by halving the dose, at weekly intervals. If you have any further questions on the use of this product, ask your doctor or pharmacist.

#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

When NEBILET is **used for the treatment of raised blood pressure**, the possible side effects are:

##### Common side effects

(may affect up to 1 in 10 people):

- headache
- dizziness
- tiredness
- an unusual itching or tingling feeling

- diarrhoea
- constipation
- nausea
- shortness of breath
- swollen hands or feet.

##### Uncommon side effects

(may affect up to 1 in 100 people):

- slow heartbeat or other heart complaints
- low blood pressure
- cramp-like leg pains on walking
- abnormal vision
- impotence
- feelings of depression
- digestive difficulties (dyspepsia), gas in stomach or bowel, vomiting
- skin rash, itchiness
- breathlessness such as in asthma, due to sudden cramps in the muscles around the airways (bronchospasm)
- nightmares.

##### Very rare side effects

(may affect up to 1 in 10,000 people):

- fainting
- worsening of psoriasis (a skin disease - scaly pink patches).

The following side effects have been reported only in some isolated cases during NEBILET treatment:

- whole-body allergic reactions, with generalised skin eruption (hypersensitivity reactions);
- rapid-onset swelling, especially around the lips, eyes, or of the tongue with possible sudden difficulty breathing (angioedema);
- kind of skin rash notable for pale red, raised, itchy bumps of allergic or non allergic causes (urticaria).

In a clinical study for **chronic heart failure**, the following side effects were seen:

##### Very common side effects

(may affect more than 1 in 10 people):

- slow heart beat
- dizziness.

##### Common side effects

(may affect up to 1 in 10 people):

- worsening of heart failure
- low blood pressure (such as feeling faint when getting up quickly)
- inability to tolerate this medicine
- a kind of light heart conduction disorder that affects heart rhythm (1st degree AV-block)
- swelling of the lower limbs (such as swollen ankles).

##### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRa Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2  
Tel: +353 1 6764971 Fax: +353 1 6762517  
Website: www.hpra.ie E-mail: medsafety@hpra.ie

By reporting side effects, you can help provide more information on the safety of this medicine.

#### 5. How to store NEBILET

Keep out of the sight and reach of children.

This medicinal product does not require any special storage precautions.

Do not take NEBILET after the expiry date (Exp) which is stated on the carton and blister pack. The expiry date refers to the last day of that month.

NEBILET should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

If your tablets appear to be discoloured, damaged or show any other signs of deterioration, please return them to your pharmacist who will advise you further.

#### 6. Contents of the pack and other information

##### What NEBILET contains

The active substance is Nebivolol 5mg.

Each tablet contains 5mg nebivolol (as nebivolol hydrochloride): 2.5mg of d-nebivolol and 2.5mg of l-nebivolol.

The other ingredients are: lactose monohydrate, polysorbate 80 (E433), hypromellose (E464), maize starch, croscarmellose sodium (E468), microcrystalline cellulose (E460), colloidal anhydrous silica (E551), magnesium stearate (E572).

##### What NEBILET looks like and contents of the pack

NEBILET tablets are white, round, cross-scored on one side and plain on the other.

NEBILET is available in blister packs of 28 tablets.

Manufactured by: Berlin Chemie AG, Glienicke Weg 125, D-12489 Berlin, Germany.

Procured from within the EU by the PPA holder: Imbat Ltd, Unit L2, North Ring Business Park, Santry, Dublin 9.

Repackaged by: Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, South Yorkshire, DN3 1QR, UK.

Distributed by: Eurodrug Ltd, Santry, Dublin 9.

PPA No: 1151/83/1

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Austria: Nomexor	Belgium: Hypoloc
Cyprus: Lobivon	Finland: Nebilet
France: Hypoloc	Germany: Hypoloc
Greece: Hypoloc	Ireland: Nebilet
Italy: Lobivon	Iceland: Nebilet
Luxembourg: Hypoloc	Portugal: Hypoloc
Spain: Silostar	The Netherlands: Hypoloc
United Kingdom: Nebilet	

Leaflet revision & issue date (ref): 13.05.15

NEBILET® is a registered trademark of JOHNSON & JOHNSON.

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### NEBILET® 5mg tablets nebivolol (as hydrochloride)

#### Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet:

- What NEBILET is and what it is used for
- What you need to know before you take NEBILET
- How to take NEBILET
- Possible side effects
- How to store NEBILET
- Contents of the pack and other information

#### 1. What NEBILET is and what it is used for

NEBILET contains nebivolol, a cardiovascular drug belonging to the group of selective beta-blocking agents (i.e. with a selective action on the cardiovascular system). It prevents increased heart rate, controls heart pumping strength. It also exerts a dilating action on blood vessels, which contributes as well to lower blood pressure. It is used to treat raised blood pressure (hypertension). NEBILET is also used to treat mild and moderate chronic heart failure in patients aged 70 or over, in addition to other therapies.

#### 2. What you need to know before you take NEBILET

##### Do not take NEBILET

- if you are allergic to nebivolol or any of the other ingredients of this medicine (listed in section 6).
- if you have one or more of the following disorders:
  - low blood pressure
  - serious circulation problems in the arms or legs
  - very slow heartbeat (less than 60 beats per minute)
  - certain other serious heart rhythm problems (e.g. 2<sup>nd</sup> and 3<sup>rd</sup> degree atrioventricular block, heart conduction disorders).
  - heart failure, which has just occurred or which has recently become worse, or you are receiving treatment for circulatory shock due to acute heart failure by intravenous drip feed to help your heart work
  - asthma or wheezing (now or in the past)
  - untreated phaeochromocytoma, a tumour located on top of the kidneys (in the adrenal glands)
  - liver function disorder
  - a metabolic disorder (metabolic acidosis), for example, diabetic ketoacidosis.

#### Warnings and precautions

Talk to your doctor or pharmacist before taking NEBILET

Inform your doctor if you have or develop one of the following problems:

- abnormally slow heartbeat
- a type of chest pain due to spontaneously occurring heart cramp called Prinzmetal angina
- untreated chronic heart failure
- 1st degree heart block (a kind of light heart conduction disorder that affects heart rhythm)
- poor circulation in the arms or legs, e.g. Raynaud's disease or syndrome, cramp-like pains when walking
- prolonged breathing problems
- diabetes: This medicine has no effect on blood sugar, but it could conceal the warning signs of a low sugar level (e.g. palpitations, fast heartbeat)
- overactive thyroid gland: This medicine may mask the signs of an abnormally fast heart rate due to this condition
- allergy: This medicine may intensify your reaction to pollen or other substances you are allergic to
- psoriasis (a skin disease - scaly pink patches) or if you have ever had psoriasis
- if you have to have surgery, always inform your anaesthetist that you are on NEBILET before being anaesthetised.

If you have serious kidney problems do not take NEBILET for heart failure and tell your doctor.

You will be regularly monitored at the beginning of your treatment for chronic heart failure by an experienced physician (see section 3).

This treatment should not be stopped abruptly unless clearly indicated and evaluated by your doctor (see section 3).

#### Children and adolescents

Because of the lack of data on the use of the product in children and adolescents, NEBILET is **not** recommended for use in them.

#### Other medicines and NEBILET

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Always tell your doctor if you are using or receiving any of the following medicines in addition to NEBILET:

- Medicines for controlling the blood pressure or medicines for heart problems (such as amiodarone, amlodipine, cibenzoline, clonidine, digoxin, diltiazem, disopyramide, felodipine, flecainide, guanfacin, hydroquinidine, lacidipine, lidocaine, methyl dopa, mexiletine, moxonidine, nicardipine, nifedipine, nimodipine, nitrendipine, propafenone, quinidine, rilmenidine, verapamil).
- Sedatives and therapies for psychosis (a mental illness) e.g. barbiturates (also used for epilepsy), phenothiazine (also used for vomiting and nausea) and thioridazine.
- Medicines for depression e.g. amitriptyline, paroxetine, fluoxetine.
- Medicines used for anaesthesia during an operation.
- Medicines for asthma, blocked nose or certain eye disorders such as glaucoma (increased pressure in the eye) or dilation (widening) of the pupil.
- Baclofen (an antispasmodic drug); Amifostine (a protective medicine used during cancer treatment)

All these drugs as well as nebivolol may influence the blood pressure and/or heart function.

- Medicines for treating excessive stomach acid or ulcers (antacid drug), e.g. cimetidine: you should take NEBILET during a meal and the antacid drug between meals.

#### NEBILET with food and drink

Please refer to section 3.

#### Pregnancy and breast-feeding

NEBILET should not be used during pregnancy, unless clearly necessary.

It is not recommended for use while breast-feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

#### Driving and using machines

This medicine may cause dizziness or fatigue. If affected, **do not** drive or operate machinery.

#### Important information about some of the ingredients of NEBILET

This product contains **lactose**. If you have been told by your doctor that you have an intolerance to some sugars, **contact your doctor before** taking this medicine.

#### 3. How to take NEBILET

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

NEBILET may be taken before, during or after the meal, but, alternatively, you can take it independently of meals. The tablet is best taken with some water.

#### Treatment of raised blood pressure (hypertension)

- The usual dose is 1 tablet per day. The dose should be taken preferably at the same time of the day.
- Elderly patients and patients with a kidney disorder will usually start with ½ (half) tablet daily.
- The therapeutic effect on blood pressure becomes evident after 1-2 weeks of treatment. Occasionally, the optimal effect is reached only after 4 weeks.

#### Treatment of chronic heart failure

- Your treatment will be started and closely supervised by an experienced physician.
- Your doctor will start your treatment with ¼ (quarter) tablet per day. This may be increased after 1-2 weeks to ½ (half) tablet per day, then to 1 tablet per day and then to 2 tablets per day until the correct dose is reached for you. Your doctor will prescribe the dose that is right for you at each step and you should closely follow his/her instructions.
- The maximum recommended dose is 2 tablets (10mg) a day.
- You will need to be under the close supervision for 2 hours by an experienced physician when you start treatment and every time your dose is increased.
- Your doctor may reduce your dose if necessary.

You should **not stop treatment abruptly** as this can make your heart failure worse.


- Patients with serious kidney problems should not take this medicine.
- Take your medicine once daily, preferably at about the same time of day.

If you have been told by your doctor to take ¼ (quarter) or ½ (half) tablet daily, please refer to the instructions on how to break NEBILET 5mg cross-scored tablets.

- Place the tablets onto a flat, hard surface (e.g. a table or worktop), with the cross score facing up.

- Break the tablet by pushing it with the index fingers of both hands placed along one breakmark (Diagrams 1 and 2).
- Tablet quarters are obtained by breaking the halves in the same way (Diagrams 3 and 4).

**Diagrams 1 and 2:** Easy breaking of the Nebivolol 5mg cross-scored tablet in half.



**Diagrams 3 and 4:** Easy breaking of half of the Nebivolol 5mg cross-scored tablet into quarters.



- Your doctor may decide to combine NEBILET tablets with other medicines to treat your condition.
- Do not use in children or adolescents.

### If you take more NEBILET than you should

If you accidentally take an overdose of this medicine, tell your doctor or pharmacist **immediately**. The most frequent symptoms and signs of a NEBILET overdose are very slow heart beat (bradycardia), low blood pressure with possible fainting (hypotension), breathlessness such as in asthma (bronchospasm), and acute heart failure. You can take activated charcoal (which is available at your pharmacy) while you wait for the arrival of the doctor.

### If you forget to take NEBILET

If you forget a dose of NEBILET, but remember a little later on that you should have taken it, take that day's dose as usual. However, if a long delay has occurred (e.g. several hours), so that the next due dose is near, skip the forgotten dose and take the next, scheduled, **normal dose** at the usual time. Do not take a double dose. Repeated skipping, however, should be avoided.

### If you stop taking NEBILET

You should always consult with your doctor before stopping NEBILET treatment, whether you are taking it for high blood pressure or chronic heart failure.

You should not stop NEBILET treatment abruptly as this can temporarily make your heart failure worse. If it is necessary to stop NEBILET treatment for chronic heart failure, the daily dose should be decreased gradually, by halving the dose, at weekly intervals. If you have any further questions on the use of this product, ask your doctor or pharmacist.

## 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

When NEBILET is **used for the treatment of raised blood pressure**, the possible side effects are:

### Common side effects

(may affect up to 1 in 10 people):

- headache
- dizziness
- tiredness
- an unusual itching or tingling feeling

- diarrhoea
- constipation
- nausea
- shortness of breath
- swollen hands or feet.

### Uncommon side effects

(may affect up to 1 in 100 people):

- slow heartbeat or other heart complaints
- low blood pressure
- cramp-like leg pains on walking
- abnormal vision
- impotence
- feelings of depression
- digestive difficulties (dyspepsia), gas in stomach or bowel, vomiting
- skin rash, itchiness
- breathlessness such as in asthma, due to sudden cramps in the muscles around the airways (bronchospasm)
- nightmares.

### Very rare side effects

(may affect up to 1 in 10,000 people):

- fainting
- worsening of psoriasis (a skin disease - scaly pink patches).

The following side effects have been reported only in some isolated cases during NEBILET treatment:

- whole-body allergic reactions, with generalised skin eruption (hypersensitivity reactions);
- rapid-onset swelling, especially around the lips, eyes, or of the tongue with possible sudden difficulty breathing (angioedema);
- kind of skin rash notable for pale red, raised, itchy bumps of allergic or non allergic causes (urticaria).

In a clinical study for **chronic heart failure**, the following side effects were seen:

### Very common side effects

(may affect more than 1 in 10 people):

- slow heart beat
- dizziness.

### Common side effects

(may affect up to 1 in 10 people):

- worsening of heart failure
- low blood pressure (such as feeling faint when getting up quickly)
- inability to tolerate this medicine
- a kind of light heart conduction disorder that affects heart rhythm (1st degree AV-block)
- swelling of the lower limbs (such as swollen ankles).

### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRAs Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2  
Tel: +353 1 6764971 Fax: +353 1 6762517  
Website: www.hpra.ie E-mail: medsafety@hpra.ie

By reporting side effects, you can help provide more information on the safety of this medicine.

## 5. How to store NEBILET

Keep out of the sight and reach of children.

This medicinal product does not require any special storage precautions.

Do not take NEBILET after the expiry date (EXP) which is stated on the carton and blister pack. The expiry date usually refers to the last day of that month.

NEBILET should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment. If your tablets appear to be discoloured, damaged or show any other signs of deterioration, please return them to your pharmacist who will advise you further.

## 6. Contents of the pack and other information

### What NEBILET contains

The active substance is nebivolol.

Each tablet contains 5mg nebivolol (as nebivolol hydrochloride): 2.5mg of d-nebivolol and 2.5mg of l-nebivolol.

The other ingredients are: lactose monohydrate, polysorbate 80, hypromellose, maize starch, croscarmellose sodium, microcrystalline cellulose, colloidal anhydrous silica, magnesium stearate.

### What NEBILET looks like and contents of the pack

NEBILET tablets are white, round, cross-scored on one side and plain on the other.

NEBILET is available in blister packs of 28 tablets.

Manufactured by: Berlin-Chemie AG, Glienicker Weg 125, 12489 Berlin, Germany.

Procured from within the EU by the PPA holder: Imbat Ltd, Unit L2, North Ring Business Park, Santry, Dublin 9.

Repackaged by: Doncaster Pharmaceuticals Group Ltd, Kirk Sandall, Doncaster, South Yorkshire, DN3 1QR, UK.

Distributed by: Eurodrug Ltd, Santry, Dublin 9.

PPA No: 1151/83/1

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Austria: Nomexor	Belgium: Hypoloc
Cyprus: Lobivon	Finland: Nebilet
France: Hypoloc	Germany: Hypoloc
Greece: Hypoloc	Ireland: Nebilet
Italy: Lobivon	Iceland: Nebilet
Luxembourg: Hypoloc	Portugal: Hypoloc
Spain: Silostar	The Netherlands: Hypoloc
United Kingdom: Nebilet	

Leaflet revision & issue date (ref): 13.05.15

NEBILET® is a registered trademark of JOHNSON & JOHNSON.