

PACKAGE LEAFLET: INFORMATION FOR THE USER

Protium® 40 mg gastro-resistant tablets

Pantoprazole

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Protium is and what it is used for
2. What you need to know before you take Protium
3. How to take Protium
4. Possible side effects
5. How to store Protium
6. Contents of the pack and other information

1. What Protium is and what it is used for

Protium contains the active substance pantoprazole. Protium is a selective "proton pump inhibitor", a medicine which reduces the amount of acid produced in your stomach. It is used for treating acid-related diseases of the stomach and intestine.

Protium is used to treat adults and adolescents 12 years of age and above for

- Reflux oesophagitis. An inflammation of your oesophagus (the tube which connects your throat to your stomach) accompanied by the regurgitation of stomach acid.

Protium is used to treat adults for

- An infection with a bacterium called *Helicobacter pylori* in patients with duodenal ulcers and stomach ulcers in combination with two antibiotics (eradication therapy). The aim is to get rid of the bacteria and so reduce the likelihood of these ulcers returning.
- Stomach and duodenal ulcers.
- Zollinger-Ellison-Syndrome and other conditions producing too much acid in the stomach.

2. What you need to know before you take Protium

Do not take Protium

- If you are allergic to Pantoprazole or to any of the other ingredients of this medicine (listed in section 6)
- If you are allergic to medicines containing other proton pump inhibitors.

Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking Protium

- If you have severe liver problems. Please tell your doctor if you ever had problems with your liver in the past. He will check your liver enzymes more frequently, especially when you are taking Protium as a long-term treatment. In the case of a rise of liver enzymes the treatment should be stopped.
- If you have reduced body stores or risk factors for reduced vitamin B12 and receive long-term treatment with pantoprazole. As with all acid reducing agents, pantoprazole may lead to a reduced absorption of vitamin B12.
- If you are taking HIV protease inhibitors such as atazanavir (for the treatment of HIV- infection) at the same time as pantoprazole, ask your doctor for specific advice.
- Taking a proton pump inhibitor like pantoprazole, especially over a period of more than one year, may slightly increase your risk of fracture in the hip, wrist or spine. Tell your doctor if you have osteoporosis or if you are taking corticosteroids (which can increase the risk of osteoporosis).
- If you are on Protium for more than three months it is possible that the levels of magnesium in your blood may fall. Low levels of magnesium can be seen as fatigue, involuntary muscle contractions, disorientation, convulsions, dizziness, increased heart rate. If you get any of these symptoms, please tell your doctor promptly. Low levels of magnesium can also lead to a reduction in potassium or calcium levels in the blood. Your doctor may decide to perform regular blood tests to monitor your levels of magnesium.
- If you have ever had a skin reaction after treatment with a medicine similar to Protium that reduces stomach acid.
- If you get a rash on your skin, especially in areas exposed to the sun tell your doctor as soon as you can, as you may need to stop your treatment with Protium. Remember to also mention any other ill-effects like pain in your joints.

Tell your doctor immediately, before or after taking this medicine, if you notice any of the following symptoms, which could be a sign of another, more serious, disease:

- an unintentional loss of weight
- vomiting, particularly if repeated
- vomiting blood; this may appear as dark coffee grounds in your vomit
- you notice blood in your stools; which may be black or tarry in appearance
- difficulty in swallowing or pain when swallowing
- you look pale and feel weak (anaemia)
- chest pain
- stomach pain
- severe and/or persistent diarrhoea, because this medicine has been associated with a small increase in infectious diarrhoea.

Your doctor may decide that you need some tests to rule out malignant disease because pantoprazole also alleviates the symptoms of cancer and could cause delay in diagnosing it. If your symptoms continue in spite of your treatment, further investigations will be considered.

If you take Protium on a long-term basis (longer than 1 year) your doctor will probably keep you under regular surveillance. You should report any new and exceptional symptoms and circumstances whenever you see your doctor.

Children and adolescents

Protium is not recommended for use in children as it has not been proven to work in children below 12 years of age.

Other medicines and Protium

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

This is because Protium may influence the effectiveness of other medicines, so tell your doctor if you are taking:

- Medicines such as ketoconazole, itraconazole and posaconazole (used to treat fungal infections) or erlotinib (used for certain types of cancer) because Protium may stop these and other medicines from working properly.
- Warfarin and phenprocoumon, which affect the thickening, or thinning of the blood. You may need further checks.
- Medicines used to treat HIV- infection such as atazanavir.
- Methotrexate (used to treat rheumatoid arthritis, psoriasis, and cancer)- if you are taking methotrexate your doctor may temporarily stop your Protium treatment because pantoprazole can increase levels of methotrexate in the blood. Fluvoxamine (used to treat depression and other psychiatric diseases- if you are taking fluvoxamine your doctor may reduce the dose.
- Rifampicin (used to treat infections)
- St John's wort (Hypericum perforatum) (used to treat mild depression).

Pregnancy and breast-feeding

There are no adequate data from the use of pantoprazole in pregnant women. Excretion into human milk has been reported.

If you are pregnant or breast feeding, think you may be pregnant, or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

You should use this medicine only if your doctor considers the benefit for you greater than the potential risk for your unborn child or baby.

Driving and using machines

Protium has no or negligible influence on the ability to drive and use machines.

If you experience side effects like dizziness or disturbed vision, you should not drive or operate machines.

3. How to take Protium

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Methods of administration

Take the tablets 1 hour before a meal without chewing or breaking them and swallow them whole with some water.

The recommended dose is:

Adults and adolescents 12 years of age and above:

-To treat reflux oesophagitis

The usual dose is one tablet a day. Your doctor may tell you to increase to 2 tablets daily. The treatment period for reflux oesophagitis is usually between 4 and 8 weeks. Your doctor will tell you how long to take your medicine.

Adults:

For the treatment of an infection with a bacterium called Helicobacter pylori in patients with duodenal ulcers and stomach ulcers in combination with two antibiotics (Eradication therapy).

One tablet, two times a day plus two antibiotic tablets of either amoxicillin, clarithromycin and metronidazole (or tinidazole), each to be taken two times a day with your pantoprazole tablet. Take the first pantoprazole tablet 1 hour before breakfast and the second pantoprazole tablet 1 hour before your evening meal. Follow your doctor's instructions and make sure you read the package leaflets for these antibiotics. The usual treatment period is one to two weeks.

-For the treatment of stomach and duodenal ulcers.

The usual dose is one tablet a day. After consultation with your doctor, the dose may be doubled.

Your doctor will tell you how long to take your medicine. The treatment period for stomach ulcers is usually between 4 and 8 weeks. The treatment period for duodenal ulcers is usually between 2 and 4 weeks.

-For the long-term treatment of Zollinger-Ellison-Syndrome and of other conditions in which too much stomach acid is produced.

The recommended starting dose is usually two tablets a day.

Take the two tablets 1 hour before a meal. Your doctor may later adjust the dose, depending on the amount of stomach acid you produce. If prescribed more than two tablets a day, the tablets should be taken twice daily.

If your doctor prescribes a daily dose of more than four tablets a day, you will be told exactly when to stop taking the medicine.

Patients with kidney problems

- If you have kidney problems, you should not take Protium for eradication of *Helicobacter pylori*.

Patients with liver problems

- If you suffer from severe liver problems, you should not take more than one tablet 20 mg pantoprazole a day (for this purpose tablets containing 20 mg pantoprazole are available).

If you suffer from moderate or severe liver problems, you should not take Protium for eradication of *Helicobacter pylori*.

Use in children and adolescents

These tablets are not recommended for use in children below 12 years.

If you take more Protium than you should

Consult your doctor or pharmacist. There are no known symptoms of overdose.

If you forget to take Protium

Do not take a double dose to make up for the forgotten dose. Take your next, normal dose at the usual time.

If you stop taking Protium

Do not stop taking these tablets without first talking to your doctor or pharmacist.

If you have any further questions about the use of this product, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, Protium can cause side effects, although not everybody gets them.

If you get any of the following side effects, stop taking these tablets and tell your doctor immediately, or contact the casualty department at your nearest hospital:

- **Serious allergic reactions (frequency rare:** may affect up to 1 in 1,000 people): swelling of the tongue and/or throat, difficulty in swallowing, hives (nettle rash), difficulties in breathing, allergic facial swelling (Quincke's oedema/ angioedema), severe dizziness with very fast heartbeat and heavy sweating.

- **Serious skin conditions (frequency not known:** frequency cannot be estimated from the available data): blistering of the skin and rapid deterioration of your general condition, erosion (including slight bleeding) of eyes, nose, mouth/lips or genitals (Stevens-Johnson-Syndrome, Lyell-Syndrome, Erythema multiforme) and sensitivity to light.

- **Other serious conditions (frequency not known:** frequency cannot be estimated from the available data): yellowing of the skin or whites of the eyes (severe damage to liver cells, jaundice) or fever, rash, and enlarged kidneys sometimes with painful urination and lower back pain (serious inflammation of the kidneys), possibly leading to kidney failure.

Other side effects are:

- **Uncommon** (may affect up to 1 in 100 people)

Headache; dizziness; diarrhoea; feeling sick, vomiting; bloating and flatulence (wind); constipation; dry mouth; abdominal pain and discomfort; skin rash, exanthema, eruption; itching; feeling weak, exhausted or generally unwell; sleep disorders; fracture in the hip, wrist or spine.

- **Rare** (may affect up to 1 in 1,000 people)

Distortion or complete lack of the sense of taste; disturbances in vision such as blurred vision; hives; pain in the joints; muscle pains; weight changes; raised body temperature; high fever; swelling of the extremities (peripheral oedema); allergic reactions; depression; breast enlargement in males.

- **Very Rare** (may affect up to 1 in 10,000 people)

Disorientation.

- **Not known** (frequency cannot be estimated from the available data)

Hallucination, confusion (especially in patients with a history of these symptoms); decreased sodium level in blood, decreased magnesium level in blood (see section 2), feeling of tingling, pricking, pins and needles, burning sensations or numbness, rash, possibly with pain in the joints.

Side effects identified through blood tests:

- **Uncommon** (may affect up to 1 in 100 people)

an increase in liver enzymes.

- **Rare** (may affect up to 1 in 1,000 people)

an increase in bilirubin; increased fat levels in the blood; sharp drop in circulating granular white blood cells, associated with high fever

- **Very Rare** (may affect up to 1 in 10,000 people)

a reduction in the number of blood platelets, which may cause you to bleed or bruise more than normal; a reduction in the number of white blood cells, which may lead to more frequent infections; coexisting abnormal reduction in the number of red and white blood cells, as well as platelets.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Protium

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date, which is stated on the carton and the container after EXP.

The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Protium contains

- The active substance is pantoprazole. Each gastro-resistant tablet contains 40 mg of pantoprazole (as sodium sesquihydrate).

- The other ingredients are:

Core: sodium carbonate (anhydrous), mannitol (E421), crospovidone, povidone K90, calcium stearate.

Coating: hypromellose, povidone K25, titanium dioxide (E171), yellow iron oxide (E172), propylene glycol, methacrylic acid-ethyl acrylate copolymer (1:1), polysorbate 80, sodium laurilsulfate, triethyl citrate.

Printing ink: shellac, red, black and yellow iron oxide (E172), titanium dioxide (E171), antifoam DC 1510 (dimeticone emulsion) and ammonia solution (concentrated).

What Protium looks like and contents of the pack

Yellow, oval, biconvex gastro-resistant coated tablet imprinted with "P 40" on one side.

Packs: blister (ALU/ALU blister) without cardboard reinforcement

Protium is available in the following pack sizes:

Packs with 28 gastro-resistant tablets.

Product procured from within the EU by the Parallel Product Authorisation Holder:

IMED Healthcare Ltd, Unit 625 Kilshane Avenue, Northwest Business Park, Ballycoolin, Dublin 15, Ireland

Manufacturer:

Takeda GmbH, Production site Oranienburg, Lehnitzstrasse 70 – 98, 16515 Oranienburg, Germany

Repackaged by:

Cast Healthcare Ltd, Unit B6 The Business Centre, 5-7 Tobermore Road, Draperstown, Northern Ireland, BT45 7AG, UK.

This medicine is authorised in the Member States of the EEA under the following names:

Austria – Pantoloc 40mg – Filmtabletten, Zurcal 40mg - Filmtabletten

Belgium – Pantozol, Zurcale

Bulgaria- Controloc

Cyprus - Controloc

Czech Republic – Controloc 40mg

Denmark – Pantoloc

Estonia – Controloc 40mg

Finland – Somac 40mg

France – Eupantol 40mg, Inipomp 40mg, Pantec 40mg, Pantipp 40mg

Germany – Pantozol 40mg, Pantoprazol NYC 40mg, Rifun 40mg,mZurcal S 40mg

Greece – Controloc, Zurcazole

Hungary – Controloc 40mg

Ireland – Protium

Italy – Pantroc, Pantopan, Pantecta, Peptazol

Latvia – Controloc 40mg

Lithuania – Controloc 40mg

Luxembourg – Pantozol-40, Panto-Byk-40

Netherlands – Pantozol 40mg, Pantoprazol Nycomed 40mg

Norway - Somac

Poland – Controloc 40

Portugal – Pantoc 40mg,Zurcal 40mg, Apton 40mg, Pantoprazole ALTANA 40mg

Romania – Controloc 40mg

Slovakia – Controloc 40mg

Slovenia – Controloc 40mg

Spain – Pantecta 40mg blister, Anagastra 40mg Blister, Ulcotenal 40mg comprimidos gastrorresistentes Blister

Sweden – Pantoloc

United Kingdom – Protium 40mg

This leaflet was last approved in May 2016

PACKAGE LEAFLET: INFORMATION FOR THE USER

Protium® 40 mg gastro-resistant tablets

Pantoprazole

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Protium is and what it is used for
2. What you need to know before you take Protium
3. How to take Protium
4. Possible side effects
5. How to store Protium
6. Contents of the pack and other information

1. What Protium is and what it is used for

Protium contains the active substance pantoprazole. Protium is a selective "proton pump inhibitor", a medicine which reduces the amount of acid produced in your stomach. It is used for treating acid-related diseases of the stomach and intestine.

Protium is used to treat adults and adolescents 12 years of age and above for

- Reflux oesophagitis. An inflammation of your oesophagus (the tube which connects your throat to your stomach) accompanied by the regurgitation of stomach acid.

Protium is used to treat adults for

- An infection with a bacterium called *Helicobacter pylori* in patients with duodenal ulcers and stomach ulcers in combination with two antibiotics (eradication therapy). The aim is to get rid of the bacteria and so reduce the likelihood of these ulcers returning.
- Stomach and duodenal ulcers.
- Zollinger-Ellison-Syndrome and other conditions producing too much acid in the stomach.

2. What you need to know before you take Protium

Do not take Protium

- If you are allergic to Pantoprazole or to any of the other ingredients of this medicine (listed in section 6)
- If you are allergic to medicines containing other proton pump inhibitors.

Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking Protium

- If you have severe liver problems. Please tell your doctor if you ever had problems with your liver in the past. He will check your liver enzymes more frequently, especially when you are taking Protium as a long-term treatment. In the case of a rise of liver enzymes the treatment should be stopped.
- If you have reduced body stores or risk factors for reduced vitamin B12 and receive long-term treatment with pantoprazole. As with all acid reducing agents, pantoprazole may lead to a reduced absorption of vitamin B12.
- If you are taking HIV protease inhibitors such as atazanavir (for the treatment of HIV- infection) at the same time as pantoprazole, ask your doctor for specific advice.
- Taking a proton pump inhibitor like pantoprazole, especially over a period of more than one year, may slightly increase your risk of fracture in the hip, wrist or spine. Tell your doctor if you have osteoporosis or if you are taking corticosteroids (which can increase the risk of osteoporosis).
- If you are on Protium for more than three months it is possible that the levels of magnesium in your blood may fall. Low levels of magnesium can be seen as fatigue, involuntary muscle contractions, disorientation, convulsions, dizziness, increased heart rate. If you get any of these symptoms, please tell your doctor promptly. Low levels of magnesium can also lead to a reduction in potassium or calcium levels in the blood. Your doctor may decide to perform regular blood tests to monitor your levels of magnesium.
- If you have ever had a skin reaction after treatment with a medicine similar to Protium that reduces stomach acid.
- If you get a rash on your skin, especially in areas exposed to the sun tell your doctor as soon as you can, as you may need to stop your treatment with Protium. Remember to also mention any other ill-effects like pain in your joints.

Tell your doctor immediately, before or after taking this medicine, if you notice any of the following symptoms, which could be a sign of another, more serious, disease:

- an unintentional loss of weight
- vomiting, particularly if repeated
- vomiting blood; this may appear as dark coffee grounds in your vomit
- you notice blood in your stools; which may be black or tarry in appearance
- difficulty in swallowing or pain when swallowing
- you look pale and feel weak (anaemia)
- chest pain
- stomach pain
- severe and/or persistent diarrhoea, because this medicine has been associated with a small increase in infectious diarrhoea.

Your doctor may decide that you need some tests to rule out malignant disease because pantoprazole also alleviates the symptoms of cancer and could cause delay in diagnosing it. If your symptoms continue in spite of your treatment, further investigations will be considered.

If you take Protium on a long-term basis (longer than 1 year) your doctor will probably keep you under regular surveillance. You should report any new and exceptional symptoms and circumstances whenever you see your doctor.

Children and adolescents

Protium is not recommended for use in children as it has not been proven to work in children below 12 years of age.

Other medicines and Protium

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

This is because Protium may influence the effectiveness of other medicines, so tell your doctor if you are taking:

- Medicines such as ketoconazole, itraconazole and posaconazole (used to treat fungal infections) or erlotinib (used for certain types of cancer) because Protium may stop these and other medicines from working properly.
- Warfarin and phenprocoumon, which affect the thickening, or thinning of the blood. You may need further checks.
- Medicines used to treat HIV- infection such as atazanavir.
- Methotrexate (used to treat rheumatoid arthritis, psoriasis, and cancer)- if you are taking methotrexate your doctor may temporarily stop your Protium treatment because pantoprazole can increase levels of methotrexate in the blood. Fluvoxamine (used to treat depression and other psychiatric diseases- if you are taking fluvoxamine your doctor may reduce the dose.
- Rifampicin (used to treat infections)
- St John's wort (Hypericum perforatum) (used to treat mild depression).

Pregnancy and breast-feeding

There are no adequate data from the use of pantoprazole in pregnant women. Excretion into human milk has been reported.

If you are pregnant or breast feeding, think you may be pregnant, or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

You should use this medicine only if your doctor considers the benefit for you greater than the potential risk for your unborn child or baby.

Driving and using machines

Protium has no or negligible influence on the ability to drive and use machines.

If you experience side effects like dizziness or disturbed vision, you should not drive or operate machines.

3. How to take Protium

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Methods of administration

Take the tablets 1 hour before a meal without chewing or breaking them and swallow them whole with some water.

The recommended dose is:

Adults and adolescents 12 years of age and above:

-To treat reflux oesophagitis

The usual dose is one tablet a day. Your doctor may tell you to increase to 2 tablets daily. The treatment period for reflux oesophagitis is usually between 4 and 8 weeks. Your doctor will tell you how long to take your medicine.

Adults:

For the treatment of an infection with a bacterium called Helicobacter pylori in patients with duodenal ulcers and stomach ulcers in combination with two antibiotics (Eradication therapy).

One tablet, two times a day plus two antibiotic tablets of either amoxicillin, clarithromycin and metronidazole (or tinidazole), each to be taken two times a day with your pantoprazole tablet. Take the first pantoprazole tablet 1 hour before breakfast and the second pantoprazole tablet 1 hour before your evening meal. Follow your doctor's instructions and make sure you read the package leaflets for these antibiotics. The usual treatment period is one to two weeks.

-For the treatment of stomach and duodenal ulcers.

The usual dose is one tablet a day. After consultation with your doctor, the dose may be doubled.

Your doctor will tell you how long to take your medicine. The treatment period for stomach ulcers is usually between 4 and 8 weeks. The treatment period for duodenal ulcers is usually between 2 and 4 weeks.

-For the long-term treatment of Zollinger-Ellison-Syndrome and of other conditions in which too much stomach acid is produced.

The recommended starting dose is usually two tablets a day.

Take the two tablets 1 hour before a meal. Your doctor may later adjust the dose, depending on the amount of stomach acid you produce. If prescribed more than two tablets a day, the tablets should be taken twice daily.

If your doctor prescribes a daily dose of more than four tablets a day, you will be told exactly when to stop taking the medicine.

Patients with kidney problems

- If you have kidney problems, you should not take Protium for eradication of *Helicobacter pylori*.

Patients with liver problems

- If you suffer from severe liver problems, you should not take more than one tablet 20 mg pantoprazole a day (for this purpose tablets containing 20 mg pantoprazole are available).

If you suffer from moderate or severe liver problems, you should not take Protium for eradication of *Helicobacter pylori*.

Use in children and adolescents

These tablets are not recommended for use in children below 12 years.

If you take more Protium than you should

Consult your doctor or pharmacist. There are no known symptoms of overdose.

If you forget to take Protium

Do not take a double dose to make up for the forgotten dose. Take your next, normal dose at the usual time.

If you stop taking Protium

Do not stop taking these tablets without first talking to your doctor or pharmacist.

If you have any further questions about the use of this product, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, Protium can cause side effects, although not everybody gets them.

If you get any of the following side effects, stop taking these tablets and tell your doctor immediately, or contact the casualty department at your nearest hospital:

- **Serious allergic reactions (frequency rare:** may affect up to 1 in 1,000 people): swelling of the tongue and/or throat, difficulty in swallowing, hives (nettle rash), difficulties in breathing, allergic facial swelling (Quincke's oedema/ angioedema), severe dizziness with very fast heartbeat and heavy sweating.

- **Serious skin conditions (frequency not known:** frequency cannot be estimated from the available data): blistering of the skin and rapid deterioration of your general condition, erosion (including slight bleeding) of eyes, nose, mouth/lips or genitals (Stevens-Johnson-Syndrome, Lyell-Syndrome, Erythema multiforme) and sensitivity to light.

- **Other serious conditions (frequency not known:** frequency cannot be estimated from the available data): yellowing of the skin or whites of the eyes (severe damage to liver cells, jaundice) or fever, rash, and enlarged kidneys sometimes with painful urination and lower back pain (serious inflammation of the kidneys), possibly leading to kidney failure.

Other side effects are:

- **Uncommon** (may affect up to 1 in 100 people)

Headache; dizziness; diarrhoea; feeling sick, vomiting; bloating and flatulence (wind); constipation; dry mouth; abdominal pain and discomfort; skin rash, exanthema, eruption; itching; feeling weak, exhausted or generally unwell; sleep disorders; fracture in the hip, wrist or spine.

- **Rare** (may affect up to 1 in 1,000 people)

Distortion or complete lack of the sense of taste; disturbances in vision such as blurred vision; hives; pain in the joints; muscle pains; weight changes; raised body temperature; high fever; swelling of the extremities (peripheral oedema); allergic reactions; depression; breast enlargement in males.

- **Very Rare** (may affect up to 1 in 10,000 people)

Disorientation.

- **Not known** (frequency cannot be estimated from the available data)

Hallucination, confusion (especially in patients with a history of these symptoms); decreased sodium level in blood, decreased magnesium level in blood (see section 2), feeling of tingling, pricking, pins and needles, burning sensations or numbness, rash, possibly with pain in the joints.

Side effects identified through blood tests:

- **Uncommon** (may affect up to 1 in 100 people)

an increase in liver enzymes.

- **Rare** (may affect up to 1 in 1,000 people)

an increase in bilirubin; increased fat levels in the blood; sharp drop in circulating granular white blood cells, associated with high fever

- **Very Rare** (may affect up to 1 in 10,000 people)

a reduction in the number of blood platelets, which may cause you to bleed or bruise more than normal; a reduction in the number of white blood cells, which may lead to more frequent infections; coexisting abnormal reduction in the number of red and white blood cells, as well as platelets.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Protium

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date, which is stated on the carton and the container after EXP.

The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Protium contains

- The active substance is pantoprazole. Each gastro-resistant tablet contains 40 mg of pantoprazole (as sodium sesquihydrate).

- The other ingredients are:

Core: sodium carbonate (anhydrous), mannitol (E421), crospovidone, povidone K90, calcium stearate.

Coating: hypromellose, povidone K25, titanium dioxide (E171), yellow iron oxide (E172), propylene glycol, methacrylic acid-ethyl acrylate copolymer (1:1), polysorbate 80, sodium laurilsulfate, triethyl citrate.

Printing ink: shellac, red, black and yellow iron oxide (E172), titanium dioxide (E171), antifoam DC 1510 (dimeticone emulsion) and ammonia solution (concentrated).

What Protium looks like and contents of the pack

Yellow, oval, biconvex gastro-resistant coated tablet imprinted with "P 40" on one side.

Packs: blister (ALU/ALU blister) without cardboard reinforcement

Protium is available in the following pack sizes:

Packs with 28 gastro-resistant tablets.

Product procured from within the EU by the Parallel Product Authorisation Holder:

IMED Healthcare Ltd, Unit 625 Kilshane Avenue, Northwest Business Park, Ballycoolin, Dublin 15, Ireland

Manufacturer:

Takeda GmbH, Production site Oranienburg, Lehnitzstrasse 70 – 98, 16515 Oranienburg, Germany

Repackaged by:

IMED Pharma Ltd, Unit 625 Kilshane Avenue, Northwest Business Park, Ballycoolin, Dublin 15, Ireland.

This medicine is authorised in the Member States of the EEA under the following names:

Austria – Pantoloc 40mg – Filmtabletten, Zurcal 40mg - Filmtabletten

Belgium – Pantozol, Zurcale

Bulgaria- Controloc

Cyprus - Controloc

Czech Republic – Controloc 40mg

Denmark – Pantoloc

Estonia – Controloc 40mg

Finland – Somac 40mg

France – Eupantol 40mg, Inipomp 40mg, Pantec 40mg, Pantipp 40mg

Germany – Pantozol 40mg, Pantoprazol NYC 40mg, Rifun 40mg,mZurcal S 40mg

Greece – Controloc, Zurcazole

Hungary – Controloc 40mg

Ireland – Protium

Italy – Pantroc, Pantopan, Pantecta, Peptazol

Latvia – Controloc 40mg

Lithuania – Controloc 40mg

Luxembourg – Pantozol-40, Panto-Byk-40

Netherlands – Pantozol 40mg, Pantoprazol Nycomed 40mg

Norway - Somac

Poland – Controloc 40

Portugal – Pantoc 40mg,Zurcal 40mg, Apton 40mg, Pantoprazole ALTANA 40mg

Romania – Controloc 40mg

Slovakia – Controloc 40mg

Slovenia – Controloc 40mg

Spain – Pantecta 40mg blister, Anagastra 40mg Blister, Ulcotenal 40mg comprimidos gastrorresistentes

Blister

Sweden – Pantoloc

United Kingdom – Protium 40mg

This leaflet was last approved in May 2016