

## Package leaflet: Information for the patient

### Darunavir Rowex 75 mg Film-coated tablets Darunavir Rowex 600 mg Film-coated tablets

darunavir

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Darunavir Rowex is and what it is used for
2. What you need to know before you take Darunavir Rowex
3. How to take Darunavir Rowex
4. Possible side effects
5. How to store Darunavir Rowex
6. Contents of the pack and other information

#### **1. What Darunavir Rowex is and what it is used for**

##### **What is Darunavir Rowex**

Darunavir Rowex contains the active substance darunavir. Darunavir Rowex is an antiretroviral medicine used in the treatment of Human Immunodeficiency Virus (HIV) infection. It belongs to a group of medicines called protease inhibitors. Darunavir Rowex works by reducing the amount of HIV in your body. This will improve your immune system and reduces the risk of developing illnesses linked to HIV infection.

##### **What is it used for?**

Darunavir Rowex is used to treat adults and children of 3 years of age and above, and at least 15 kilogram body weight who are infected by HIV and who have already used other antiretroviral medicines.

Darunavir Rowex must be taken in combination with a low dose of ritonavir and other anti-HIV medicines. Your doctor will discuss with you which combination of medicines is best for you.

#### **2. What you need to know before you take Darunavir Rowex**

##### **Do not take Darunavir Rowex**

- if you are **allergic** to darunavir or any of the other ingredients of this medicine (listed in section 6) or to ritonavir.
- if you have **severe liver problems**. Ask your doctor if you are unsure about the severity of your liver disease. Some additional tests might be necessary.

##### **Do not combine Darunavir Rowex with any of the following medicines**

If you are taking any of these, ask your doctor about switching to another medicine.

Medicine	Purpose of the medicine
<i>Avanafil</i>	to treat erectile dysfunction

<i>Astemizole or terfenadine</i>	to treat allergy symptoms
<i>Triazolam and oral (taken by mouth) midazolam</i>	to help you sleep and/or relieve anxiety
<i>Cisapride</i>	to treat some stomach conditions
<i>Colchicine</i> (if you have kidney and/or liver problems)	to treat gout or familial Mediterranean fever
<i>Lurasidone, Pimozide, quetiapine or sertindole</i>	to treat psychiatric conditions
<i>Ergot alkaloids</i> like <i>ergotamine, dihydroergotamine, ergometrine and methylergonovine</i>	to treat migraine headaches
<i>Amiodarone, bepridil, dronedarone, quinidine, ranolazine</i>	to treat certain heart disorders e.g. abnormal heartbeat
<i>Lovastatin and simvastatin and lomitapide</i>	to lower cholesterol levels
<i>Rifampicin</i>	to treat some infections such as tuberculosis
The combination product <i>lopinavir/ritonavir</i>	this anti-HIV medicine belongs to the same class as Darunavir Rowex
<i>Elbasvir/grazoprevir</i>	to treat hepatitis C infection
<i>Alfuzosin</i>	to treat enlarged prostate
<i>Sildenafil</i>	to treat high blood pressure in the pulmonary circulation
<i>Dabigatran, Ticagrelor</i>	to help stop the clumping of platelets in the treatment of patients with a history of a heart attack

Do not combine Darunavir Rowex with products that contain St John's wort (*Hypericum perforatum*).

### Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking Darunavir Rowex.

Darunavir Rowex is not a cure for HIV infection. You can still pass on HIV when taking this medicine, although the risk is lowered by effective antiretroviral therapy. Discuss with your physician the precautions needed to avoid infecting other people.

People taking Darunavir Rowex may still develop infections or other illnesses associated with HIV infection. You must keep in regular contact with your doctor.

People taking Darunavir Rowex may develop a skin rash. Infrequently a rash may become severe or potentially life-threatening. Please contact your doctor whenever you develop a rash.

In patients taking Darunavir Rowex and raltegravir (for HIV infection), rashes (generally mild or moderate) may occur more frequently than in patients taking either medicine separately.

### Tell your doctor about your situation BEFORE and DURING your treatment

Make sure that you check the following points and tell your doctor if any of these apply to you.

- Tell your doctor if you have had **problems with your liver** before, including hepatitis B or C infection. Your doctor may evaluate how severe your liver disease is before deciding if you can take Darunavir Rowex.
- Tell your doctor if you have **diabetes**. Darunavir Rowex might increase sugar levels in the blood.
- Tell your doctor immediately if you notice any **symptoms of infection** (for example

enlarged lymph nodes and fever). In some patients with advanced HIV infection and a history of opportunistic infection, signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.

- In addition to the opportunistic infections, autoimmune disorders (a condition that occurs when the immune system attacks healthy body tissue) may also occur after you start taking medicines for the treatment of your HIV infection. Autoimmune disorders may occur many months after the start of treatment. If you notice any symptoms of infection or other symptoms such as muscle weakness, weakness beginning in the hands and feet and moving up towards the trunk of the body, palpitations, tremor or hyperactivity, please inform your doctor immediately to seek necessary treatment.
- Tell your doctor if you have **haemophilia**. Darunavir Rowex might increase the risk of bleeding.
- Tell your doctor if you are **allergic to sulphonamides** (e.g. used to treat certain infections).
- Tell your doctor if you notice any **musculoskeletal problems**. Some patients taking combination antiretroviral therapy may develop a bone disease called osteonecrosis (death of bone tissue caused by loss of blood supply to the bone). The length of combination antiretroviral therapy, corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index, among others, may be some of the many risk factors for developing this disease. Signs of osteonecrosis are joint stiffness, aches and pains (especially of the hip, knee and shoulder) and difficulty in movement. If you notice any of these symptoms please inform your doctor.

### **Elderly**

Darunavir Rowex has only been used in limited numbers of patients 65 years or older. If you belong to this age group, please discuss with your doctor if you can use Darunavir Rowex.

### **Children**

Darunavir Rowex is not for use in children younger than 3 years of age or weighing less than 40 kilograms.

### **Other medicines and Darunavir Rowex**

Tell your doctor or pharmacist if you are taking, or have recently taken any other medicines.

There are some medicines that **you must not combine** with Darunavir Rowex. These are mentioned above under the heading 'Do not combine Darunavir Rowex with any of the following medicines:'

In most cases, Darunavir Rowex can be combined with anti-HIV medicines belonging to another class [e.g. NRTIs (nucleoside reverse transcriptase inhibitors), NNRTIs (non-nucleoside reverse transcriptase inhibitors), CCR5 antagonists and FIs (fusion inhibitors)]. Darunavir with ritonavir has not been tested with all PIs (protease inhibitors) and must not be used with other HIV PIs. In some cases dosage of other medicines might need to be changed. Therefore always tell your doctor if you take other anti-HIV medicines and follow your doctor's instruction carefully on which medicines can be combined.

The effects of Darunavir Rowex might be reduced if you take any of the following products. Tell your doctor if you take:

- *Phenobarbital, phenytoin* (to prevent seizures)
- *Dexamethasone* (corticosteroid)
- *Efavirenz* (HIV infection)
- *Boceprevir* (hepatitis C infection)
- *Rifapentine, rifabutin* (medicines to treat some infections such as tuberculosis)

- *Saquinavir* (HIV infection).

The effects of other medicines might be influenced if you take Darunavir Rowex.

Tell your doctor if you take:

- *Amlodipine, diltiazem, disopyramide, carvedilol, felodipine, flecainide, lidocaine, metoprolol, mexiletine, nifedipine, nicardipine, propafenone, timolol, verapamil* (for heart disease) as the therapeutic effect or side effects of these medicines may be increased.
- *Apixaban, edoxaban, rivaroxaban, warfarin* (to reduce clotting of the blood) as their therapeutic effect or side effects may be altered; your doctor may have to check your blood.
- Oestrogen-based hormonal contraceptives and hormonal replacement therapy. Darunavir Rowex might reduce its effectiveness. When used for birth control, alternative methods of non-hormonal contraception are recommended.
- Ethinylestradiol/drospirenone. Darunavir Rowex might increase the risk for elevated potassium levels by drospirenone.
- *Atorvastatin, pravastatin, rosuvastatin* (to lower cholesterol levels). The risk of muscle damage might be increased. Your doctor will evaluate which cholesterol lowering regimen is best for your specific situation.
- *Clarithromycin* (antibiotic)
- *Ciclosporin, everolimus, tacrolimus, sirolimus* (for dampening down immune system) as the therapeutic effect or side effects of these medicines might be increased. Your doctor might want to do some additional tests.
- Corticosteroids including betamethasone, budesonide, fluticasone, mometasone, prednisone, triamcinolone. These medicines are used to treat allergies, asthma, inflammatory bowel diseases, inflammatory conditions of the eyes, joints and muscles and other inflammatory conditions. If alternatives cannot be used, its use should only take place after medical evaluation and under close monitoring by your doctor for corticosteroid side effects.
- *Buprenorphine/naloxone* (medicines to treat opiod dependence)
- *Salmeterol* (medicine to treat asthma)
- *Artemether/lumefantrine* (a combination medicine to treat malaria).
- *Dasatinib, everolimus, nilotinib, vinblastine, vincristine* (to treat cancer)
- *Sildenafil, tadalafil, vardenafil* (for erectile dysfunction or to treat a heart and lung disorder called pulmonary arterial hypertension).
- Glecaprevir/pibrentasvir, simeprevir (to treat hepatitis C infection)
- Fentanyl, oxycodone, tramadol (to treat pain).
- 

The dosage of other medicines might need to be changed since either their own or Darunavir Rowex's therapeutic effect or side effects may be influenced when combined.

Tell your doctor if you take:

- *Alfentanil* (injectable strong and short-acting painkiller that is used for surgical procedures)
- *Digoxin* (to treat certain heart disorders)
- *Clarithromycin* (antibiotic)
- *Itraconazole, isavuconazole, fluconazole, posaconazole, clotrimazole* (to treat fungal infections). Voriconazole should only be taken after medical evaluation.
- *Rifabutin* (against bacterial infections)
- *Sildenafil, vardenafil, tadalafil* (for erectile dysfunction or high blood pressure in the pulmonary circulation)
- *Amitriptyline, desipramine, imipramine, nortriptyline, paroxetine, sertraline, trazodone* (to treat depression and anxiety)
- *Maraviroc* (to treat HIV infection)
- *Methadone* (to treat opiate dependence)
- *Carbamazepine, clonazepam* (to prevent seizures or to treat certain types of nerve pain)

- *Colchicine* (to treat gout or familial Mediterranean fever)
- *Bosentan* (to treat high blood pressure in the pulmonary circulation)
- *Bupirone, clorazepate, diazepam, estazolam, flurazepam, midazolam when used as injection, zoldipem* (sedative agents)
- *Perphenazine, risperidone, thioridazine* (to treat psychiatric conditions).

This is **not** a complete list of medicines. Tell your healthcare provider about **all** medicines that you are taking.

### **Darunavir Rowex with food and drink**

See section 3 'How to take Darunavir Rowex.'

### **Pregnancy and breast-feeding**

Tell your doctor immediately if you are pregnant, planning to become pregnant or if you are breast-feeding. Pregnant or breast-feeding mothers should not take Darunavir Rowex with ritonavir unless specifically directed by the doctor. Pregnant or breast-feeding mothers should not take Darunavir Rowex with cobicistat.

It is recommended that HIV infected women must not breast-feed their infants because of both the possibility of your baby becoming infected with HIV through your breast milk and because of the unknown effects of the medicine on your baby.

### **Driving and using machines**

Do not operate machines or drive if you feel dizzy after taking Darunavir Rowex.

**Darunavir Rowex 300 mg film-coated tablets contain sunset yellow FCF (E110)** which may cause allergic reactions.

**Darunavir Rowex 600 mg film-coated tablets contain sunset yellow FCF (E110)** which may cause allergic reactions.

## **3. How to take Darunavir Rowex**

Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. Check with your doctor, pharmacist or nurse if you are not sure. Even if you feel better, do not stop taking Darunavir Rowex and ritonavir without talking to your doctor.

After therapy has been initiated, the dose or dosage form must not be changed or therapy must not be stopped without instruction of the doctor.

### **Dose for children of 3 years of age and above, weighing at least 15 kilograms who have not taken antiretroviral medicines before (your child's doctor will determine this)**

The doctor will work out the right once daily dose based on the weight of the child (see table below). This dose must not exceed the recommended adult dose, which is 800 milligram Darunavir Rowex together with 100 milligram ritonavir once a day.

The doctor will inform you on how much Darunavir Rowex tablets and how much ritonavir (capsules, tablets or solution) the child must take.

<b>Weight</b>	<b>One darunavir dose is</b>	<b>One ritonavir<sup>a</sup> dose is</b>
between 15 and 30 kilograms	600 milligram	100 milligram
between 30 and 40 kilograms	675 milligram	100 milligram
more than 40 kilograms	800 milligram	100 milligram

<sup>a</sup> ritonavir oral solution: 80 milligram per milliliter

**Dose for children of 3 years of age and above, weighing at least 15 kilograms who have taken antiretroviral medicines before (your child's doctor will determine this)**

The doctor will work out the right dose based on the weight of the child (see table below). The doctor will determine if once daily dosing or twice daily dosing is appropriate for the child. This dose must not exceed the recommended adult dose, which is 600 milligram Darunavir Rowex together with 100 milligram ritonavir two times per day or 800 milligram Darunavir Rowex together with 100 milligram ritonavir once a day. The doctor will inform you on how many Darunavir Rowex tablets and how much ritonavir (capsules, tablets or solution) the child must take. Tablets of other strengths are available and your doctor may have prescribed a certain combination of tablets to construct the appropriate dosing regimen.

**Twice daily dosing**

<b>Weight</b>	<b>One dose is</b>
between 15 and 30 kilograms	375 milligram darunavir + 50 milligram ritonavir twice a day
between 30 and 40 kilograms	450 milligram darunavir + 60 milligram ritonavir twice a day
more than 40 kilograms*	600 milligram darunavir + 100 milligram ritonavir twice a day

\* For children aged 12 or more and weighing at least 40 kilograms, your child's doctor will determine if Darunavir Rowex 800 milligram once daily dosing may be used. This cannot be administered with these tablets. Other strengths of Darunavir Rowex are available.

**Once daily dosing**

<b>Weight</b>	<b>One darunavir dose is</b>	<b>One ritonavir<sup>a</sup> dose is</b>
between 15 and 30 kilograms	600 milligram	100 milligram
between 30 and 40 kilograms	675 milligram	100 milligram
more than 40 kilograms	800 milligram	100 milligram

<sup>a</sup> ritonavir oral solution: 80 milligram per milliliter

**Instructions for children**

- The child must take Darunavir Rowex always together with ritonavir. Darunavir Rowex cannot work properly without ritonavir.
- The child must take the appropriate doses of Darunavir Rowex and ritonavir two times per day or once a day. If prescribed Darunavir Rowex twice daily the child must take one dose in the morning, and one dose in the evening. Your child's doctor will determine the appropriate dosing regimen for your child.
- The child must take Darunavir Rowex with food. Darunavir Rowex cannot work properly without food. The type of food is not important.
- The child must swallow the tablets with a drink such as water or milk.
- Darunavir Rowex 75 milligram and 150 milligram tablets have been developed for use in children weighing less than 40 kilograms, but can also be used in some cases.

**Dose for adults who have not taken antiretroviral medicines before (your doctor will determine this)**

You will require a different dose of Darunavir Rowex which cannot be administered with these tablets. Other strengths of Darunavir Rowex are available.

**Dose for adults who have taken antiretroviral medicines before (your doctor will**

**determine this)**

The dose is either:

- 600 milligram Darunavir Rowex (2 tablets containing 300 milligram of Darunavir Rowex or 1 tablet containing 600 milligram of Darunavir Rowex) together with 100 milligram ritonavir twice daily.
- OR
- 800 milligram Darunavir Rowex (2 tablets containing 400 milligram of Darunavir Rowex or 1 tablet containing 800 milligram of Darunavir Rowex) together with 100 milligram ritonavir once daily. Darunavir Rowex 400 milligram and 800 milligram tablets are only to be used to construct the once daily 800 milligram regimen.

Please discuss with your doctor which dose is right for you.

**Instructions for adults**

- Take Darunavir Rowex always together with ritonavir. Darunavir Rowex cannot work properly without ritonavir.
- In the morning, take 600 milligram Darunavir Rowex together with 100 milligram ritonavir.
- In the evening, take 600 milligram Darunavir Rowex tablet together with 100 milligram ritonavir.
- Take Darunavir Rowex with food. Darunavir Rowex cannot work properly without food. The type of food is not important.
- Swallow the tablets with a drink such as water or milk.
- Darunavir Rowex 75 milligram and 150 milligram tablets have been developed for use in children, but can also be used in adults in some cases.

**Removing the child resistant cap**

The plastic bottle comes with a child resistant cap and must be opened as follows:

Push the plastic screw cap down while turning it counter clockwise.

Remove the unscrewed cap.

**If you take more Darunavir Rowex than you should**

Contact your doctor, pharmacist or nurse immediately.

**If you forget to take Darunavir Rowex**

If you notice **within 6 hours**, you must take the tablets immediately. Always take with ritonavir and food. If you notice **after 6 hours**, then skip the intake and take the next doses as usual. Do not take a double dose to make up for a forgotten dose.

**Do not stop taking Darunavir Rowex without talking to your doctor first**

Anti-HIV medicines may make you feel better. Even when you feel better, do not stop taking Darunavir Rowex. Talk to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

**4. Possible side effects**

During HIV therapy there may be an increase in weight and in levels of blood lipids and

glucose. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Tell your doctor if you develop any of the following side effects.**

Liver problems that may occasionally be severe have been reported. Your doctor should do blood tests before you start Darunavir Rowex. If you have chronic hepatitis B or C infection, your doctor should check your blood tests more often because you have an increased chance of developing liver problems. Talk to your doctor about the signs and symptoms of liver problems. These may include yellowing of your skin or whites of your eyes, dark (tea coloured) urine, pale coloured stools (bowel movements), nausea, vomiting, loss of appetite, or pain, aching, or pain and discomfort on your right side below your ribs.

Skin rash (more often when used in combination with raltegravir), itching. The rash is usually mild to moderate. A skin rash might also be a symptom of a rare severe situation. It is therefore important to talk to your doctor if you develop a rash. Your doctor will advise you how to deal with your symptoms or whether Darunavir Rowex must be stopped.

Other severe side effects were diabetes (common) and inflammation of the pancreas (uncommon).

**Very common side effects** (may affect more than 1 in 10 people)

- diarrhoea.

**Common side effects** (may affect up to 1 in 10 people)

- vomiting, nausea, abdominal pain or distension, dyspepsia, flatulence
- headache, tiredness, dizziness, drowsiness, numbness, tingling or pain in hands or feet, loss of strength, difficulty falling asleep.

**Uncommon side effects** (may affect up to 1 in 100 people)

- chest pain, changes in electrocardiogram, rapid heart beating
- decreased or abnormal skin sensibility, pins and needles, attention disturbance, loss of memory, problems with your balance
- difficulty breathing, cough, nosebleed, throat irritation
- inflammation of the stomach or mouth, heartburn, retching, dry mouth, discomfort of the abdomen, constipation, belching
- kidney failure, kidney stones, difficult discharge of urine, frequent or excessive passage of urine, sometimes at night
- urticaria, severe swelling of the skin and other tissues (most often the lips or the eyes), eczema, excessive sweating, night sweats, hair loss, acne, scaly skin, colouration of nails
- muscle pain, muscle cramps or weakness, pain in extremity, osteoporosis
- slowing down of the thyroid gland function. This can be seen in a blood test.
- high blood pressure, flushing
- red or dry eyes
- fever, swelling of lower limbs due to fluids, malaise, irritability, pain
- symptoms of infection, herpes simplex
- erectile dysfunction, enlargement of breasts
- sleeping problems, sleepiness, depression, anxiety, abnormal dreams, decrease in sexual drive

**Rare side effects** (may affect up to 1 in 1,000 people)

- a reaction called DRESS [severe rash, which may be accompanied by fever, fatigue, swelling of the face or lymph glands, increase of eosinophils (type of white blood cells), effects on liver, kidney or lung]

- heart attack, slow heart beating, palpitations
- visual disturbance
- chills, feeling abnormal
- a feeling of confusion or disorientation, altered mood, restlessness
- fainting, epileptic fits, changes or loss of taste
- mouth sores, vomiting blood, inflammation of the lips, dry lips, coated tongue
- running nose
- skin lesions, dry skin
- stiffness of muscles or joints, joint pain with or without inflammation
- changes in some values of your blood cells or chemistry. These can be seen in the results of blood and/or urine tests. Your doctor will explain these to you. Examples are: increase in some white blood cells.

Some side effects are typical for anti-HIV medicines in the same family as Darunavir Rowex. These are:

- muscle pain, tenderness or weakness. On rare occasions, these muscle disorders have been serious (rhabdomyolysis).

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRAs Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: [www.hpra.ie](http://www.hpra.ie); E-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Darunavir Rowex**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, the blister and on the bottle after EXP. The expiry date refers to the last day of that month.

After first opening of the bottle: 6 months, when not stored above 25°C.

For blister: Do not store above 30°C

For bottle: This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away any medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Darunavir Rowex contains**

#### **Darunavir Rowex 75 mg Film-coated tablets**

The active substance is darunavir.

Each film-coated tablet contains 75 mg of darunavir.

The other ingredients are:

- Tablet core: cellulose, microcrystalline (E460), silica, colloidal anhydrous (E551), crospovidone (type A) (E1202), magnesium stearate (E470b).
- Tablet coating: poly (vinyl alcohol) (E1203), macrogol (3350) (E1521), titanium dioxide (E171), talc (E553b).

#### **Darunavir Rowex 600 mg Film-coated tablets**

The active substance is darunavir.

Each film-coated tablet contains 600 mg of darunavir.

The other ingredients are:

- Tablet core: cellulose, microcrystalline (E460), silica, colloidal anhydrous (E551), crospovidone (type A) (E1202), magnesium stearate (E470b).
- Tablet coating: poly (vinyl alcohol) (E1203), macrogol (3350) (E1521), titanium dioxide (E171), talc (E553b), sunset yellow FCF (E110).

### **What Darunavir Rowex looks like and contents of the pack**

#### **Darunavir Rowex 75 mg film-coated tablets**

White, caplet shaped film-coated tablet, debossed with '75' on one side and plain on the other side.

Dimensions: approximately 8.6 mm x 4.6 mm

#### **Darunavir Rowex 600 mg film-coated tablets**

Orange oval shaped film-coated tablet, debossed with '600' on one side and plain on the other side.

Dimensions: approximately 20.1 mm x 10.1 mm

Darunavir Rowex is packed in Aluminium-PVC/PE/PVDC perforated blisters or in HDPE bottles stoppered with polypropylene (PP), child resistant closure.

#### Pack size:

Bottle packs 75 mg: 480 film-coated tablets

Bottle packs 600 mg: 60, 120 (2x60), 180 (3x60), 240 (4x60) film-coated tablets

Perforated blister packs 75 mg: 10, 30, 60, 90, 120 film-coated tablets

Perforated blister packs 600 mg: 10, 30, 60, 90, 120 film-coated tablets

Perforated unit dose blister packs 600 mg: 60x1 film-coated tablets

### **Marketing Authorisation Holder and Manufacturers**

#### **Marketing Authorisation Holder**

Rowex Ltd., Bantry, Co. Cork, Ireland.

#### **Manufacturers**

Lek Pharmaceuticals d.d., Verovškova ulica 57, Ljubljana, 1526, Slovenia.

Salutas Pharma GmbH, Otto-von-Guericke-Allee 1, Sachsen-Anhalt, 39179 Barleben, Germany.

Remedica Ltd, Aharon Street, Limassol Industrial Estate, Limassol 3056, Cyprus.

### **This medical product is authorized in the Member State of the EEA under the following names:**

The Netherlands:	Darunavir Sandoz 75 mg, filmomhulde tabletten Darunavir Sandoz 600 mg, filmomhulde tabletten
Austria:	Darunavir Sandoz 75 mg – Filmtabletten Darunavir Sandoz 600 mg – Filmtabletten
Belgium:	Darunavir Sandoz 75 mg filmomhulde tabletten Darunavir Sandoz 600 mg filmomhulde tabletten
Croatia:	Darunavir Sandoz 75 mg filmom obložene tablete Darunavir Sandoz 600 mg filmom obložene tablete
Czech Republic:	Darunavir Sandoz
Denmark:	Darunavir Sandoz 75 mg filmovertrukne tabletter Darunavir Sandoz 600 mg filmovertrukne tabletter
Estonia:	Darunavir Sandoz
Finland:	Darunavir Sandoz 75 mg tabletti, kalvopäällysteinen Darunavir Sandoz 600 mg tabletti, kalvopäällysteinen

France:	Darunavir Sandoz 75 mg, comprimé pelliculé Darunavir Sandoz 600 mg, comprimé pelliculé
Germany:	Darunavir – 1 A Pharma 75 mg Filmtabletten Darunavir – 1 A Pharma 600 mg Filmtabletten
Hungary:	Darunavir Sandoz 75 mg filmtabletta Darunavir Sandoz 600 mg filmtabletta
Ireland:	Darunavir Rowex 75 mg Film-coated tablets Darunavir Rowex 600 mg Film-coated tablets
Latvia:	Darunavir Sandoz 75 mg apvalkotās tabletes Darunavir Sandoz 600 mg apvalkotās tabletes
Lithuania:	Darunavir Sandoz 75 mg plėvele dengtos tabletės Darunavir Sandoz 600 mg plėvele dengtos tabletės
Poland:	Darunavir Sandoz
Portugal:	Darunavir Sandoz
Romania:	Darunavir Sandoz 75 mg comprimate filmate Darunavir Sandoz 600 mg comprimate filmate
Slovenia:	Darunavir Sandoz 75 mg filmsko obložene tablete Darunavir Sandoz 600 mg filmsko obložene tablete
Spain:	Darunavir Sandoz 75 mg comprimidos recubiertos con película EFG Darunavir Sandoz 600 mg comprimidos recubiertos con película EFG
Sweden:	Darunavir Sandoz 75 mg filmdragerade tabletter Darunavir Sandoz 600mg filmdragerade tabletter
United Kingdom:	Darunavir Sandoz 75mg film-coated tablets Darunavir Sandoz 600mg film-coated tablets

**This leaflet was last revised in 12/2018.**