

## **Package leaflet: Information for the patient**

### **Pantoprazole Teva Pharma 40 mg Gastro-resistant Tablets**

pantoprazole

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Pantoprazole Teva Pharma is and what it is used for
2. What you need to know before you take Pantoprazole Teva Pharma
3. How to take Pantoprazole Teva Pharma
4. Possible side effects
5. How to store Pantoprazole Teva Pharma
6. Contents of the pack and other information

#### **1. What Pantoprazole Teva Pharma is and what it is used for**

Pantoprazole Teva Pharma contains the active substance pantoprazole. Pantoprazole Teva Pharma is a selective “proton pump inhibitor”, a medicine which reduces the amount of acid produced in your stomach. It is used for treating acid-related diseases of the stomach and intestine.

Pantoprazole Teva Pharma is used to treat adults and adolescents 12 years of age and above for:

- Reflux oesophagitis. An inflammation of your oesophagus (the tube which connects your throat to your stomach) accompanied by the regurgitation of stomach acid.

Pantoprazole Teva Pharma is used to treat adults for:

- An infection with a bacterium called *Helicobacter pylori* in patients with duodenal ulcers and stomach ulcers in combination with two antibiotics (Eradication therapy). The aim is to get rid of the bacteria and so reduce the likelihood of these ulcers returning.
- Stomach and duodenal ulcers.
- Zollinger-Ellison-Syndrome and other conditions producing too much acid in the stomach.

#### **2. What you need to know before you take Pantoprazole Teva Pharma**

##### **Do not take Pantoprazole Teva Pharma**

- If you are allergic to pantoprazole or to any of the other ingredients of this medicine (listed in section 6).
- If you are allergic to medicines containing other proton pump inhibitors.

##### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Pantoprazole Teva Pharma

- If you have severe liver problems. Please tell your doctor if you have ever had problems with your liver in the past. Your doctor will check your liver enzymes more frequently, especially when you

are taking Pantoprazole Teva Pharma as a long-term treatment. In the case of a rise of liver enzymes the treatment should be stopped.

- If you have reduced body stores or risk factors for reduced vitamin B12 and receive long-term treatment with pantoprazole. As with all acid reducing agents, pantoprazole may lead to a reduced absorption of vitamin B12. Please contact your doctor if you notice any of the following symptoms, which could indicate low levels of Vitamin B12:
  - Extreme tiredness or lack of energy
  - Pins and needles
  - Sore or red tongue, mouth ulcers
  - Muscle weakness
  - Disturbed vision
  - Problems with memory, confusion, depression
- If you are taking HIV protease inhibitors such as atazanavir (for the treatment of HIV-infection) at the same time as pantoprazole, ask your doctor for specific advice.
- Taking a proton pump inhibitor like pantoprazole, especially over a period of more than one year, may slightly increase your risk of fracture in the hip, wrist or spine. Tell your doctor if you have osteoporosis (reduced bone density) or if you have been told that you are at risk of getting osteoporosis (for example, if you are taking steroids).
- If you are on pantoprazole for more than three months it is possible that the levels of magnesium in your blood may fall. Low levels of magnesium can be seen as fatigue, involuntary muscle contractions, disorientation, convulsions, dizziness or increased heart rate. If you get any of these symptoms, please tell your doctor promptly. Low levels of magnesium can also lead to a reduction in potassium or calcium levels in the blood. Your doctor may decide to perform regular blood tests to monitor your levels of magnesium.
- If you have ever had a skin reaction after treatment with a medicine similar to Pantoprazole Teva Pharma that reduces stomach acid.
- If you get a rash on your skin, especially in areas exposed to the sun tell your doctor as soon as you can, as you may need to stop your treatment with Pantoprazole Teva Pharma. Remember to also mention any other ill-effects like pain in your joints.
- Serious skin reactions including Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms (DRESS) and erythema multiforme, have been reported in association with pantoprazole treatment. Stop using pantoprazole and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.
- If you are due to have a specific blood test (Chromogranin A).

**Tell your doctor immediately** before or after taking this medicine, if you notice any of the following symptoms, which could be a sign of another, more serious, disease:

- an unintentional loss of weight
- vomiting, particularly if repeated
- vomiting blood; this may appear as dark coffee grounds in your vomit
- you notice blood in your stools; which may be black or tarry in appearance
- difficulty in swallowing or pain when swallowing
- you look pale and feel weak (anaemia)
- chest pain
- stomach pain
- severe and/or persistent diarrhoea, because this medicine has been associated with a small increase in infectious diarrhoea

Your doctor may decide that you need some tests to rule out malignant disease because pantoprazole also alleviates the symptoms of cancer and could cause delay in diagnosing it. If your symptoms continue in spite of your treatment, further investigations will be considered.

If you take Pantoprazole Teva Pharma on a long-term basis (longer than 1 year) your doctor will probably keep you under regular surveillance. You should report any new and exceptional symptoms and circumstances whenever you see your doctor.

### **Children and adolescents**

Pantoprazole Teva Pharma is not recommended for use in children as it has not been proven to work in children below 12 years of age.

### **Other medicines and Pantoprazole Teva Pharma**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

This is because Pantoprazole Teva Pharma may influence the effectiveness of other medicines, so tell your doctor if you are taking:

- Medicines such as ketoconazole, itraconazole and posaconazole (used to treat fungal infections) or erlotinib (used for certain types of cancer) because Pantoprazole Teva Pharma may stop these and other medicines from working properly.
- Warfarin and phenprocoumon, which affect the thickening, or thinning of the blood. You may need further checks.
- Medicines used to treat HIV-infection, such as atazanavir.
- Methotrexate (used to treat rheumatoid arthritis, psoriasis and cancer) – if you are taking methotrexate your doctor may temporarily stop your treatment with Pantoprazole Teva Pharma because pantoprazole can increase levels of methotrexate in the blood.
- Fluvoxamine (used to treat depression and other psychiatric diseases) – if you are taking fluvoxamine your doctor may reduce the dose.
- Rifampicin (used to treat infections).
- St John's wort (*Hypericum perforatum*) (used to treat mild depression).

Talk to your doctor before taking Pantoprazole Teva Pharma if you are due to have a specific urine test (for THC; Tetrahydrocannabinol).

### **Pregnancy and breast-feeding**

There are no adequate data from the use of pantoprazole in pregnant women. Excretion into human milk has been reported.

If you are pregnant or breast feeding, think you may be pregnant, or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

You should use this medicine, only if your doctor considers the benefit for you greater than the potential risk for your unborn child or baby.

### **Driving and using machines**

Pantoprazole Teva Pharma has no or negligible influence on the ability to drive and use machines. If you experience side effects like dizziness or disturbed vision, you should not drive or operate machines.

### **Pantoprazole Teva Pharma contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per gastro-resistant tablet, that is to say essentially 'sodium-free'.

## **3. How to take Pantoprazole Teva Pharma**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

### **Method of administration**

Take the tablets 1 hour before a meal without chewing or breaking them and swallow them whole with some water.

The recommended dose is:

**Adults and adolescents 12 years of age and above**

- *To treat reflux oesophagitis*

The recommended dose is one tablet a day. Your doctor may tell you to increase to 2 tablets daily. The treatment period for reflux oesophagitis is usually between 4 and 8 weeks. Your doctor will tell you how long to take your medicine.

**Adults**

- *For the treatment of an infection with a bacterium called Helicobacter pylori in patients with duodenal ulcers and stomach ulcers in combination with two antibiotics (Eradication therapy).*

One tablet, two times a day plus two antibiotic tablets of either amoxicillin, clarithromycin and metronidazole (or tinidazole), each to be taken two times a day with your pantoprazole tablet. Take the first pantoprazole tablet 1 hour before breakfast and the second pantoprazole tablet 1 hour before your evening meal. Follow your doctor's instructions and make sure you read the package leaflets for these antibiotics. The usual treatment period is one to two weeks.

- *For the treatment of stomach and duodenal ulcers.*

The recommended dose is one tablet a day. After consultation with your doctor, the dose may be doubled. Your doctor will tell you how long to take your medicine. The treatment period for stomach ulcers is usually between 4 and 8 weeks. The treatment period for duodenal ulcers is usually between 2 and 4 weeks.

- *For the long-term treatment of Zollinger-Ellison-Syndrome and of other conditions in which too much stomach acid is produced.*

The recommended starting dose is usually two tablets a day.

Take the two tablets 1 hour before a meal. Your doctor may later adjust the dose, depending on the amount of stomach acid you produce. If prescribed more than two tablets a day, the tablets should be taken twice daily.

If your doctor prescribes a daily dose of more than four tablets a day, you will be told exactly when to stop taking the medicine.

**Patients with kidney problems**

If you have kidney problems, you should not take pantoprazole for eradication of Helicobacter pylori.

**Patients with liver problems**

If you suffer from severe liver problems, you should not take more than one tablet 20 mg pantoprazole a day (for this purpose tablets containing 20 mg pantoprazole are available).

If you suffer from moderate or severe liver problems, you should not take pantoprazole for eradication of Helicobacter pylori.

**Use in children and adolescents**

These tablets are not recommended for use in children below 12 years.

**If you take more Pantoprazole Teva Pharma than you should**

Consult your doctor or pharmacist. There are no known symptoms of overdose.

**If you forget to take Pantoprazole Teva Pharma**

Do not take a double dose to make up for a forgotten dose. Take your next, normal dose at the usual time.

**If you stop taking Pantoprazole Teva Pharma**

Do not stop taking these tablets without first talking to your doctor or pharmacist.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**If you get any of the following side effects, stop taking these tablets and tell your doctor immediately, or contact the casualty department at your nearest hospital:**

- **Serious allergic reactions (frequency rare:** may affect up to 1 in 1,000 people): swelling of the tongue and/or throat, difficulty in swallowing, hives (nettle rash), difficulties in breathing, allergic facial swelling (Quincke's oedema/ angioedema), severe dizziness with very fast heartbeat and heavy sweating.

**Serious skin conditions (frequency not known:** frequency cannot be estimated from the available data): you may notice one or more of the following - blistering of the skin and rapid deterioration of your general condition, erosion (including slight bleeding) of eyes, nose, mouth/lips or genitals, or skin sensitivity/rash, particularly in areas of skin exposed to light/the sun. You may also have joint pain or flu-like symptoms, a fever, swollen glands (e.g. in the armpit) and blood tests may show changes in certain white blood cells or liver enzymes - reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (Stevens-Johnson syndrome, toxic epidermal necrolysis) - widespread rash, high body temperature and enlarged lymph nodes (DRESS syndrome or drug hypersensitivity syndrome).

- **Other serious conditions (frequency not known:** frequency cannot be estimated from the available data): yellowing of the skin or whites of the eyes (severe damage to liver cells, jaundice) or fever, rash, and enlarged kidneys sometimes with painful urination, and lower back pain (serious inflammation of the kidneys), possibly leading to kidney failure.

Other side effects are:

- **Common** (may affect up to 1 in 10 people)  
Benign polyps in the stomach.
- **Uncommon** (may affect up to 1 in 100 people) Headache; dizziness; diarrhoea; feeling sick, vomiting; bloating and flatulence (wind); constipation; dry mouth; abdominal pain and discomfort; skin rash, exanthema, eruption; itching; feeling weak, exhausted or generally unwell; sleep disorders; fracture in the hip, wrist or spine.
- **Rare** (may affect up to 1 in 1,000 people)  
Distortion or complete lack of the sense of taste; disturbances in vision such as blurred vision; hives; pain in the joints; muscle pains; weight changes; raised body temperature; high fever; swelling of the extremities (peripheral oedema); allergic reactions; depression; breast enlargement in males.
- **Very rare** (may affect up to 1 in 10,000 people)  
Disorientation.
- **Not known** (frequency cannot be estimated from the available data)  
Hallucination, confusion (especially in patients with a history of these symptoms); muscle spasms due to electrolyte disturbances (changes in the salt levels in the body); feeling of tingling, prickling, pins and needles, burning sensation or numbness; inflammation in the large bowel, that causes persistent watery diarrhoea.

**Side effects identified through blood tests:**

- **Uncommon** (may affect up to 1 in 100 people)  
An increase in liver enzymes.
- **Rare** (may affect up to 1 in 1,000 people)

- An increase in bilirubin; increased fat levels in blood; sharp drop in circulating granular white blood cells, associated with high fever.
- **Very rare** (may affect up to 1 in 10,000 people)  
A reduction in the number of blood platelets, which may cause you to bleed or bruise more than normal; a reduction in the number of white blood cells, which may lead to more frequent infections; coexisting abnormal reduction in the number of red and white blood cells, as well as platelets, which may lead to tiredness, being short of breath and looking pale.
  - **Not known** (frequency cannot be estimated from the available data)  
Decreased level of sodium, magnesium, calcium or potassium in blood (see section 2)

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance Website: [www.hpra.ie](http://www.hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Pantoprazole Teva Pharma**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date, which is stated on the blister, HDPE tablet container or outer carton after EXP. The expiry date refers to the last day of that month.

Store below 30 °C.

### HDPE tablet container:

After first opening the container, use Pantoprazole Teva Pharma within 60 days.

Do not throw away any medicines via wastewater <or household waste>. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Pantoprazole Teva Pharma contains**

The active substance is pantoprazole.

Each gastro-resistant tablet contains 40 mg pantoprazole (as sodium sesquihydrate).

The other ingredients are:

#### **Tablet core:**

Disodium Phosphate,  
Mannitol,  
Cellulose Microcrystalline,  
Croscarmellose Sodium,  
Magnesium Stearate,  
Hypromellose,  
Triethyl Citrate  
Sodium Starch Glycolate;

#### **Film-coating:**

Methacrylic acid-Ethyl acrylate copolymer,  
Yellow Iron Oxide,

Triethyl Citrate

### **What Pantoprazole Teva Pharma looks like and contents of the pack**

Pantoprazole Teva Pharma Gastro-resistant Tablets are oval, yellow tablets and are available in either aluminium blister packs of 7, 14, 15, 28, 30, 50x1, 56, 90, 98,100 & 120 tablets or HDPE containers with child resistant closure of 28, 50, 56 and 100 (50x2) tablets.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

Teva Pharma B.V.  
Swensweg 5,  
2031GA Haarlem,  
The Netherlands

### **Manufacturer**

Teva Pharma, S.L.U  
Poligono Industrial Malpica,  
calle C, Numero 4, 50016  
Zaragoza, Spain

TEVA Pharmaceutical Works Private Limited Company  
Pallagi út 13, 4042 Debrecen  
Hungary

Pharmachemie B.V.  
Swensweg 5, 2031 GA Haarlem  
Netherlands

**This medicine is authorised in the Member States of the European Economic Area under the following names:**

Ireland	Pantoprazole Teva Pharma 40 mg Gastro-resistant Tablets
Belgium	Pantoprazole Teva 40 mg maagsapresistente tabletten
Bulgaria	PRAZOLPAN 40 mg gastro-resistant tablets
Denmark	Pantoprazole Teva
Greece	Pantoprazole Teva Pharma 40 mg Γαστροανθεκτικά Δισκία
Finland	Pantoprazole ratiopharm 40 mg enterotablettia
Hungary	Pantoprazole-Teva 40 mg gyomornedv-ellenálló tableta
Italy	Pantoprazolo Teva Italia
Netherlands	Pantoprazol Teva 40 mg, maagsapresistente tabletten
Romania	Pantoprazole Teva 40 mg comprimate gastrorezistente
Sweden	Ulzate 40 mg enterotabletter
Slovenia	Pantoprazol Teva 40 mg gastrorezistentne tablete

**This leaflet was last revised in 02/2024.**