

PATIENT ALERT CARD

Dabigatran Etxilate Krka
hard capsules
(dabigatran etexilate)



XXXXXX

- ◆ This card should be with you / the caregiver at all times
- ◆ Make sure to use the latest version

Dear Patient / Caregiver of a paediatric patient,

Your / your child's doctor has initiated treatment with Dabigatran Etxilate Krka.

In order to use Dabigatran Etxilate Krka safely, please consider the important information in the package leaflet.

As this patient alert card contains

important information about your / your child's treatment, this card should be with you / your child at all times to inform healthcare professionals about your / your child's intake of Dabigatran Etxilate Krka.



Dabigatran Etxilate Krka Information for Patients / Caregivers of paediatric patients

About your / your child's treatment

- ◆ Dabigatran Etxilate Krka thins the blood. It is used to treat existing blood clots or to prevent the formation of dangerous blood clots.
- ◆ Follow your / your child's doctor's instructions when taking Dabigatran Etxilate Krka. Never skip a dose or stop the intake of Dabigatran Etxilate Krka without talking to your / your child's doctor.
- ◆ Inform your / your child's doctor about all medicines you / your child are / is currently taking.

- ◆ Inform your / your child's doctor about the intake of Dabigatran Etxilate Krka before any surgery / invasive procedure.
- ◆ Dabigatran Etxilate Krka capsules can be taken with or without food. The capsule should be swallowed whole with a glass of water. The capsule must not be broken or chewed and the pellets must not be emptied from the capsule.

When to seek medical advice

- ◆ Taking Dabigatran Etxilate Krka may increase the risk of bleeding. Speak to your / your child's doctor immediately if you / your child experience(s) signs and symptoms of bleeding such as: swelling, discomfort, unusual pain or headache,

dizziness, paleness, weakness, unusual bruising, nosebleeds, bleeding of gums, unusual long bleeding cuts, abnormal menstrual flow or vaginal bleeding, blood in the urine which may be pink or brown, red/black stools, coughing up blood, vomiting blood or coffee ground like material.

- ◆ In case of fall or injury, especially if the head is hit, urgently seek medical advice.
- ◆ Do not stop intake of Dabigatran Etxilate Krka without talking to your / your child's doctor, if you / your child experience(s) heartburn, nausea, vomiting, stomach discomfort, bloating or upper abdominal pain.

Dabigatran Etxilate Krka Information for Healthcare Professionals

- ◆ Dabigatran Etxilate Krka is an oral anticoagulant (direct thrombin inhibitor).
- ◆ Dabigatran Etxilate Krka may need to be stopped in advance of surgical or other invasive procedures.
- ◆ In case of major bleeding events, Dabigatran Etxilate Krka must be stopped immediately.
- ◆ A specific reversal agent (idarucizumab) is available for adult patients. The efficacy and safety of the specific reversal agent idarucizumab have not been established in paediatric patients.

For details and more advice to antagonise the anticoagulant effect of Dabigatran Etexilate Krka please refer to the Summary of Product Characteristics of Dabigatran Etexilate Krka and idarucizumab.

◆ Dabigatran Etexilate Krka is mainly eliminated by the kidneys; adequate diuresis must be maintained. Dabigatran Etexilate Krka is dialyzable.

Please complete this section or ask your / your child's doctor to do it.
Patient Information

Name of the patient

Date of birth

Indication for anticoagulation

Dose of Dabigatran Etexilate Krka