UPTRAVI FILM-COATED TABLETS selexipag

Patient Titration Guide: Starting Treatment With Uptravi

Please read the accompanying patient information leaflet before starting treatment.

Tell your doctor if you experience side effects, take any other medicines or have any other health issues as they may need to make an adjustment to your Uptravi dose.



CONTENTS

•	How should you take UPTRAVI?	. 5
•	Talk to your doctor	.6
•	How should you step up your dose?	.8
•	What are the steps? 1	10
•	When should you step down?	12
•	Stepping down	14

•	When you move to your maintenance dose	.16
•	If you forget to take UPTRAVI	. 18
•	If you stop taking UPTRAVI	. 19
•	Titration diary	20

UPTRAVI is a medicine taken every morning and evening for the treatment of pulmonary arterial hypertension, also called PAH.

The starting dose for UPTRAVI is 200 micrograms **once in the morning and once in the evening.** The first intake of UPTRAVI should be in the evening. You should take each dose with a glass of water, preferably during a meal. Your doctor may recommend to take UPTRAVI only once daily in certain circumstances. Talk to your doctor before taking UPTRAVI if:

- You are taking other medicines, e.g.:
 - Clopidogrel (medicine used to inhibit blood clots in coronary artery disease)
 - Deferasirox (medicine used to remove iron from the blood stream)
 - Teriflunomide (medicine used to treat relapsing-remitting multiple sclerosis)
- You have or have had problems with your liver not working properly

There are two phases of treatment with UPTRAVI:

TITRATION

In the first several weeks, you will work with your doctor to find the dose of UPTRAVI that is right for you. Your doctor may have you step up from the starting dose to higher doses of UPTRAVI. Your doctor may step you down to a lower dose. This process is called titration. It lets your body gradually adjust to the medicine.

MAINTENANCE

Once your doctor has found the dose that is right for you, this will be the dose you take on a regular basis. This is called the maintenance dose.

HOW SHOULD YOU STEP UP YOUR DOSE?

You will start at the 200 microgram dose in the morning and in the evening, and after discussing with your doctor or nurse, step up to the next dose. The first intake of the increased dose should be in the evening. Each step usually lasts about 1 week. It could take several weeks to find the dose that is right for you.

The goal is to reach the dose that is most appropriate to treat you. This dose will be your maintenance dose.

Every patient with PAH is different. **Not everyone will end up on the same maintenance dose.**

Some patients may have 200 micrograms in the morning and in the evening as their maintenance dose, while some will reach the highest dose of 1,600 micrograms in the morning and in the evening. Others may reach a maintenance dose somewhere in between. What is important is that you reach the dose that is most appropriate to treat you.

WHAT ARE THE STEPS?

200 microgram tablet





800 microgram tablet (Use with steps 5 through 8 to reduce the number of tablets needed per dose.) HIGHEST DOSE

STEP 5 Morning: One 800 and one 200 microgram tablet Evening:

One 800 and one 200 microgram tablet (total daily dose: 2,000 microgram)

STEP 6 Morning: One 800 and two 200 microgram tablets Evening: One 800 and two 200 microgram tablets (total daily dose:

2,400 microgram)

STEP 7 Morning: One 800 and three 200 microgram tablets Evening: One 800 and three 200 microgram tablets (total daily dose: 2.800 microgram)

STEP 8 Morning:

One 800 and four 200 microgram tablets **Evening:**

One 800 and four 200 microgram tablets (total daily dose: 3,200 microgram)

Each dosing step lasts about 1 week.

As with all medicines, you may experience side effects with UPTRAVI as you step up to higher doses. Talk to your doctor or nurse if you have side effects. There are treatments available that can help relieve them.

The most common side effects (may affect more than 1 in 10 people) you may experience while taking UPTRAVI are:

- Headache Diarrhoea Nausea Vomiting Jaw pain Muscle pain
- Leg pain Joint pain Facial redness Nasopharyngitis (stuffy nose)

For a full list of side effects see package leaflet for further information.

If you cannot tolerate the side effects even after your doctor or nurse has tried

to treat them, he or she may recommend you step down to a lower dose.

If your doctor or nurse tells you to step down to a lower dose, you should take one less 200 microgram tablet in the morning and one less in the evening.

You should only step down after speaking with your PAH doctor or nurse. This stepping-down process will help you find the dose that is right for you, also called your maintenance dose.

AN EXAMPLE OF STARTING & STEPPING DOWN TREATMENT

Each dosing step lasts about 1 week.





Each dosing step lasts about 1 week.

800 microgram tablet (Use with steps 5 through 8 to reduce the number of tablets needed per dose.)



WHEN YOU MOVE TO YOUR MAINTENANCE DOSE

The highest dose that you can tolerate during titration will become your **maintenance dose.** Your maintenance dose is the dose you should continue to take on a regular basis. Your doctor or nurse can prescribe an equivalent **single-tablet strength** for your maintenance dose.

This lets you take just one tablet in the morning and one in the evening, instead of multiple tablets for each dose.

For example, if your highest tolerated dose during titration was 1,200 micrograms once in the morning and once in the evening:



Over time, your doctor or nurse may adjust your maintenance dose as needed.

IF YOU FORGET TO TAKE UPTRAVI

If you miss a dose, take the dose as soon as you remember, then continue to take your tablets at the usual times. If it is within 6 hours of when you would normally take your next dose, you should skip the missed dose and continue to take your medicine at the usual time.

Do not take a double dose to make up for a forgotten tablet.

Do not stop taking UPTRAVI unless your doctor or nurse tells you to. If, for any reason, you stop taking UPTRAVI for more than 3 consecutive days (if you missed six doses in a row), **contact your PAH doctor or nurse immediately as your dose may need to be adjusted to avoid side effects.**

Your doctor or nurse may have you resume treatment at a lower dose, gradually increasing to your previous maintenance dose.

Please read the instructions in the package leaflet carefully.

The following diary pages help you keep track of the number of tablets you need to take in the morning and evening during titration. Use them to write down the number of tablets you take in the morning and the evening.

Each step usually lasts about 1 week, unless your doctor or nurse instructs you otherwise. If your titration steps last longer than 1 week there are additional diary pages to track this.



Use pages 22 to 31 to track the first weeks of treatment, when you are using 200 microgram tablets only (steps 1–4).



If you have been prescribed both 200 and 800 microgram tablets, use pages 32 to 41 (steps 5–8).


Remember to talk to your PAH doctor or nurse regularly.

Write down your doctor or nurse's instructions:

Doctor's office telephone and e-mail:

Pharmacist's telephone:

Notes:



WEEK Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY




The first intake of an increased dose of UPTRAVI should be in the evening



WEEK Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY



WEEK Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY



WEEK Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY



WEEK

Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY



WEEK Every day write down in the boxes below how many tablets you take in the morning and evening. I spoke with my doctor or nurse on DD/MM/YY



Use the following diary pages if your doctor or nurse prescribes 800 microgram tablets in addition to your 200 microgram tablets.

On the diary pages, check off that you have taken **one** 800 microgram tablet every day in the morning and in the evening with your prescribed number of 200 microgram tablets.



200 microgram tablets



800 microgram tablet

(Use with steps 5 through 8 to reduce the number of tablets needed per dose.)


Remember to talk to your PAH doctor or nurse regularly.

Write down your doctor or nurse's instructions:


Doctor's office telephone and e-mail:

Pharmacist's telephone:


Notes:


















NOTES

You can help by reporting any side effects you may get. If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any side effects not listed in the package leaflet. You can also report side effects via HPRA Pharmacovigilance, website www.hpra.ie.

 Marketing Authorisation Holder: JANSSEN-CILAG INTERNATIONAL NV, Turnhoutseweg 30, B-2340 Beerse, Belgium

 EM-83143
 December 2021
 © 2021 Janssen Sciences Ireland UC