



WITH VITAMIN C TABLETS

Paracetamol 500mg • Diphenhydramine Hydrochloride 15mg
Phenylephrine Hydrochloride 10mg • Caffeine 30mg
Ascorbic Acid 300mg

Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to take Uniflu with Vitamin C Tablets carefully to get the best results.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve after 3 days.
- If any of the side effects gets serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Uniflu with Vitamin C Tablets are and what they are used for
2. Before you take Uniflu with Vitamin C Tablets
3. How to take Uniflu with Vitamin C Tablets
4. Possible side effects
5. How to store Uniflu with Vitamin C Tablets
6. Further information

1. What Uniflu with Vitamin C Tablets are and what they are used for

Uniflu Tablets are for the symptomatic relief from the discomfort associated with influenza and colds i.e. nasal and sinus congestion, headache, fever, aching limbs, coughing and running nose.

Vitamin C Tablets are used as a vitamin C supplement during cold and influenza (flu) infections.

2. Before you take Uniflu with Vitamin C Tablets

Do not take Uniflu Tablets

- if you are already taking other paracetamol-containing products
- if you are allergic to paracetamol or any of the other ingredients
- if you have an over-active thyroid, high blood pressure or heart disease
- if you are currently taking monoamine oxidase inhibitors (MAOIs) (a type of antidepressant) or within 14 days of stopping such treatment
- if you are taking other sympathomimetic drugs (stimulants or appetite suppressants and drugs used to treat congestion and asthma)
- if you are suffering from persistent or chronic cough such as occurs with smoking, asthma or emphysema (serious lung condition causing shortness of breath), or if cough is accompanied by excessive mucus (phlegm).
- if you have glaucoma (high pressure in the eye)
- if you suffer from urinary retention (difficulty passing urine)
- if you have breathing problems

Do not take Vitamin C Tablets:

- If you are allergic to ascorbic acid or any of the other ingredients

Take special care with Uniflu with Vitamin C Tablets

Talk to your doctor before taking Uniflu with Vitamin C Tablets:

- if you have liver problems, such as alcoholic liver disease - you have an increased risk of overdose

- if you have kidney problems
- if you have an enlarged prostate gland causing difficulty in passing urine
- if you suffer from epilepsy
- if you suffer from angina.

Taking other medicines

Please tell your doctor or pharmacist before taking Uniflu with Vitamin C Tablets if you are taking or have recently taken any other medicines, including:

- blood thinning tablets (warfarin and other coumarins)
- chloramphenicol (antibiotic)
- cholestyramine (cholesterol lowering medication)
- phenothiazines or antidepressants
- medicines to treat high blood pressure
- medicines to treat nausea and vomiting (such as metoclopramide and domperidone)
- heart medication such as digitalis, beta-adrenergic blockers, guanethidine, reserpine or methyldopa
- a general anaesthetic
- atropine
- medicines obtained without a prescription
- flucloxacillin (antibiotic), due to a serious risk of blood and fluid abnormality (high anion gap metabolic acidosis) that must have urgent treatment and which may occur particularly in case of severe renal impairment, sepsis (when bacteria and their toxins circulate in the blood leading to organ damage), malnutrition, chronic alcoholism, and if the maximum daily doses of paracetamol are used.

Taking Uniflu with Vitamin C Tablets with food and drink

The sedative effect of Uniflu Tablets is increased by alcohol. Therefore, alcoholic drinks should be avoided whilst you are taking Uniflu with Vitamin C Tablets.

Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Uniflu Tablets may make you feel sleepy. If affected:

- do not drive because Uniflu Tablets could stop you driving safely
- do not use any tools or machines.

Important information about some of the ingredients of Uniflu with Vitamin C Tablets

Uniflu Tablets contain Carmoisine (E122) which may cause allergic reactions.

Vitamin C Tablets contain sucrose and lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Uniflu with Vitamin C Tablets

Adults and Elderly:	Swallow 1 Uniflu tablet and 1 Vitamin C tablet (swallow, suck or chew) with water every six hours. Do not take more than 4 tablets in 24 hours.
Children aged 12 and over:	Swallow 1 Uniflu tablet and 1 Vitamin C tablet (swallow, suck or chew) with water every eight hours. Do not take more than 3 tablets in 24 hours.
Children under 12 years:	Not recommended.

If you do not feel any better after taking this medicine for 3 days, consult your doctor or pharmacist.

If you take more Uniflu with Vitamin C Tablets than you should
Immediate medical advice should be sought in the event of an overdose, even if you feel well, because of the risk of delayed, serious liver damage.

Overdose may lead to increased heart rate, high blood pressure, sickness, liver damage and breathing problems.

If you forget to take Uniflu with Vitamin C Tablets

Do not take a double dose to make up for a missed dose.

4. Possible side effects

Like all medicines, Uniflu with Vitamin C Tablets can cause side effects, although not everybody gets them.

Tell your doctor if you notice any of the following mild side effects:

- difficulty passing urine
- headache
- dry mouth
- blurred vision
- stomach upsets (nausea, diarrhoea, abdominal cramps, stomach pains)
- tremors
- rashes
- high blood pressure.
- blood disorders which can lead to a reduction in blood platelets, which increases risk of bleeding or bruising (thrombocytopenia), or reduction in the number of white blood cells which makes infections more likely (agranulocytosis).
- insomnia and nervousness
- very rare cases of serious skin reactions have been reported.

Reporting of side effect

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL – Dublin; Tel: +353 1 6764971; Fax: +353 1 6762517; Website: www.hpra.ie; E-mail: medsafety@hpra.ie

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Uniflu with Vitamin C Tablets

Keep out of the sight and reach of children.

Do not take Uniflu Tablets with Vitamin C after the expiry date which is stated on the packaging.

Do not store above 25°C. Store in the original package to protect from moisture. Do not use if foil strip is damaged.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Further information

What Uniflu with Vitamin C Tablets contain:

The active substances per Uniflu Tablet:

Paracetamol500 mg
Diphenhydramine hydrochloride 15 mg
Phenylephrine hydrochloride 10 mg
Caffeine30 mg

The other ingredients are:

Core: Acacia; Protein S (Byco C) (Hydrolysed Gelatin); Alginate Acid (E400); Magnesium Stearate; Stearic Acid; Sodium Starch Glycolate (Type A)

Coating: Opadry II Pink (includes Polyvinyl Alcohol, Talc (E553b), Titanium Dioxide (E171), Macrogol, Lecithin, Carmoisine Aluminium Lake, FD&C Blue Indigo Carmine Aluminium Lake (E132)).

The active substances per Vitamin C Tablet:

Ascorbic Acid (Vitamin C) 300 mg
(as Ascorbic Acid /Sodium Ascorbate)

The other ingredients are:

Saccharin Sodium; Stearic Acid; Magnesium Stearate; Compressible Sugar; Quinoline Yellow Lake 66154 (E104); Hypromellose; Pectin; Lemon Flavour (E2954) (includes Maltodextrin, Acacia, Lactose).

What Uniflu with Vitamin C Tablets look like and contents of the pack

Uniflu Tablets are pink, oblong, film coated tablets. They are only available in combined blister packs with Vitamin C Tablets.

Vitamin C Tablets are a yellow, biconvex, circular, plain, lemon flavoured, chewable tablet.

There are 6 dose packs (containing 6 Uniflu Tablets and 6 Vitamin C Tablets) or 12 dose packs (containing 12 Uniflu Tablets and 12 Vitamin C Tablets).

Marketing Authorisation Holder

PHOENIX LABS
Suite 12, Bunkilla Plaza
Bracetown Business Park,
Clonee, Co. Meath


Manufacturer:

SIDEFARMA
Sociedade Industrial de Expansão Farmacêutica, SA
Rua da Guiné, 26
2689 - 514 Prior Velho
Portugal

Date of review of Patient Information Leaflet

Date of revision June 2023

Legal Status

 Pharmacy-Only-Medicine

Marketing Authorisation Number

Ireland: PA 1113/5/1