

## Package leaflet: information for the patient

### Risperidone 0.5mg, 1mg, 2mg, 3mg, 4mg, and 6mg film-coated Tablets

Risperidone

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, nurse or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor, nurse or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4

#### **What is in this leaflet**

1. What Risperidone Tablets are and what they are used for
2. What do you need to know before you take Risperidone Tablets
3. How to take Risperidone Tablets
4. Possible side effects
5. How to store Risperidone Tablets
6. Contents of the pack and other information

#### **1. What Risperidone Tablets are and what they are used for**

Risperidone belongs to a group of medicines called antipsychotics.

Risperidone is used to treat the following:

- Schizophrenia, where you may see, hear or feel things that are not there, believe things that are not true or feel unusually suspicious, or confused.
- Mania, where you may feel very excited, elated, agitated, enthusiastic or hyperactive. Mania occurs in an illness called “bipolar disorder”.
- Short-term treatment (up to six weeks) of long-term aggression in people with Alzheimer’s dementia, who harm themselves or others. Alternative (non-drug) treatments should have been used previously.
- Short-term treatment (up to six weeks) of long-term aggression in intellectually disabled children (at least five years of age) and adolescents with conduct disorder.

Risperidone can help alleviate the symptoms of your disease and stop your symptoms from coming back.

#### **2. What do you need to know before you take Risperidone Tablets**

##### **Do not take Risperidone Tablets:**

If you are allergic to risperidone or any of the other ingredients of this medicine (listed in section 6) If you are not sure if the above applies to you, talk to your doctor or pharmacist before using Risperidone Tablets.

#### **Warnings and precautions**

Talk to your doctor or pharmacist before taking Risperidone Tablets if:

- you have a heart problem. Examples include an irregular heart rhythm or if you are prone to low blood pressure or if you are using medicines for your blood pressure. Risperidone Tablets may cause low blood pressure. Your dose may need to be adjusted
- you know of any factors which would favour you having a stroke, such as high blood pressure, cardiovascular disorder or blood vessel problems in the brain
- you have ever experienced involuntary movements of the tongue, mouth and face

- you have ever had a condition whose symptoms include high temperature, muscle stiffness, sweating or a lowered level of consciousness (also known as Neuroleptic Malignant Syndrome)
- you have Parkinson's disease or dementia
- you know that you have had low levels of white blood cells in the past (which may or may not have been caused by other medicines)
- you are diabetic
- you have epilepsy
- you are a man and you have ever had a prolonged or painful erection
- you have problems controlling your body temperature or overheating
- you have kidney problems
- you have liver problems
- you have an abnormally high level of the hormone prolactin in your blood or if you have a possible prolactin-dependent tumour
- you or someone else in your family has a history of blood clots, as antipsychotics have been associated with formation of blood clots.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking Risperidone Tablets.

As dangerously low numbers of a certain type of white blood cell needed to fight infection in your blood has been seen very rarely with patients taking risperidone, your doctor may check your white blood cell counts.

Risperidone Tablets may cause you to gain weight. Significant weight gain may adversely affect your health. Your doctor should regularly measure your body weight.

As diabetes mellitus or worsening of pre-existing diabetes mellitus have been seen with patients taking risperidone, your doctor should check for signs of high blood sugar. In patients with pre-existing diabetes mellitus blood glucose should be monitored regularly.

Risperidone commonly raises levels of a hormone called "prolactin". This may cause side effects such as menstrual disorders or fertility problems in women, breast swelling in men (see Possible side effects). If such side effects occur, evaluation of the prolactin level in the blood is recommended.

During an operation on the eye for cloudiness of the lens (cataract), the pupil (the black circle in the middle of your eye) may not increase in size as needed. Also, the iris (the coloured part of the eye) may become floppy during surgery and that may lead to eye damage. If you are planning to have an operation on your eye, make sure you tell your eye doctor that you are taking this medicine.

### **Elderly people with dementia**

In elderly patients with dementia, there is an increased risk of stroke. You should not take risperidone if you have dementia caused by stroke.

During treatment with risperidone you should frequently see your doctor.

Medical treatment should be sought straight away if you or your care-giver notice a sudden change in your mental state or sudden weakness or numbness of your face, arms or legs, especially on one side, or slurred speech, even for a short period of time. These may be signs of a stroke.

### **Children and adolescents**

Before treatment is started for conduct disorder, other causes of aggressive behaviour should have been ruled out.

If during treatment with risperidone tiredness occurs, a change in the time of administration might improve attention difficulties.

Before treatment is started your, or your child's body weight may be measured and it may be regularly monitored during treatment.

A small and inconclusive study has reported an increase in height in children who took risperidone, but whether this is an effect of the drug or due to some other reason is not known.

### **Other medicines and Risperidone Tablets**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

#### **It is especially important to talk to your doctor or pharmacist if you are taking any of the following:**

- medicines that work on your brain such as to help you calm down (benzodiazepines) or some medicines for pain (opiates), medicines for allergy (some antihistamines), as risperidone may increase the sedative effect of all of these
- medicines that may change the electrical activity of your heart, such as medicines for malaria, heart rhythm problems, allergies (antihistamines), some antidepressants or other medicines for mental problems
- medicines that cause a slow heart beat
- medicines that cause low blood potassium (such as certain diuretics)
- medicines to treat raised blood pressure. Risperidone can lower blood pressure
- medicines for Parkinson's disease (such as levodopa)
- water tablets (diuretics) used for heart problems or swelling of parts of your body due to a build up of too much fluid (such as furosemide or chlorothiazide). Risperidone taken by itself or with furosemide, may have an increased risk of stroke or death in elderly people with dementia
- medicines that increase the activity of the central nervous system (psychostimulants, such as methylphenidate).

Risperidone taken by itself or with furosemide, may have an increased risk of stroke or death in elderly people with dementia.

The following medicines may reduce the effect of risperidone:

- rifampicin (a medicine for treating some infections)
- carbamazepine, phenytoin (medicines for epilepsy)
- phenobarbital.

If you start or stop taking such medicines you may need a different dose of risperidone

The following medicines may increase the effect of risperidone:

- quinidine (used for certain types of heart disease)
- antidepressants such as paroxetine, fluoxetine, tricyclic antidepressants
- medicines known as beta blockers (used to treat high blood pressure)
- phenothiazines (such as medicines used to treat psychosis or to calm down)
- cimetidine, ranitidine (blockers of the acidity of stomach)
- itraconazole and ketoconazole (medicines for treating fungal infections)
- certain medicines used in the treatment of HIV/AIDS, such as ritonavir
- verapamil, a medicine used to treat high blood pressure, and/or abnormal heart rhythm
- sertraline and fluvoxamine, medicines used to treat depression and other psychiatric disorders.

If you start or stop taking such medicines you may need a different dose of risperidone

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before using risperidone.

### **Risperidone Tablets with food, drink and alcohol**

You can take this medicine with or without food. You should avoid drinking alcohol when taking Risperidone.

### **Pregnancy, breast-feeding and fertility**

- If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. Your doctor will decide if you can take it.

- The following symptoms may occur in newborn babies, of mothers that have used risperidone in the last trimester (last three months of their pregnancy): shaking, muscle stiffness and/or weakness, sleepiness, agitation, breathing problems, and difficulty in feeding. If your baby develops any of these symptoms you may need to contact your doctor.

- Risperidone can raise your levels of a hormone called "prolactin" that may impact fertility (see Possible side effects)

### **Driving and using machines**

Dizziness, tiredness, and vision problems may occur during treatment with Risperidone Tablets. Do not drive or use any tools or machines without talking to your doctor first.

### **Risperidone Tablets contain lactose.**

This product contains lactose, a type of sugar. If you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking this medicinal product.

The 2mg and 6mg tablets contain sunset yellow FCF (E110) this may cause allergic reactions in some patients.

### **3. How to take Risperidone Tablets**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

#### **The recommended dose is as follows:**

#### **For the treatment of schizophrenia**

##### **Adults**

- The usual starting dose is 2mg per day, this may be increased to 4mg per day on the second day
- Your dose may then be adjusted by your doctor depending on how you respond to the treatment
- Most people feel better with daily doses of 4 to 6mg
- This total daily dose can be divided into either one or two doses a day. Your doctor will tell you which is the best for you.

##### **Elderly people**

- Your starting dose will normally be 0.5mg twice a day
- Your dose may then be gradually increased by your doctor to 1mg to 2mg twice a day
- Your doctor will tell you which is the best for you.

#### **For the treatment of mania**

##### **Adults**

- Your starting dose will usually be 2mg once a day
- Your dose may then be gradually adjusted by your doctor depending on how you respond to the treatment
- Most people feel better with doses of 1 to 6mg once a day.

##### **Elderly people**

- Your starting dose will usually be 0.5mg twice a day
- Your dose may then be gradually adjusted by your doctor to 1mg to 2mg twice a day depending on how much you respond to the treatment.

#### **For the treatment of long-standing aggression in people with Alzheimer's dementia**

##### **Adults (including elderly people)**

- Your starting dose will normally be 0.25mg twice a day
- Your dose may then be gradually adjusted by your doctor depending on how you respond to the

treatment

- Most people feel better with 0.5mg twice a day. Some patients may need 1mg twice a day  
Treatment duration in patients with Alzheimer's dementia should be not more than six weeks.

### **Use in children and adolescents**

Children and adolescents under 18 years old should not be treated with Risperidone for schizophrenia or mania.

### **For the treatment of conduct disorder**

The dose will depend on your child's weight

For children who weigh less than 50kg

- The starting dose will normally be 0.25mg once a day
- The dose may be increased every other day in steps of 0.25mg per day
- The usual maintenance dose is 0.25mg to 0.75mg once a day.

For children who weigh 50kg or more

- The starting dose will normally be 0.5mg once a day
- The dose may be increased every other day in steps of 0.5mg per day
- The usual maintenance dose is 0.5mg to 1.5mg once a day.

Treatment duration in patients with conduct disorder should be not more than six weeks.

Children under 5 years old should not be treated with Risperidone Tablets for conduct disorder.

### **People with kidney or liver problems**

Regardless of the disease to be treated, all starting doses and following doses of risperidone should be halved. Dose increases should be slower in these patients.

Risperidone should be used with caution in this patient group.

### **Method of administration**

#### **For oral use**

You should swallow your tablet with a drink of water.

### **If you take more Risperidone Tablets than you should**

- See a doctor right away. Take the medicine pack with you
- In case of overdose you may feel sleepy or tired, or have abnormal body movements, problems standing and walking, feel dizzy due to low blood pressure, or have abnormal heart beats or fits.

### **If you forget to take Risperidone Tablets**

- If you forget to take a dose, take it as soon as you remember it. However, if it is almost time for your next dose, skip the missed dose and continue as usual. If you miss two or more doses, contact your doctor
- **Do not take a double dose (two doses at the same time) to make up for a forgotten dose.**

### **If you stop taking Risperidone Tablets**

You should not stop taking this medicine unless told to do so by your doctor. Your symptoms may return.

If your doctor decides to stop this medicine, your dose may be decreased gradually over a few days. If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. POSSIBLE SIDE EFFECTS**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Tell your doctor immediately if you experience any of the following uncommon side effects (may affect up to 1 in 100 people):**

- Have dementia and experience a sudden change in your mental state or sudden weakness or numbness of your face, arms or legs, especially on one side, or slurred speech, even for a short period of time. These may be signs of a stroke
- Experience tardive dyskinesia (twitching or jerking movements that you cannot control in your face, tongue, or other parts of your body). Tell your doctor immediately if you experience involuntary rhythmic movements of the tongue, mouth and face. Withdrawal of risperidone may be needed.

**Tell your doctor immediately if you experience any of the following rare side effects (may affect up to 1 in 1,000 people):**

- Experience blood clots in the veins, especially in the legs (symptoms include swelling, pain and redness in the leg), which may travel through blood vessels to the lungs causing chest pain and difficulty breathing. If you notice any of these symptoms seek medical advice immediately
- Experience fever, muscle stiffness, sweating or a lowered level of consciousness (a disorder called "Neuroleptic Malignant Syndrome"). Immediate medical treatment may be needed
- Are a man and experience prolonged or painful erection. This is called priapism. Immediate medical treatment may be needed
- Experience severe allergic reaction characterised by fever, swollen mouth, face, lip or tongue, shortness of breath, itching, skin rash or drop in blood pressure.

The following side effects may happen:

**Very common (may affect more than 1 in 10 people):**

- Difficulty falling or staying asleep
- Parkinsonism. This condition may include: slow or impaired movement, sensation of stiffness or tightness of the muscles (making your movements jerky), and sometimes even a sensation of movement "freezing up" and then restarting. Other signs of parkinsonism include a slow shuffling walk, a tremor while at rest, increased saliva and/or drooling, and a loss of expression on the face.
- Feeling sleepy, or less alert
- Headache.

**Common (may affect up to 1 in 10 people):**

- Pneumonia, infection of the chest (bronchitis), common cold symptoms, sinus infection, urinary tract infection, ear infection, feeling like you have the flu
- Raised levels of a hormone called "prolactin" found in a blood test (which may or may not cause symptoms). Symptoms of high prolactin occur uncommonly and may include in men breast swelling, difficulty in getting or maintaining erections, decreased sexual desire or other sexual dysfunction. In women they may include breast discomfort, leakage of milk from the breasts, missed menstrual periods, or other problems with your cycle or fertility problems.
- Weight gain, increased appetite, decreased appetite
- Sleep disorder, irritability, depression, anxiety, restlessness
- Dystonia: This is a condition involving slow or sustained involuntary contraction of muscles. While it can involve any part of the body (and may result in abnormal posture), dystonia often involves muscles of the face, including abnormal movements of the eyes, mouth, tongue or jaw.
- Dizziness
- Dyskinesia: This is a condition involving involuntary muscle movements, and can include repetitive, spastic or writhing movements, or twitching.
- Tremor (shaking)
- Blurry vision, eye infection or "pink eye"
- Rapid heart rate, high blood pressure, shortness of breath
- Sore throat, cough, nosebleeds, stuffy nose

- Abdominal pain, abdominal discomfort, vomiting, nausea, constipation, diarrhea, indigestion, dry mouth, toothache
- Rash, skin redness
- Muscle spasms, bone or muscle ache, back pain, joint pain
- Incontinence (lack of control) of urine
- Swelling of the body, arms or legs, fever, chest pain, weakness, fatigue (tiredness), pain
- Fall.

**Uncommon (may affect up to 1 in 100 people):**

- Infection of the breathing passages, bladder infection, 'eye infection, tonsillitis, fungal infection of the nails, infection of the skin, an infection confined to a single area of skin or part of the body, viral infection, skin inflammation caused by mites
- Decrease in the type of white blood cells that help to protect you against infection, white blood cell count decreased, decrease in platelets (blood cells that help you stop bleeding), anemia, decrease in red blood cells, increase in eosinophils (a type of white blood cell) in your blood
- Allergic reaction
- Diabetes or worsening of diabetes, high blood sugar, excessive drinking of water
- Weight loss, loss of appetite resulting in malnutrition and low body weight
- Increased cholesterol in your blood
- Elated mood (mania), confusion, decreased sexual drive, nervousness, nightmares
- Unresponsive to stimuli, loss of consciousness, low level of consciousness
- Convulsion (fits), fainting
- A restless urge to move parts of your body, balance disorder, abnormal coordination, dizziness upon standing, disturbance in attention, problems with speech, loss or abnormal sense of taste, reduced sensation of skin to pain and touch, a sensation of tingling, pricking, or numbness skin
- Oversensitivity of the eyes to light, dry eye, increased tears, redness of the eyes
- Sensation of spinning (vertigo), ringing in the ears, ear pain
- Atrial fibrillation (an abnormal heart rhythm), an interruption in conduction between the upper and lower parts of the heart, abnormal electrical conduction of the heart, prolongation of the QT interval from your heart, slow heart rate, abnormal electrical tracing of the heart (electrocardiogram or ECG), a fluttering or pounding feeling in your chest (palpitations)
- Low blood pressure, low blood pressure upon standing (consequently, some people taking risperidone may feel faint, dizzy, or may pass out when they stand up or sit up suddenly, flushing
- Pneumonia caused by inhaling food, lung congestion, congestion of breathing passages, crackly lung sounds, wheezing, voice disorder, breathing passage disorder
- Stomach or intestinal infection, stool incontinence, very hard stool, difficulty swallowing, excessive passing of gas or wind
- Hives (or "nettle rash"), itching, hair loss, thickening of skin, eczema, dry skin, skin discoloration, acne, flaky, itchy scalp or skin, skin disorder, skin lesion
- An increase of CPK (creatine phosphokinase) in your blood, an enzyme which is sometimes released with muscle breakdown
- Abnormal posture, joint stiffness, joint swelling, muscle weakness, neck pain
- Frequent passing of urine, inability to pass urine, pain when passing urine
- Erectile dysfunction, ejaculation disorder
- Loss of menstrual periods, missed menstrual periods or other problems with your cycle (females)
- Development of breasts in men, leakage of milk from the breasts, sexual dysfunction, breast pain, breast discomfort, vaginal discharge
- Swelling of the face, mouth, eyes, or lips
- Chills, an increase in body temperature
- A change in the way you walk
- Feeling thirsty, feeling unwell, chest discomfort, Feeling "out of sorts", discomfort
- Increased liver transaminases in your blood, increased GGT (a liver enzyme called gamma-glutamyltransferase) in your blood, increased liver enzymes in your blood
- Procedural pain.

**Rare (may affect up to 1 in 1,000 people):**

- Infection

- Inappropriate secretion of a hormone that controls urine volume
- Sleep walking
- Sleep-related eating disorder
- Sugar in the urine, low blood sugar, high blood triglycerides (a fat)
- Lack of emotion, inability to reach orgasm
- Neuroleptic malignant syndrome (confusion, reduced or loss of consciousness, high fever, and severe muscle stiffness)
- Blood vessel problems in the brain
- Coma due to uncontrolled diabetes
- Shaking of the head
- Glaucoma (increased pressure within the eyeball), problems with movement of your eyes, eye rolling, eyelid margin crusting
- Eye problems during cataract surgery. During cataract surgery, a condition called intraoperative floppy iris syndrome (IFIS) can happen if you take or have taken risperidone. If you need to have cataract surgery, be sure to tell your eye doctor if you take or have taken this medicine
- Dangerously low numbers of a certain type of white blood cell needed to fight infection in your blood
- Dangerously excessive intake of water
- Irregular heart beat
- Trouble breathing during sleep (sleep apnea), fast, shallow breathing
- Inflammation of the pancreas, a blockage in the bowels
- Swollen tongue, chapped lips, rash on skin related to drug
- Dandruff
- Breakdown of muscle fibers and pain in muscles (rhabdomyolysis)
- A delay in menstrual periods, enlargement of the glands in your breasts, breast enlargement, discharge from the breasts
- Increased insulin (a hormone that controls blood sugar levels) in your blood
- Hardening of the skin
- Decreased body temperature, coldness in arms and legs
- Symptoms of drug withdrawal
- Yellowing of the skin and the eyes (jaundice).

**Very rare (may affect up to 1 in 10,000 people):**

- Life threatening complications of uncontrolled diabetes
- Serious allergic reaction with swelling that may involve the throat and lead to difficulty breathing
- Lack of bowel muscle movement that causes blockage.

The following side effect has been seen with the use of another medicine called paliperidone that is very similar to risperidone, so these can also be expected with risperidone: Rapid heartbeat upon standing.

**Additional side effects in children and adolescents**

In general, side effects in children are expected to be similar to those in adults.

The following side effects were reported more often in children and adolescents (5 to 17 years) than in adults: feeling sleepy, or less alert, fatigue (tiredness), headache, increased appetite, vomiting, common cold symptoms, nasal congestion, abdominal pain, dizziness, cough, fever, tremor (shaking), diarrhoea, and incontinence (lack of control) of urine.



## Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting systems listed below.

### United Kingdom

Yellow Card Scheme

[www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

### Ireland

HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517; Website: [www.hpra.ie](http://www.hpra.ie) ; e-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie)

By reporting side effects you can help provide more information on the safety of this medicine.

## 5. HOW TO STORE RISPERIDONE TABLETS

Keep this medicine out of the sight and reach of children.

Store the tablets below 30°C.

Store in the original package.

Do not use this medicine after the expiry date which is stated on the blister and the carton.

The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

## 6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Risperidone Tablets contain

The active substance is risperidone. Each tablet contains 0.5mg, 1mg, 2mg, 3mg, 4mg or 6mg risperidone.

Other ingredients are sodium laurilsulphate, lactose monohydrate, microcrystalline cellulose, sodium starch glycolate, colloidal anhydrous silica, hypromellose and magnesium stearate.

The tablets also contain:

0.5mg: hypromellose 6 cP, titanium dioxide (E171), talc, propylene glycol (E1520) and iron oxide red (E172)

1mg: hypromellose 15 cP, propylene glycol (E1520)

2mg: titanium dioxide (E171), hypromellose 15 cP, propylene glycol (E1520), talc, sunset yellow FCF (E110)

3mg: hypromellose, quinoline yellow (E104), titanium dioxide (E171), propylene glycol (E1520), talc, yellow iron oxide (E172)

4mg: hypromellose, quinoline yellow (E104), propylene glycol (E1520), titanium dioxide (E171), talc, indigo carmine (E132)

6mg: hypromellose, titanium dioxide (E171), quinolone yellow (E104), propylene glycol (E1520), talc, sunset yellow FCF (E110)

What Risperidone Tablets look like and contents of the pack

0.5mg: brown, capsule-shaped, film-coated tablets with breakline on one side and plain on the other side – 20 tablet pack

1mg: white, capsule-shaped, film-coated tablets with breakline on one side and plain on the other side – 20 or 60 tablet pack

2mg: orange, capsule-shaped, film-coated tablets with breakline on one side and plain on the other side – 60 tablet pack

3mg: yellow, capsule-shaped, film-coated tablets with breakline on one side and plain on the other side – 60 tablet pack

4mg: green, capsule-shaped, film-coated tablets with breakline on one side and plain on the other side  
– 60 tablet pack

6mg: orange, round, film-coated tablets – 28 tablet pack

The 0.5mg, 1mg, 2mg, 3mg & 4mg tablets can be divided into equal halves.

### **Risperidone Tablets – PIL Information**

To listen to or request a copy of this leaflet in Braille, large print or audio please call, free of charge:

**0800 198 5000 (UK only)**

Please be ready to give the following information:

<b>Product name</b>	<b>Reference number</b>
Risperidone 0.5mg Film-Coated Tablets	29831/0343
Risperidone 1mg Film-Coated Tablets	29831/0344
Risperidone 2mg Film-Coated Tablets	29831/0345
Risperidone 3mg Film-Coated Tablets	29831/0346
Risperidone 4mg Film-Coated Tablets	29831/0347
Risperidone 6mg Film-Coated Tablets	29831/0348

This is a service provided by the Royal National Institute of Blind People.

For the Republic of Ireland please call +44 1978 661 261

### **Marketing Authorisation Holder and Manufacturer**

**Marketing Authorisation Holder:** Wockhardt UK Ltd, Ash Road North, Wrexham, LL13 9UF, UK

**Manufacturer:** CP Pharmaceuticals Ltd, Ash Road North, Wrexham, LL13 9UF, UK

**This leaflet was last revised in 11/2018.**