

**PACKAGE LEAFLET**

## **Package leaflet: Information for the patient**

### **Risedronate Pinewood Once a week 35 mg film-coated tablets** risedronate sodium

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Risedronate Pinewood is and what it is used for
2. What you need to know before you take Risedronate Pinewood
3. How to take Risedronate Pinewood
4. Possible side effects
5. How to store Risedronate Pinewood
6. Contents of the pack and other information

## **1. What Risedronate Pinewood is and what it is used for**

### **What Risedronate Pinewood is**

Risedronate Pinewood belongs to a group of non-hormonal medicines called bisphosphonates which are used to treat bone diseases. It works directly on your bones to make them stronger and therefore less likely to break.

Bone is a living tissue. Old bone is constantly removed from your skeleton and replaced with new bone.

Postmenopausal osteoporosis is a condition occurring in women after the menopause where the bones become weaker, more fragile and more likely to break after a fall or strain.

Osteoporosis can also occur in men due to a number of causes including ageing and/or a low level of the male hormone, testosterone.

Osteoporosis is more likely to occur in women who have reached the menopause early and also in patients treated long-term with steroids.

The spine, hip and wrist are the most likely bones to break, although this can happen to any bone in your body. Osteoporosis related fractures can also cause back pain, height loss and a curved back. Many patients with osteoporosis have no symptoms and you may not even have known that you had it.

### **What Risedronate Pinewood is used for**

The treatment of osteoporosis:

- in **postmenopausal women**, even if osteoporosis is severe. It reduces the risk of spinal and hip fractures.
- in men at high risk of fractures.

## **2. What you need to know before you take Risedronate Pinewood**

### **Do not take Risedronate Pinewood**

- if you are **allergic** to risedronate sodium or any of the other ingredients of this medicine (listed in section 6).
- if your doctor has told you that you have a condition called **hypocalcaemia** (a low blood calcium level).
- if you may be **pregnant**, are pregnant or are planning to become pregnant.
- if you are **breast-feeding**.
- if you have **severe kidney problems**.

### **Warnings and precautions**

Talk to your doctor or pharmacist before taking Risedronate Pinewood

- If you are unable to stay in an upright position (sitting or standing) for at least 30 minutes.
- If you have abnormal bone and mineral metabolism (for example lack of vitamin D, parathyroid hormone abnormalities, both leading to a low blood calcium level).
- If you have or have had problems in the past with your oesophagus (the tube that connects your mouth with your stomach). For instance you may have or have had pain or difficulty in swallowing food or you have previously been told that you have Barrett's oesophagus (a condition associated with changes in the cells that line the lower oesophagus).
- If you have been told by your doctor that you have an intolerance to some sugars (such as lactose).
- If you have or have had pain, swelling or numbness of the jaw or a “heavy jaw feeling” or loosening of a tooth.
- If you are under dental treatment or will undergo dental surgery, tell your dentist that you are being treated with Risedronate Pinewood.

Your doctor will advise you on what to do when taking Risedronate Pinewood if you have any of the above.

### **Children and adolescents**

Risedronate Pinewood is not recommended for use in children below 18 years of age due to insufficient data on safety and efficacy.

### **Other medicines and Risedronate Pinewood**

Medicines containing one of the following lessen the effect of Risedronate Pinewood if taken at the same time:

- calcium
- magnesium
- aluminium (for example some indigestion mixtures)
- iron

Take these medicines at least 30 minutes after taking Risedronate Pinewood.

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

### **Risedronate Pinewood with food and drink**

It is very important that you do NOT take your Risedronate Pinewood tablet with food or drinks (other than plain water) so that it can work properly. In particular do not take this medicine at the same time as dairy products (such as milk) as they contain calcium (see section 2, “Other medicines and Risedronate Pinewood”).

Take food and drinks (other than plain water) at least 30 minutes after your Risedronate Pinewood tablet.

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Do NOT take Risedronate Pinewood, if you may be pregnant, are pregnant or are planning to become pregnant (see section 2, “Do not take Risedronate Pinewood”). The potential risk associated with the use of risedronate sodium (active substance in Risedronate Pinewood) in pregnant women is unknown.

Do NOT take Risedronate Pinewood if you are breast-feeding (see section 2 “Do not take Risedronate Pinewood”).

Risedronate Pinewood should only be used to treat postmenopausal women and men.

### **Driving and using machines**

Risedronate Pinewood is not known to affect your ability to drive and use machines.

### **Risedronate Pinewood contains lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine (see section 2, “Warnings and precautions”).

### **Risedronate Pinewood contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially ‘sodium-free’.

## **3. How to take Risedronate Pinewood**

### **Dose**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

### Recommended dose

Take ONE Risedronate Pinewood tablet (35 mg of risedronate sodium) once a week.

Choose one day of the week that best fits your schedule. Every week, take the Risedronate Pinewood tablet on your chosen day.

There are boxes/spaces on the carton. Please mark the day of the week you have chosen to take your Risedronate Pinewood tablet. Also, write in the dates you will take the tablet.

### **When to take your Risedronate Pinewood tablet:**

Take your Risedronate Pinewood tablet at least 30 minutes before the first food, drink (other than plain water) or other medicine of the day.

### **How to take your Risedronate Pinewood tablet**

- Take the tablet whilst you are in an **upright position** (you may sit or stand) to avoid heartburn.
- **Swallow** it with at least one **glass** (120 mL) **of plain water**.
- **Swallow it whole**. Do not suck or chew it.
- **Do not lie down for 30 minutes after** taking the tablet.

Your doctor will tell you if you need calcium and vitamin supplements, if you are not taking enough from your diet.

### **If you take more Risedronate Pinewood than you should**

If you or somebody else has accidentally taken more Risedronate Pinewood tablets than prescribed, drink one full **glass of milk** and **seek medical attention**.

### **If you forget to take Risedronate Pinewood**

If you have forgotten to take your tablet at your regular time, you can take it at the next possible time according to the instruction above (i.e. before breakfast, between meals, or in the evening).

Return to taking one tablet once a week on the day the tablet is normally taken.

Do not take a double dose to make up for a forgotten tablet.

### **If you stop taking Risedronate Pinewood**

If you stop treatment, you may begin to lose bone mass. Please talk to your doctor before you consider stopping treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Stop taking Risedronate Pinewood and contact a doctor immediately** if you experience any of the following:

- Symptoms of a severe allergic reaction such as;
  - Swelling of the face, tongue or throat
  - Difficulties in swallowing
  - Hives and difficulties in breathing
- Severe skin reactions that can include blistering of the skin.

**Tell your doctor promptly**, if you experience the following side effects:

- Eye inflammation, usually with pain, redness and light sensitivity.
- Bone necrosis of the jaw (osteonecrosis) associated with delayed healing and infection, often following tooth extraction (see section 2, “Warnings and precautions”).
- Symptoms from oesophagus such as pain when you swallow, difficulties in swallowing, chest pain or new/worsened heartburn.

Unusual fracture of the thigh bone particularly in patients on long-term treatment for osteoporosis may occur rarely. Contact your doctor if you experience pain, weakness or discomfort in your thigh, hip or groin as this may be an early indication of a possible fracture of the thigh bone.

However in clinical studies the other side effects that were observed were usually mild and did not cause the patient to stop taking their tablets.

**Common side effects** (may affect up to 1 in 10 people):

- Indigestion, feeling sick, stomach ache, stomach cramps or discomfort, constipation, feelings of fullness, bloating, diarrhoea.
- Pain in your bones, muscles or joints.
- Headache.

**Uncommon side effects** (may affect up to 1 in 100 people):

- Inflammation or ulcer of the oesophagus (the tube that connects your mouth with your stomach) causing difficulty and pain in swallowing (see also section 2, “Warnings and precautions”), inflammation of the stomach and duodenum (bowel draining the stomach).
- Inflammation of the coloured part of the eye (iris) (red painful eyes with a possible change in vision).

**Rare side effects** (may affect up to 1 in 1,000 people):

- Inflammation of the tongue (red swollen, possibly painful), narrowing of the oesophagus (the tube that connects your mouth with your stomach).
- Abnormal liver tests have been reported. These can only be diagnosed from a blood test.

During post-marketing experience, the following have been reported

- Very rare side effects (may affect up to 1 in 10,000 people): Talk to your doctor if you have ear pain, discharge from the ear, and/or an ear infection. These could be signs of bone damage in the ear.
  -
- Not known (frequency cannot be estimated from the available data):

- A serious illness called Stevens-Johnson syndrome (SJS) characterized by blistering of the skin, mouth, eyes and other moist body surfaces (genitals), a severe illness called toxic epidermal necrolysis (TEN) which causes red rash over many parts of the body and/or loss of the outer layer of the skin, inflammation of small blood vessels characterized by palpable red spots on the skin (leukocytoclastic vasculitis).
- Hair loss.

Liver disorders, some cases were severe

Rarely, at the beginning of treatment, a patient's blood calcium and phosphate levels may fall.

These changes are usually small and cause no symptoms.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517; Website: [www.hpra.ie](http://www.hpra.ie); e-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Risedronate Pinewood**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after "EXP". The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Risedronate Pinewood contains**

- The active substance is risedronate sodium. Each film-coated tablet contains 35 mg risedronate sodium, equivalent to 32.5 mg risedronic acid.
- The other ingredients are:  
Tablet core: Starch, pregelatinised (maize); Cellulose, microcrystalline; Crospovidone; Magnesium stearate.  
Film coating: Hypromellose, Lactose monohydrate, Titanium dioxide (E171), Macrogol 4000.

### **What Risedronate Pinewood looks like and contents of the pack**

Risedronate Pinewood are white round biconvex film-coated tablets with diameter of 11.2 mm, 5.0 mm in thickness and embossed with "35" on one side. It is supplied in blister packs containing: 1, 2, 4, 10, 12, 16 film-coated tablets.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

Pinewood Laboratories Limited,  
Ballymacarbry,  
Clonmel,  
Co. Tipperary,  
Ireland.

**Manufacturer**

Batch releaser  
Pharmathen S.A.  
6, Dervenakion str.  
153 51 Pallini, Attiki  
Greece

**This medicinal product is authorised in the Member States of the EEA under the following names:**

DK: Karadronat 35 mg Filmovertrukne tabletter  
IE: Risedronate Pinewood Once a week 35 mg film-coated tablets  
IT: Risedronato Accord 35 mg compresse rivestite con film

**This leaflet was last revised in 09/2019**