

Package leaflet: Information for the user

Gabin 100 mg capsules, hard

Gabin 300 mg capsules, hard

Gabin 400 mg capsules, hard

gabapentin

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Gabin is and what it is used for
2. What you need to know before you take Gabin
3. How to take Gabin
4. Possible side effects
5. How to store Gabin
6. Contents of the pack and other information

1. What Gabin is and what it is used for

Gabin belongs to a group of medicines used to treat epilepsy and peripheral neuropathic pain (long lasting pain caused by damage to the nerves).

The active substance in Gabin is gabapentin.

Gabin is used to treat:

Various forms of epilepsy (seizures that are initially limited to certain parts of the brain, whether the seizure spreads to other parts of the brain or not).

The doctor treating you or your child 6 years of age and older will prescribe Gabin to help treat epilepsy when the current treatment is not fully controlling the condition. You or your child 6 years of age and older should take Gabin in addition to the current treatment unless told otherwise. Gabin can also be used on its own to treat adults and children over 12 years of age.

Peripheral neuropathic pain (long lasting pain caused by damage to the nerves). A variety of different diseases can cause peripheral neuropathic pain (primarily occurring in the legs and/or arms), such as diabetes or shingles. Pain sensations may be described as hot, burning, throbbing, shooting, stabbing, sharp, cramping, aching, tingling, numbness, pins and needles etc.

2. What you need to know before you take Gabin

Do not take Gabin

- if you are allergic to gabapentin or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor or pharmacist before taking Gabapentin

- if you suffer from kidney problems your doctor may prescribe a different dosing schedule

- if you are on haemodialysis (to remove waste products because of kidney failure), tell your doctor
- if you develop muscle pain and/or weakness
- if you develop signs such as persistent stomach pain, feeling sick and being sick contact your doctor immediately as these may be symptoms of acute pancreatitis (an inflamed pancreas).
- if you have nervous system disorders, respiratory disorders, or you are more than 65 years old, your doctor may prescribe you a different dosing regimen
- Before taking this medicine, tell your doctor if you have ever abused or been dependent on alcohol, prescription medicines or illegal drugs; it may mean you have a greater risk of becoming dependent on Gabin.

Cases of abuse and dependence have been reported for gabapentin from the post-marketing experience. Talk to your doctor if you have a history of abuse or dependence.

Dependence

Some people may become dependent on Gabin (a need to keep taking the medicine). They may have withdrawal effects when they stop using Gabin (see section 3, “How to take Gabin” and “If you stop taking Gabin”). If you have concerns that you may become dependent on Gabin, it is important that you consult your doctor.

If you notice any of the following signs whilst taking Gabin, it could be a sign that you have become dependent.

- You feel you need to take the medicine for longer than advised by your prescriber
- You feel you need to take more than the recommended dose
- You are using the medicine for reasons other than prescribed
- You have made repeated, unsuccessful attempts to quit or control the use of the medicine
- When you stop taking the medicine you feel unwell, and you feel better once taking the medicine again

If you notice any of these, speak to your doctor to discuss the best treatment pathway for you, including when it is appropriate to stop and how to do this safely.

A small number of people being treated with anti-epileptics such as gabapentin have had thoughts of harming or killing themselves. If at any time you have these thoughts, immediately contact your doctor.

Important information about potentially serious reactions

Serious skin rashes including Stevens-Johnson syndrome, toxic epidermal necrolysis and drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in association with gabapentin. Stop using gabapentin and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.

Read the description of these symptoms in section 4 of this leaflet under ‘Contact your doctor immediately if you experience any of the following symptoms after taking this medicine as they can be serious’

Muscle weakness, tenderness or pain and particularly, if at the same time, you feel unwell or have a high temperature it may be caused by an abnormal muscle breakdown which can be life-threatening and lead to kidney problems. You may also experience discoloration of your urine, and a change in blood test results (notably blood creatine phosphokinase increased). If you experience any of these signs or symptoms, please contact your doctor immediately.

Notice: The HDPE bottle contains desiccant. Do not swallow.

Other medicines and Gabin

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. In particular, tell your doctor or pharmacist if you are taking or have been recently taking

any medicines for convulsions, sleeping disorders, depression, anxiety, or any other neurological or psychiatric problems.

Medicines containing opioids such as morphine

If you are taking any medicines containing opioids (such as morphine), please tell your doctor or pharmacist as opioids may increase the effect of Gabin. In addition, combination of Gabapentin with opioids may cause sleepiness, sedation, decrease in breathing, or death.

Antacids for indigestion

If Gabin and antacids containing aluminium and magnesium are taken at the same time, absorption of Gabin from the stomach may be reduced. It is therefore recommended that Gabin is taken at the earliest two hours after taking an antacid.

Gabapentin:

- is not expected to interact with other antiepileptic drugs or the oral contraceptive pill.
- may interfere with some laboratory tests, if you require a urine test tell your doctor or hospital what you are taking.

Gabin with food, drink and alcohol

Gabin can be taken with or without food.

Pregnancy, breast-feeding and fertility

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy

Gabin should not be taken during pregnancy, unless you are told otherwise by your doctor. Effective contraception must be used by women of child-bearing potential.

There have been no studies specifically looking at the use of gabapentin in pregnant women, but other medications used to treat seizures have reported an increased risk of harm to the developing baby, particularly when more than one seizure medication is taken at the same time. Therefore, whenever possible you should try to take only one seizure medication during pregnancy and only under the advice of your doctor.

Contact your doctor immediately if you become pregnant, think you might be pregnant or are planning to become pregnant while taking Gabin. Do not suddenly discontinue taking this medicine as this may lead to a breakthrough seizure, which could have serious consequences for you and your baby.

If used during pregnancy, gabapentin may lead to withdrawal symptoms in newborn infants. This risk might be increased when gabapentin is taken together with opioid analgesics (drugs for treatment of severe pain).

Breast-feeding

Gabapentin, the active substance of Gabin, is passed on through human milk.

Because the effect on the baby is unknown, it is not recommended to breast-feed while using Gabin.

Fertility

There is no effect on fertility in animal studies.

Driving and using machines

Gabin may produce dizziness, drowsiness and tiredness. You should not drive, operate complex machinery or take part in other potentially hazardous activities until you know whether this medication affects your ability to perform these activities.

Gabin contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

3. How to take Gabin

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. Do not take more medicine than prescribed.

Your doctor will determine what dose is appropriate for you.

Epilepsy, the recommended dose is:

Adults and adolescents:

Take the number of capsules as instructed. Your doctor will usually build up your dose gradually. The starting dose will generally be between 300 mg and 900 mg each day. Thereafter, the dose may be increased as instructed by your doctor up to a maximum of 3600 mg each day and your doctor will tell you to take this in 3 separate doses, i.e. once in the morning, once in the afternoon and once in the evening.

Children aged 6 years and above:

The dose to be given to your child will be decided by your doctor as it is calculated against your child's weight. The treatment is started with a low initial dose which is gradually increased over a period of approximately 3 days. The usual dose to control epilepsy is 25-35 mg per kg per day. It is usually given in 3 separate doses, by taking the capsule(s) each day, usually once in the morning, once in the afternoon and once in the evening.

Gabin is not recommended for use in children below 6 years of age.

Peripheral Neuropathic Pain, the recommended dose is:

Adults

Take the number of capsules as instructed by your doctor. Your doctor will usually build up your dose gradually. The starting dose will generally be between 300 mg and 900 mg each day. Thereafter, the dose may be increased as instructed by your doctor up to a maximum of 3600 mg each day and your doctor will tell you to take this in 3 separate doses, i.e. once in the morning, once in the afternoon and once in the evening.

If you have kidney problems or are receiving haemodialysis

Your doctor may prescribe a different dosing schedule and/or dose if you have problems with your kidneys or are undergoing haemodialysis.

If you are an elderly patient (over 65 years of age)

You should take the normal dose of Gabapentin unless you have problems with your kidneys. Your doctor may prescribe a different dosing schedule and/or dose if you have problems with your kidneys.

If you have the impression that the effect of Gabapentin is too strong or too weak, talk to your doctor or pharmacist as soon as possible.

Method of Administration

Gabapentin is for oral use. Always swallow the capsules whole with plenty of water.

Continue taking Gabapentin until your doctor tells you to stop

If you take more Gabin than you should

Higher than recommended doses may result in an increase in side effects including loss of consciousness, dizziness, double vision, slurred speech, drowsiness and diarrhoea. Call your doctor or go to the nearest hospital emergency unit immediately if you take more Gabapentin than your doctor prescribed. Take along any capsules or tablets that you have not taken, together with the container and the label so that the hospital can easily tell what medicine you have taken.

If you forget to take Gabin

If you forget to take a dose, take it as soon as you remember unless it is time for your next dose. Do not take a double dose to make up for a forgotten dose.

If you stop taking Gabin

Do not suddenly stop taking Gabin. If you want to stop taking Gabin, discuss this with your doctor first. They will tell you how to do this.. If your treatment is stopped it should be done gradually over a minimum of 1 week. After stopping a short or long-term treatment with Gabin, you need to know that you may experience certain side effects, so-called withdrawal effects. These effects can include seizures, anxiety, difficulty sleeping, feeling sick (nausea), pain, sweating, shaking, headache, depression, feeling abnormal, dizziness, and feeling generally unwell. These effects usually occur within 48 hours after stopping Gabin. If you experience withdrawal effects, you should contact your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Stop using Gabin and seek medical attention immediately if you notice any of the following symptoms:

- reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (Stevens-Johnson- syndrome, toxic epidermal necrolysis).
- Widespread rash, high body temperature and enlarged lymph nodes (DRESS syndrome or drug hypersensitivity syndrome).

Contact your doctor immediately if you experience any of the following symptoms after taking this medicine as they can be serious:

- severe skin reactions that require immediate attention, swelling of the lips and face, skin rash and redness, and/or hair loss (these may be symptoms of a serious allergic reaction)
- persistent stomach pain, feeling sick and being sick as these may be symptoms of acute pancreatitis (an inflamed pancreas)
- breathing problems, which if severe you may need emergency and intensive care to continue breathing normally
- Gabin may cause a serious or life-threatening allergic reaction that may affect your skin or other parts of your body such as your liver or blood cells. You may or may not have rash when you get this type of reaction. It may cause you to be hospitalized or to stop Gabin. Call your doctor right away if you have any of the following symptoms:
 - skin rash
 - hives
 - fever
 - swollen glands that do not go away
 - swelling of your lip and tongue
 - yellowing of your skin or of the whites of the eyes
 - unusual bruising or bleeding

- **severe fatigue or weakness**
- **unexpected muscle pain**
- **frequent infections**

These symptoms may be the first signs of a serious reaction. A doctor should examine you to decide if you should continue taking Gabin.

- If you are on haemodialysis, tell your doctor if you develop muscle pain and/or weakness.

Other side effects include:

Very common (may affect more than 1 in 10 people):

- Viral infection
- Feeling drowsy, dizziness, lack of coordination
- Feeling tired, fever

Common (may affect up to 1 in 10 people):

- Pneumonia, respiratory infection, urinary tract infection, inflammation of the ear or other infections
- Low white blood cell counts
- Anorexia, increased appetite
- Anger towards others, confusion, mood changes, depression, anxiety, nervousness. , difficulty with thinking
- Convulsions, jerky movements, difficulty with speaking, loss of memory, tremor, difficulty sleeping, headache, sensitive skin, decreased sensation (numbness), difficulty with coordination, unusual eye movement, increased, decreased or absent reflexes
- Blurred vision, double vision
- Vertigo
- High blood pressure, flushing or dilation of blood vessels
- Difficulty breathing, bronchitis, sore throat, cough, dry nose
- Vomiting (being sick), nausea (feeling sick), problems with teeth, inflamed gums, diarrhoea, stomach pain, indigestion, constipation, dry mouth or throat, flatulence
- Facial swelling, bruises, rash, itch, acne
- Joint pain, muscle pain, back pain, twitching
- Difficulties with erection (impotence)
- Swelling in the legs and arms, difficulty with walking, weakness, pain, feeling unwell, flu-like symptoms
- Decrease in white blood cells, increase in weight
- Accidental injury, fracture, abrasion

Additionally in clinical studies in children, aggressive behaviour and jerky movements were reported commonly.

Uncommon (may affect up to 1 in 100 people):

- Agitation (a state of chronic restlessness and unintentional and purposeless motions)
- Allergic reaction such as hives
- Decreased movement
- Racing heartbeat
- Difficulty swallowing
- Swelling that may involve the face, trunk and limbs
- Abnormal blood test results suggesting problems with the liver.
- Mental impairment
- Fall
- Increase in blood glucose levels (most often observed in patients with diabetes)

Rare: (may affect up to 1 in 1,000 people)

- Decrease in blood glucose levels (most often observed in patients with diabetes)
- Loss of consciousness
- Trouble breathing, shallow breaths (respiratory depression)

After marketing Gabapentin the following side effects have been reported:

Not known: (frequency cannot be estimated from the available data)

- Decreased platelets (blood clotting cells)
- Suicidal thoughts, hallucinations
- Problems with abnormal movements such as writhing, jerking movements and stiffness
- Ringing in the ears
- Yellowing of the skin and eyes (jaundice)
- Inflammation of the liver
- Acute kidney failure, incontinence
- Increased breast tissue, breast enlargement
- Adverse events following the abrupt discontinuation of gabapentin (anxiety, difficulty sleeping, feeling sick, pain, sweating), chest pain
- Breakdown of muscle fibers (rhabdomyolysis)
- Change in blood test results (creatine phosphokinase increased)
- Problems with sexual functioning including inability to achieve a sexual climax, delayed ejaculation
- Low blood sodium level
- Anaphylaxis (serious, potentially life threatening allergic reaction including difficulty breathing, swelling of the lips, throat, and tongue, and hypotension requiring emergency treatment)
- Becoming dependent on Gabin ('drug dependence')

After stopping a short or long-term treatment with Gabin, you need to know that you may experience certain side effects, so-called withdrawal effects (see "If you stop taking Gabin").

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance. Website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Gabin

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date, which is stated on the carton/label/bottle after EXP. The expiry date refers to the last date of that month.

Store below 25°C.

Blister pack: Store in the original package.

HDPE bottle pack: Store in the original container.

Use within 12 months after first opening the HDPE bottle.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Gabin contains

- The active substance is gabapentin.

Each 100 mg hard capsule contains 100 mg gabapentin.

Each 300 mg hard capsule contains 300 mg gabapentin.

Each 400 mg hard capsule contains 400 mg gabapentin.

- The other ingredients are:

Capsule contents: Maize starch and talc.

Capsule shell:

100 mg: Titanium dioxide (E171), Sodium lauril sulfate, gelatin.

300 mg: Yellow iron oxide (E172), Titanium dioxide (E171), Sodium lauril sulfate, gelatin.

400 mg: Red iron oxide (E172), Yellow iron oxide (E172), Titanium dioxide (E171), Sodium lauril sulfate, gelatin.

Printing ink: Shellac, propylene glycol, black iron oxide, potassium hydroxide

What Gabin looks like and contents of the pack

Capsule, hard.

Gabin 100 mg capsules imprinted with 'D' on white cap and '02' on white body, containing white to off-white crystalline powder.

Gabin 300 mg capsules imprinted with 'D' on yellow cap and '03' on yellow body, containing white to off-white crystalline powder.

Gabin 400 mg capsules imprinted with 'D' on orange cap and '04' on orange body, containing white to off-white crystalline powder.

Gabin 100 mg capsules are available in:

- Clear PVC/PVdC – Aluminium foil blister packs of 10, 20, 30, 50, 60, 90, 100 and 200 capsules, hard.
- HDPE bottle with polypropylene closure containing silica gel desiccant: 100, 200 and 1000 capsules, hard.

Gabin 300 mg capsules are available in:

- Clear PVC/PVdC – Aluminium foil blister packs of 10, 20, 30, 50, 60, 90, 100 and 200 capsules, hard.
- HDPE bottle with polypropylene closure containing silica gel desiccant: 100, 200 and 1000 capsules, hard.

Gabin 400 mg capsules are available in:

- Clear PVC/PVdC – Aluminium foil blister packs of 10, 20, 30, 50, 60, 90, 100, 200 and 300 capsules, hard.
- HDPE bottle with polypropylene closure containing silica gel desiccant: 100, 200, 300 and 500 capsules, hard.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Aurobindo Pharma (Malta) Limited
Vault 14, Level 2, Valletta Waterfront
Floriana FRN 1913
Malta

Manufacturer

APL Swift Services (Malta) Limited
HF26, Hal Far Industrial Estate, Hal Far

Birzebbugia, BBG 3000
Malta

or

Generis Farmacêutica S.A.,
Rua João de Deus, nº 19,
Venda Nova, 2700-487
Amadora, Portugal

This medicinal product is authorised in the Member States of the EEA under the following names:

Czech Republic	Gabapentin Aurovitas 100 mg/ 300 mg/ 400mg tvrdé tobolky
France	Gabapentine Arrow Génériques 100 mg/ 300 mg/ 400mg, gélules
Germany	Gabapentin Aurobindo 100 mg/ 300 mg/ 400mg Hartkapseln
Ireland	Gabin 100 mg/ 300 mg/ 400mg capsules, hard
Italy	Gabapentin Aurobindo 100 mg/ 300 mg/ 400mg capsule rigide
Latvia	Gabapentin Aurobindo 300 mg/ 400mg cietās kapsulas
Netherlands	Gabapentine Aurobindo 100 mg/ 300 mg/ 400mg, capsules, hard
Portugal	Gabapentina Aurobindo
Romania	Gabapentin Aurobindo 100 mg/ 300 mg/ 400mg capsule
Slovak Republic	Gabapentin Vipfarm 100 mg/ 300 mg/ 400mg tvrdé kapsuly
Spain	Gabapentina Aurobindo 100 mg/ 400mg cápsulas duras
United Kingdom	Gabapentin Milpharm 100 mg/ 300 mg/ 400mg capsules, hard

This leaflet was last revised in 01/2023