

Package leaflet: Information for the user

Zineryt® 40 mg + 12 mg powder and solvent for cutaneous solution erythromycin/zinc acetate dihydrate

Read all of this leaflet carefully before you start using this medicine because it contains important information for you

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

In this leaflet:

1. What Zineryt® is and what it is used for
2. What you need to know before you use Zineryt®
3. How to use Zineryt®
4. Possible side effects
5. How to store Zineryt®
6. Contents of the pack and other information
7. Information for healthcare professionals

1. What Zineryt® is and what it is used for

The active substance of Zineryt belongs to the group of antibiotics for treatment of acne, often referred to as pimples or spots. These medicines are used to reduce the growth of bacteria which cause acne.

2. What you need to know before you use Zineryt®

Do not use Zineryt

- if you are allergic (hypersensitive) to erythromycin, or zinc or to any of the other ingredients of Zineryt (see list in section 6 'Contents of the pack and other information'). Your doctor may decide to give you another medicine instead.
- if you are allergic to antibiotics of the macrolide group. Ask your doctor or pharmacist if you are unsure.

Warnings and precautions

- Avoid contact with the eyes and mucous membranes of the nose and mouth. However, if this occurs, then rinse the affected area immediately with cold or lukewarm water for several minutes. Contact your doctor for further advice.

Other medicines and Zineryt

Please tell your doctor or pharmacist if you are taking or have recently taken, used or might use any other medicines, including medicines obtained without a prescription.

Pregnancy, breast-feeding and fertility

Ask your doctor or pharmacist for advice before taking any medicine.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

This product should **not** be used in pregnancy unless your doctor advises otherwise.

Zineryt should be used with caution in breast-feeding patients but do not use this medicine on the area of your chest.

Driving and using machines

It is not likely that Zineryt affects your ability to drive or use machines.

Zineryt contains Ethanol

This medicine contains alcohol (ethanol). It may cause burning sensation on damaged skin.

3. How to use Zineryt®


Instructions for proper use

Wash the affected area first before using Zineryt.

If you want to apply make-up, use the medicine first and let it dry.

Zineryt comes to you in a bottle with a special applicator pad protected by a cap which you should remove before use and replace firmly after each use. This pad makes it easy to apply the solution to the whole of the affected area of the skin to be treated.

To use Zineryt, unscrew the protective cap, hold the bottle upside down, place the pad against the skin, and spread the solution over the whole of the affected area (including the surrounding skin). The flow of solution can be increased by pressing the pad a little more firmly against the skin.



If you use the applicator properly, just the right quantity should be applied. But if you do put too much on, blot it off with some toilet tissue, and make sure it does not run into the eyes, nose or mouth. If it does, wash well with plenty of water (see also Warnings and precautions under section 2 'What you need to know before you use Zineryt').

Always use Zineryt exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Frequency of administration

Usually, Zineryt should be applied all over the affected area and the surrounding skin twice a day.

Duration of treatment

Successful treatment of acne may take several weeks or even months. The normal treatment time is 10 to 12 weeks.

Consult your doctor if you still have acne at the end of the prescribed course of treatment.

If you use more Zineryt than you should

If you have applied too much Zineryt, blot it off with some toilet tissue. If you begin to show signs of hypersensitivity (allergic reaction) you should wash the treated area with copious amount of water and simple soap.

Swallowing the entire contents of a single pack of Zineryt would mainly lead to symptoms associated with alcohol intake.

You should contact your doctor or pharmacist if you or somebody else accidentally swallows this medicine.

If you forget to use Zineryt

If you have missed an application, use Zineryt when you remember, but leave about three hours before you use it again.

If you stop using Zineryt

If you stop using Zineryt, the original symptoms may come back. Always contact your doctor if you want to stop using Zineryt.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, Zineryt can cause side effects, although not everybody gets them.

Contact a doctor immediately if you experience a serious skin reaction: a red, scaly rash with bumps under the skin and blisters (exanthematous pustulosis). The frequency of this side effect is not known (cannot be estimated from the available data).

Zineryt may very rarely cause allergic reactions. Symptoms may include rash, itching, swelling of the mouth, face and lips or tongue and difficulty breathing.

Please contact your doctor immediately if you experience any of these side effects.

Zineryt may cause the following uncommon side effects (likely to affect more than 1 in 1000 and less than 1 in 100 patients):

- Temporary stinging or slight redness of the skin
- Dry skin
- Skin exfoliation
- Itching
- Skin irritation

If you get a very painful rash or burning feeling after using Zineryt which does not go away in a few hours, or gets worse each time you use it, stop using it and tell your doctor as soon as possible.

Long-term use of an antibiotic such as Zineryt may result in an infection due to the development of resistance to the antibiotic product.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Zineryt®

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the label and carton after EXP. The expiry date refers to the last day of that month.

Do not store above 25°C. Store in the original package.

Do not use this medicine after the "Use before" date which has been put on the label by your pharmacist. The "Use before" date is 5 weeks after the pharmacist has prepared this medicine into a solution.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you

no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Zineryt contains

- The active substances are erythromycin (40 mg/ml) and zinc acetate dihydrate (12 mg/ml).
- The other ingredients are di-isopropyl sebacate and anhydrous ethanol.

What Zineryt looks like and contents of the pack

Zineryt is made available as a cutaneous solution. The packaging contains a bottle of white crystalline powder, a bottle of clear colourless solvent and an applicator in a plastic holder. The contents of the bottles will be mixed together by your pharmacist to form a solution. Your pharmacist will also fit the special applicator top to the bottle containing the mixed solution. The applicator top is protected by a cap. Zineryt is available in packs which, when made up, will contain 30 ml of the solution.

Parallel Product Authorisation Holder: IMED Healthcare Ltd, Unit 625 Kilshane Avenue, Northwest Business Park, Ballycoolin, Dublin 15, Ireland

Manufacturer: ASTELLAS PHARMA EUROPE B.V. Hogemaat 2, 7942 JG Meppel, Holland and LEO Pharma A/S, Industriparken 55, DK-2750 Ballerup, Denmark

Repackaged by: Cast Healthcare Ltd, Unit E, The Business Centre, 5-7 Tobermore Road, Draperstown, Magherafelt, BT45 7AG, UK(NI) or IMED Healthcare Ltd, Unit 625 Kilshane Avenue, Northwest Business Park, Ballycoolin, Dublin 15, Ireland

This leaflet was last revised in May 2023.

Information for healthcare professionals

The following information is intended for medical or healthcare professionals only:

Instructions for use and handling

The powder is reconstituted with the solvent prior to dispensing as follows:

1. Remove the caps from the powder bottle and the solution bottle; retain the cap of the powder bottle.
2. Pour the contents of the solution bottle into the powder bottle and recap the latter.
3. Immediately, shake well for one minute. Remove and retain cap.
4. Open the plastic holder containing the applicator assembly and use the holder to position the applicator assembly over the neck of the bottle and to push the applicator assembly firmly into the neck of the bottle.
5. Remove the plastic holder and ensure that the applicator fits firmly into the neck of the bottle. Discard the plastic holder.
6. Replace the cap on the now constituted application.
7. Add the expiry date to the bottle label; this date is 5 weeks from the date of preparation.
8. Ensure that the patient information leaflet is given to the patient.

After reconstitution, a clear to hazy liquid is formed.