


ZINC Ref: <b>EAS1555g LEA EZETIMIBE/SIMVASTATIN TEVA PHARMA 10/20MG 10/40MG 10/80MG IE BALK</b>	Version: <b>2</b> 23 February 2021		
	Length: <b>600 mm</b> Width: <b>190 mm</b> Depth: <b>-</b>		
Colours Used:	<table border="1"> <tr> <td>BLACK</td> </tr> <tr> <td>Template</td> </tr> </table>	BLACK	Template
BLACK			
Template			

**Ezetimibe/Simvastatin Teva Pharma 10 mg/20 mg Tablets**  
**Ezetimibe/Simvastatin Teva Pharma 10 mg/40 mg Tablets**  
**Ezetimibe/Simvastatin Teva Pharma 10 mg/80 mg Tablets**  
ezetimibe/simvastatin

**Package leaflet: Information for the patient**

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Ezetimibe/Simvastatin Teva Pharma is and what it is used for
2. What you need to know before you take Ezetimibe/Simvastatin Teva Pharma
3. How to take Ezetimibe/Simvastatin Teva Pharma
4. Possible side effects
5. How to store Ezetimibe/Simvastatin Teva Pharma
6. Contents of the pack and other information

**1 What Ezetimibe/Simvastatin Teva Pharma is and what it is used for**

Ezetimibe/Simvastatin Teva Pharma contains the active substances ezetimibe and simvastatin. Ezetimibe/Simvastatin Teva Pharma is a medicine used to lower levels of total cholesterol, "bad" cholesterol (LDL cholesterol), and fatty substances called triglycerides in the blood. In addition, Ezetimibe/Simvastatin Teva Pharma raises levels of "good" cholesterol (HDL cholesterol).

Ezetimibe/Simvastatin Teva Pharma works to reduce your cholesterol in two ways. The active ingredient ezetimibe reduces the cholesterol absorbed in your digestive tract. The active ingredient simvastatin belonging to the class of "statins" inhibits the production of the cholesterol your body makes by itself.

Cholesterol is one of several fatty substances found in the bloodstream. Your total cholesterol is made up mainly of LDL and HDL cholesterol.

LDL cholesterol is often called "bad" cholesterol because it can build up in the walls of your arteries forming plaque. Eventually this plaque build-up can lead to a narrowing of the arteries. This narrowing can slow or block blood flow to vital organs such as the heart and brain. This blocking of blood flow can result in a heart attack or stroke.

HDL cholesterol is often called "good" cholesterol because it helps keep the bad cholesterol from building up in the arteries and protects against heart disease.

Triglycerides are another form of fat in your blood that may increase your risk for heart disease.

Ezetimibe/Simvastatin Teva Pharma is used for patients who cannot control their cholesterol levels by diet alone. You should stay on a cholesterol-lowering diet while taking this medicine.

Ezetimibe/Simvastatin Teva Pharma is used in addition to your cholesterol-lowering diet if you have:

- a raised cholesterol level in your blood (primary hypercholesterolaemia (heterozygous familial and non-familial)) or elevated fat levels in your blood (mixed hyperlipidaemia):
  - that is not well controlled with a statin alone,
  - for which you have used a statin and ezetimibe as separate tablets.
- a hereditary illness (homozygous familial hypercholesterolaemia) that increases the cholesterol level in your blood. You may also receive other treatments.
- heart disease, Ezetimibe/Simvastatin Teva Pharma reduces the risk of heart attack, stroke, surgery to increase heart blood flow, or hospitalization for chest pain.

Ezetimibe/Simvastatin Teva Pharma does not help you lose weight.

**2 What you need to know before you take Ezetimibe/Simvastatin Teva Pharma**

**Do not take Ezetimibe/Simvastatin Teva Pharma if**

- you are allergic to ezetimibe, simvastatin, or any of the other ingredients of this medicine (listed in section 6).

- you currently have liver problems.
- you are pregnant or breast-feeding.
- you are taking medicine(s) with one or more than one of the following active ingredients:
  - itraconazole, ketoconazole, posaconazole or voriconazole (used to treat fungal infections).
  - erythromycin, clarithromycin, or telithromycin (used to treat infections).
  - HIV protease inhibitors such as indinavir, nelfinavir, ritonavir, and saquinavir (HIV protease inhibitors are used to treat HIV infections).
  - boceprevir or telaprevir (used to treat hepatitis C virus infections).
  - nefazodone (used to treat depression).
  - cobicistat.
  - gemfibrozil (used to lower cholesterol).
  - ciclosporin (often used in organ transplant patients).
  - danazol (a man-made hormone used to treat endometriosis, a condition in which the lining of the uterus grows outside the uterus).
- you are taking or have taken in the last 7 days a medicine called fusidic acid (a medicine for bacterial infection) orally or by injection. The combination of fusidic acid and Ezetimibe/Simvastatin Teva Pharma can lead to serious muscle problems (rhabdomyolysis).

Do not take more than 10/40 mg Ezetimibe/Simvastatin Teva Pharma if you are taking lomitapide (used to treat a serious and rare genetic cholesterol condition).

Ask your doctor if you are not sure if your medicine is listed above.

**Warnings and precautions**

- Talk to your doctor or pharmacist before taking Ezetimibe/Simvastatin Teva Pharma
- about all your medical conditions including allergies.
  - if you drink large amounts of alcohol or have ever had liver disease. Ezetimibe/Simvastatin Teva Pharma may not be right for you.
  - if you are due to have an operation. You may need to stop taking Ezetimibe/Simvastatin Teva Pharma tablets for a short time.
  - if you are Asian, because a different dose may be applicable to you.

Your doctor should do a blood test before you start taking Ezetimibe/Simvastatin Teva Pharma and if you have any symptoms of liver problems while you take Ezetimibe/Simvastatin Teva Pharma. This is to check how well your liver is working.

Your doctor may also want you to have blood tests to check how well your liver is working after you start taking Ezetimibe/Simvastatin Teva Pharma.

While you are on this medicine your doctor will monitor you closely if you have diabetes or are at risk of developing diabetes. You are likely to be at risk of developing diabetes if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure.

Tell your doctor if you have severe lung disease.

The combined use of Ezetimibe/Simvastatin Teva Pharma and fibrates (certain medicines for lowering cholesterol) should be avoided since the combined use of Ezetimibe/Simvastatin Teva Pharma and fibrates has not been studied.

**Contact your doctor immediately if you experience unexplained muscle pain, tenderness, or weakness. This is because on rare occasions, muscle problems can be serious, including muscle breakdown resulting in kidney damage; and very rare deaths have occurred.**

The risk of muscle breakdown is greater at higher doses of Ezetimibe/Simvastatin Teva Pharma, particularly the 10 mg/80 mg dose. The risk of muscle breakdown is also greater in certain patients. Talk with your doctor if any of the following applies:

- you have kidney problems
- you have thyroid problems
- you are 65 years or older
- you are female
- you have ever had muscle problems during treatment with cholesterol lowering medicines called "statins" (like simvastatin, atorvastatin, and rosuvastatin) or fibrates (like gemfibrozil and bezafibrate)
- you or close family members have a hereditary muscle disorder

Also tell your doctor or pharmacist if you have a muscle weakness that is constant. Additional tests and medicines may be needed to diagnose and treat this.

**Children and adolescents**

Ezetimibe/Simvastatin Teva Pharma is not recommended for children under age 10.

**Other medicines and Ezetimibe/Simvastatin Teva Pharma**

Tell your doctor if you are taking, have recently taken or might take any other medicine(s) with any of the following active ingredients. Taking Ezetimibe/Simvastatin Teva Pharma with any of the following medicines can increase the risk of muscle problems (some of these have already been listed in the above section "Do not take Ezetimibe/Simvastatin Teva Pharma if").

- **If you need to take oral fusidic acid to treat a bacterial infection you will need to temporarily stop using this medicine. Your doctor will tell you when it is safe to restart Ezetimibe/Simvastatin Teva Pharma. Taking Ezetimibe/Simvastatin Teva Pharma with fusidic acid may rarely lead to muscle weakness, tenderness or pain (rhabdomyolysis). See more information regarding rhabdomyolysis in section 4.**
- ciclosporin (often used in organ transplant patients)
- danazol (a man-made hormone used to treat endometriosis, a condition in which the lining of the uterus grows outside the uterus)
- medicines with an active ingredient like itraconazole, ketoconazole, fluconazole, posaconazole, or voriconazole (used to treat fungal infections)
- fibrates with active ingredients like gemfibrozil and bezafibrate (used to lower cholesterol)
- erythromycin, clarithromycin or telithromycin (used to treat bacterial infections)
- HIV protease inhibitors such as indinavir, nelfinavir, ritonavir, and saquinavir (used to treat AIDS)
- Hepatitis C antiviral agents such as boceprevir, telaprevir, elbasvir or grazoprevir (used to treat hepatitis C virus infections)
- nefazodone (used to treat depression)
- medicines with the active ingredient cobicistat
- amiodarone (used to treat an irregular heartbeat)
- verapamil, diltiazem, or amlodipine (used to treat high blood pressure, chest pain associated with heart disease, or other heart conditions)
- lomitapide (used to treat a serious and rare genetic cholesterol condition)
- daptomycin (a drug used to treat complicated skin and skin structure infections and bacteraemia). It is possible that side effects affecting the muscles may be higher when this medicine is taken during treatment with simvastatin (e.g. Ezetimibe/Simvastatin Teva Pharma). Your doctor may decide that you stop taking Ezetimibe/Simvastatin Teva Pharma for a while.
- large amounts (1 gram or more each day) of niacin or nicotinic acid (also used to lower cholesterol)
- colchicine (used to treat gout).

As well as the medicines listed above, tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. In particular, tell your doctor if you are taking any of the following:

- medicines with an active ingredient to prevent blood clots, such as warfarin, flunindione, phenprocoumon or acenocoumarol (anticoagulants)
- cholestyramine (also used to lower cholesterol), because it affects the way Ezetimibe/Simvastatin Teva Pharma works
- fenofibrate (also used to lower cholesterol)
- rifampicin (used to treat tuberculosis)
- ticagrelor (antiplatelet medicine).

You should also tell any doctor who is prescribing a new medicine for you that you are taking Ezetimibe/Simvastatin Teva Pharma.

**Ezetimibe/Simvastatin Teva Pharma with food and drink**  
Grapefruit juice contains one or more components that alter the metabolism of some medications, including Ezetimibe/Simvastatin Teva Pharma. Consuming grapefruit juice should be avoided as it may increase your risk of muscle problems.

**Pregnancy and breast-feeding**

Do not take Ezetimibe/Simvastatin Teva Pharma if you are pregnant, are trying to get pregnant or think you may be pregnant. If you get pregnant while taking Ezetimibe/Simvastatin Teva Pharma, stop taking it immediately and tell your doctor. Do not take Ezetimibe/Simvastatin Teva Pharma if you are breast-feeding, because it is not known if the medicine is passed into breast milk.

Ask your doctor or pharmacist for advice before taking any medicine.

**Driving and using machines**

Ezetimibe/Simvastatin Teva Pharma is not expected to interfere with your ability to drive or to use machinery. However, it should be taken into account that some people get dizzy after taking Ezetimibe/Simvastatin Teva Pharma.

**Ezetimibe/Simvastatin Teva Pharma contains lactose**  
If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

**Ezetimibe/Simvastatin Teva Pharma contains sodium**  
This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

**3 How to take Ezetimibe/Simvastatin Teva Pharma**

Your doctor will determine the appropriate tablet strength for you, depending on your current treatment and your personal risk status.

The tablets are not scored and should not be divided.

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

- Before starting Ezetimibe/Simvastatin Teva Pharma, you should be on a diet to lower your cholesterol.
- You should keep on this cholesterol-lowering diet while taking Ezetimibe/Simvastatin Teva Pharma.

Adults: The dose is 1 **tablet** by mouth once a day. Adolescents (10 to 17 years of age): The dose is 1 **tablet** by mouth once a day (a maximum dose of 10 mg/40 mg once daily must not be exceeded).

The Ezetimibe/Simvastatin Teva Pharma 10 mg/80 mg dose is only recommended for adult patients with very high cholesterol levels and at high risk of heart disease problems who have not reached their cholesterol goal on lower doses.

Take Ezetimibe/Simvastatin Teva Pharma in the evening. You can take it with or without food.

If your doctor has prescribed Ezetimibe/Simvastatin Teva Pharma along with another medicine for lowering cholesterol containing the active ingredient cholestyramine or any other bile acid sequestrant, you should take Ezetimibe/Simvastatin Teva Pharma at least 2 hours before or 4 hours after taking the bile acid sequestrant.

**If you take more Ezetimibe/Simvastatin Teva Pharma than you should**  
Please contact your doctor or pharmacist.

**If you forget to take Ezetimibe/Simvastatin Teva Pharma**  
Do not take a double dose to make up for a forgotten tablet, just take your normal amount of Ezetimibe/Simvastatin Teva Pharma at the usual time the next day.

**If you stop taking Ezetimibe/Simvastatin Teva Pharma**  
Talk to your doctor or pharmacist because your cholesterol may rise again.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4 Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them (see section 2 "What you need to know before you take Ezetimibe/Simvastatin Teva Pharma").

The following common side effects were reported (may affect up to 1 in 10 people):

- muscle aches
- elevations in laboratory blood tests of liver (transaminases) and/or muscle (CK) function

The following uncommon side effects were reported (may affect up to 1 in 100 people):

- elevations in blood uric acid; elevations in the time it takes for blood to clot; protein in urine; weight decreased
- dizziness; headache; tingling sensation
- abdominal pain; indigestion; flatulence; nausea; vomiting; abdominal bloating; diarrhoea; dry mouth; heartburn
- rash; itching; hives
- joint pain; muscle pain, tenderness, weakness or spasms; neck pain; pain in arms and legs; back pain
- unusual tiredness or weakness; feeling tired; chest pain; swelling, especially in the hands and feet
- sleep disorder; trouble sleeping

Additionally, the following side effects have been reported in people taking either combination of ezetimibe and simvastatin or medicines containing the active ingredients ezetimibe or simvastatin:

- low red blood cell count (anaemia); reduction in blood cell counts, which may cause bruising/bleeding (thrombocytopenia)
- numbness or weakness of the arms and legs; poor memory, memory loss, confusion
- breathing problems including persistent cough and/or shortness of breath or fever
- constipation
- inflammation of the pancreas often with severe abdominal pain
- inflammation of the liver with the following symptoms: yellowing of the skin and eyes, itching, dark coloured urine or pale coloured stool, feeling tired or weak, loss of appetite; liver failure; gallstones or inflammation of the gallbladder (which may cause abdominal pain, nausea, vomiting)
- hair loss; raised red rash, sometimes with target-shaped lesions (erythema multiforme)
- blurred vision and impaired vision (which each may affect up to 1 in 1,000 people)

- rash that may occur on the skin or sores in the mouth (lichenoid drug eruptions) (which each may affect up to 1 in 10,000 people)
- hypersensitivity reactions including some of the following: allergic reactions including swelling of the face, lips, tongue and/or throat which may cause difficulty in breathing or swallowing and requires treatment immediately (angioedema), pain or inflammation of the joints, inflammation of blood vessels, unusual bruising, skin eruptions and swelling, hives, skin sensitivity to the sun, fever, flushing, shortness of breath and feeling unwell, lupus-like disease picture (including rash, joint disorders, and effects on white blood cells). A serious very rare allergic reaction (which may affect up to 1 in 10,000 people) may occur that causes difficulty in breathing or dizziness and requires immediate treatment (anaphylaxis).
- muscle pain, tenderness, weakness or cramps; muscle breakdown; muscle rupture (which may affect up to 1 in 10,000 people); tendon problems, sometimes complicated by rupture of the tendon.
- gynecomastia (breast enlargement in men) (which may affect up to 1 in 10,000 people)
- decreased appetite
- hot flush; high blood pressure
- pain
- erectile dysfunction
- depression
- alterations in some laboratory blood tests for liver function

Additional possible side effects reported with some statins:

- sleep disturbances, including nightmares
- sexual difficulties
- diabetes. This is more likely if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure. Your doctor will monitor you while you are taking this medicine.
- muscle pain, tenderness, or weakness that is constant that may not go away after stopping Ezetimibe/Simvastatin Teva Pharma (frequency not known: cannot be estimated from the available data).

**Contact your doctor immediately if you experience unexplained muscle pain, tenderness, or weakness. This is because on rare occasions, muscle problems can be serious, including muscle breakdown resulting in kidney damage; and very rare deaths have occurred.**

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRa Pharmacovigilance Website: [www.hpra.ie](http://www.hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

**5 How to store Ezetimibe/Simvastatin Teva Pharma**

Keep this medicine out of the sight and reach of children.

Do not use Ezetimibe/Simvastatin Teva Pharma after the expiry date which is stated on the carton or container after EXP. The expiry date refers to the last day of that month.

Do not store above 25°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

**6 Contents of the pack and other information**

**What Ezetimibe/Simvastatin Teva Pharma contains**  
The active substances are ezetimibe and simvastatin. Each tablet contains 10 mg ezetimibe and 20 mg, 40 mg or 80 mg simvastatin.

The other ingredients are: *Tablet core:* lactose monohydrate, hypromellose, croscarmellose sodium, microcrystalline cellulose, ascorbic acid, citric acid, butylated hydroxyanisole, propyl gallate, magnesium stearate. *Pigment blend:* PB-220001 Yellow containing

lactose monohydrate, iron oxide yellow, iron oxide red and iron oxide black.

**What Ezetimibe/Simvastatin Teva Pharma looks like and contents of the pack**  
Ezetimibe/Simvastatin Teva Pharma 10 mg/20 mg tablets are light tan, mottled, round, 8 mm in diameter, biconvex tablets, plain on one side and marked with "512" on one side.  
Ezetimibe/Simvastatin Teva Pharma 10 mg/40 mg tablets are light tan, mottled, round, 10 mm in diameter, biconvex tablets, plain on one side and marked with "513" on one side.  
Ezetimibe/Simvastatin Teva Pharma 10 mg/80 mg tablets are light tan, mottled, capsule shaped, 17.5 x 7.55 mm, biconvex tablets, plain on one side and marked with "515" on one side.

**Pack sizes:**  
Blister packs:  
Ezetimibe/Simvastatin Teva Pharma 10 mg/80 mg tablets: 30 and 100 tablets  
Ezetimibe/Simvastatin Teva Pharma 10 mg/20 mg tablets and Ezetimibe/Simvastatin Teva Pharma 10 mg/40 mg tablets: 30, 98 and 100 tablets

HDPE tablet container: 100 and 180 tablets  
The HDPE-bottles contain a container with a silica gel desiccant. This container must not be removed from the bottle.

Not all packs may be marketed.

**Marketing Authorisation Holder and Manufacturer**

**Marketing Authorisation Holder**  
Teva B.V.  
Swensweg 5  
2031GA Haarlem  
Netherlands

**Manufacturer**  
Balkanpharma-Dupnitsa AD  
3 Samokovsko Shosse Str.  
Dupnitsa 2600,  
Bulgaria

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Austria	Ezetimib/Simvastatin Actavis 10 mg/20 mg, 10 mg/40 mg & 10 mg/80 mg Tabletten
Iceland	Ezetimib/Simvastatin Actavis
Ireland	Ezetimibe/Simvastatin Teva Pharma 10 mg/20 mg, 10 mg/40 mg & 10 mg/80 mg Tablets
Slovakia	Ezetimibe/Simvastatin Teva B.V.

**This leaflet was last revised in February 2021.**

