

Package leaflet: information for the user

Fampridine Rowa 10mg prolonged-release tablets

fampridine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Fampridine Rowa is and what it is used for
2. What you need to know before you take Fampridine Rowa
3. How to take Fampridine Rowa
4. Possible side effects
5. How to store Fampridine Rowa
6. Contents of the pack and other information

1. What Fampridine Rowa is and what it is used for

Fampridine Rowa contains the active substance fampridine which belongs to a group of medicines called potassium channel blockers. They work by stopping potassium leaving the nerve cells which have been damaged by MS. This medicine is thought to work by letting signals pass down the nerve more normally, which allows you to walk better.

Fampridine Rowa is a medicine used to improve walking in adults (18 years and over) with Multiple Sclerosis (MS) related walking disability. In multiple sclerosis, inflammation destroys the protective sheath around the nerves leading to muscle weakness, muscle stiffness and difficulty walking.

2. What you need to know before you take Fampridine Rowa

Do not take Fampridine Rowa

- if you are **allergic** to fampridine or any of the other ingredients of this medicine (listed in section 6)
- if you have a seizure or have ever had a **seizure** (also referred to as a fit or convulsion)
- if you have **kidney problems**
- if you are taking a medicine called cimetidine
- if you are **taking any other medicine containing fampridine**. This may increase your risk of serious side effects

Tell your doctor and do not take Fampridine Rowa if any of these apply to you.

Warnings and precautions

Talk to your doctor or pharmacist before taking Fampridine Rowa

- if you feel aware of your heartbeat (*palpitations*)
- if you are prone to infections
- if you have any factors or are taking any medicine which affects your risk of fits (*seizure*).
- if you have been told by a doctor that you have mild problems with your kidneys
- if you have a history of allergic reactions

You should use a walking aid, such as a cane, as needed, because this medicine may make you feel dizzy or unsteady this may result in an increased risk of falls.

Tell your doctor before you take Fampridine Rowa if any of these apply to you.

Children and in adolescents

Do not give Fampridine Rowa to children or adolescents under the age of 18 years.

Older people

Before starting treatment and during treatment your doctor may check that your kidneys are working properly.

Other medicines and Fampridine Rowa

Tell your doctor or pharmacist if you are taking, have recently taken or might take **any other medicines**.

Do not take Fampridine Rowa if you are taking any other medicine containing fampridine.

Other medicines that affect the kidneys

Your doctor will be especially careful if fampridine is given at the same time as any medicine which may affect how your kidneys eliminate medicines for example carvedilol, propranolol and metformin.

Fampridine Rowa with food and drink

Fampridine Rowa should be taken without food, on an empty stomach.

Pregnancy and breast-feeding

If you are pregnant, breast-feeding, think you may be pregnant or are planning to have a baby pregnant, ask your doctor or pharmacist for advice before taking this medicine
Fampridine Rowa is not recommended during pregnancy.

Your doctor will consider the benefit of you being treated with Fampridine Rowa against the risk to your baby.

You should not breast-feed whilst taking this medicine.

Driving and using machines

Fampridine Rowa may have an effect on people's ability to drive or use machines, it can cause dizziness. Make sure you're not affected before you start driving or use machinery.

3. How to take Fampridine Rowa

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. Fampridine is only available by prescription and under the supervision of doctors experienced in MS.

Your doctor will give you an initial prescription for 2 to 4 weeks. After 2 to 4 weeks the treatment will be reassessed.

The recommended dose is:

One tablet in the morning and **one** tablet in the evening (12 hours apart). Do not take more than two tablets in a day. **You must leave 12 hours** between each tablet. Do not take the tablets more often than every 12 hours.

Swallow each tablet whole, with a drink of water. Do not divide, crush, dissolve, suck or chew the tablet. This may increase your risk of side effects.

How to get the tablet out from the blister

Instructions on how to open the blister

1. Remove one blister by tearing down perforations.
2. Starting at one corner, carefully peel back the foil to reveal the tablet:

Note: The tablet cannot be pushed through the foil!



If you take more Fampridine Rowa than you should

Contact your doctor immediately if you take too many tablets.

Take the Fampridine Rowa box with you if you go to see the doctor.

In overdose you may notice sweating, minor shaking (*tremor*), confusion, memory loss (*amnesia*) and fits (*seizure*). You may also notice other effects not listed here.

If you forget to take Fampridine Rowa

If you forget to take a tablet, do not take two tablets at once to make up for a missed dose. You must **always leave 12 hours** between each tablet.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you have a seizure, stop taking Fampridine Rowa and tell your doctor immediately.

If you experience one or more of the following allergic (*hypersensitivity*) symptoms: swollen face, mouth, lips, throat or tongue, reddening or itching of the skin, chest tightness and breathing problems **stop taking Fampridine Rowa** and **see** your doctor immediately.

Side effects are listed below by frequency:

Very common side effects

May affect more than 1 in 10 people:

- Urinary tract infection

Common side effects

May affect up to 1 in 10 people:

- Feeling unsteady
- Dizziness
- Spinning sensation (vertigo)
- Headache
- Feeling weak and tired
- Difficulty sleeping
- Anxiety
- Minor shaking (*tremor*)
- Numbness or tingling of skin
- Sore throat
- Common cold (*nasopharyngitis*)
- Flu (*influenza*)
- Viral infection
- Difficulty breathing (shortness of breath)
- Feeling sick (*nausea*)
- Being sick (*vomiting*)
- Constipation
- Upset stomach
- Back pain
- Heartbeat that you can feel (*palpitations*)

Uncommon side effects

May affect up to 1 in 100 people

- Fits (*seizure*)
- Allergic reaction (*hypersensitivity*) Severe allergy (anaphylactic reaction)
- Swelling of the face, lips, mouth or tongue (angioedema)
- New onset or worsening of nerve pain in the face (*trigeminal neuralgia*)
- Fast heart rate (*tachycardia*)
- Dizziness or loss of consciousness (hypotension)
- Rash/itchy rash (urticaria)
- Chest discomfort

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance. Website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Fampridine Rowa

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the packaging after EXP. The expiry date refers to the last day of that month.

Store below 25°C. Store in the original package in order to protect from moisture

Do not throw away any medicines via waste water or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6 Contents of the pack and other information

What Fampridine Rowa contains

- **The active substance** is fampridine.
- Each prolonged-release tablet contains 10 mg of fampridine
- **The other ingredients** are:
- Tablet core: hypromellose, microcrystalline cellulose, silica colloidal anhydrous, magnesium stearate; film coat: Opadry White (Hypromellose, Titanium Dioxide (E171), Macrogol)

What Fampridine Rowa looks like and contents of the pack

Fampridine Rowa is a white to off-white colored, approx. 13 mm x 8 mm, oval shaped film coated tablets debossed with “L10” on one side and plain on the other.

Fampridine Rowa comes in Aluminium/Aluminium peelable blister with desiccant layer packed in cartons containing:

- 14 prolonged-release tablets
- 28 prolonged-release tablets
- 56 prolonged-release tablets
- 98 prolonged-release tablets

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:

Rowa Pharmaceuticals Ltd.,
Newtown,
Bantry,
Co.Cork.,
Ireland.

Manufacturer:

Laboratorios Liconsa, SA
PI Miralcampo, Av Miralcampo 7,
Azuqueca de Henares- Guadalajara,
Spain.

This medicinal product is authorised in the Member States of the EEA under the following

names:

Finland	Fampridine Laboratorios Liconsa 10 mg depottabletti
Germany	Fampiridin beta 10 mg Retardtabletten
Ireland	Fampridine Rowa 10 mg prolonged-release tablets

This leaflet was last revised in 07/2022