This EM contains important safety information about Invented name (Emtricitabine/tenofovir) and advice on risk minimisation.

This booklet was developed by Mylan Pharmaceuticals Limited.

DOB:

Patient Initials:

If you get any side effects talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. Please report side effects with any medicine directly via the national reporting system HPRA Pharmacovigilance, website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine. You should also report any side effects to Mylan Pharmaceuticals Limited Pharmacovigilance: pv.ireland@viatris.com.

## **Reminder Card**

Gender: M F

Age:

|                   |  | 9 |
|-------------------|--|---|
| een prescribed    |  |   |
| phylaxis) to help |  |   |

exposure prophylaxis) to help reduce the risk of getting HIV-1 infection. In order for this medicine to work effectively it is important that you don't miss or skip doses. The recommended dose is one Emtricitabine/Tenofovir disoproxil tablet every day. Take Emtricitabine/Tenofovir disoproxil exactly as your doctor has told you.

Whenever possible, Emtricitabine/Tenofovir disoproxil should be taken with food. To help you remember to take Emtricitabine/Tenofovir disoproxil:

- You can make taking medication part of your routine, for example taking it with your lunch, or when you brush your teeth after breakfast. It is important to find a time that works for you.
- You can add a reminder to your mobile phone or any other device can alert you when it is time to take Emtricitabine/Tenofovir disoproxil each day
- You can also use a small weekly pill box and put in the necessary tablets every week
- You might also find it helpful to use the calendar to tick off each day after you take Emtricitabine/Tenofovir disoproxil. Starting in your first week, tick the day when you start Emtricitabine/Tenofovir disoproxil, then tick the relevant box every day after you have taken Emtricitabine/Tenofovir disoproxil. You can also make note of the date when you took the first Emtricitabine/Tenofovir disoproxil tablet from a new bottle. If you are not sure if you took the tablet, you could count the number of tablets left in the bottle (each new bottle contains 30 tablets)

After 30 days (one bottle of tablets), you will need to start a new calendar.

If you already use a calendar on your phone or PC, you could add a daily reminder for Emtricitabine/Tenofovir disoproxil.

|        | Sun | Mon | Tue | Wed | Thu | Fri | Sat |
|--------|-----|-----|-----|-----|-----|-----|-----|
| Week 1 |     |     |     |     |     |     |     |
| Week 2 |     |     |     |     |     |     |     |
| Week 3 |     |     |     |     |     |     |     |
| Week 4 |     |     |     |     |     |     |     |
| Week 5 |     |     |     |     |     |     |     |

Date when you took the 1<sup>st</sup> Emtricitabine/Tenofovir disoproxil tablet from a new bottle: (day/month/year): / /

Appointment reminder

Your next clinic appointment is:

| Appointment | Date | Time | Location |
|-------------|------|------|----------|
|             |      |      |          |
|             |      |      |          |
|             |      |      |          |

Local version 1.0

NCA approval date 22-Dec-2021

