

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Anti-Hist Allergy 10mg Film-coated tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains Cetirizine dihydrochloride 10mg.

Excipient(s) with known effect

Each tablet contains 101.83 mg lactose monohydrate.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Film-coated tablet.

White circular biconvex film-coated tablets, embossed 'A' on one side and a deep score on the other.

The tablet can be divided into equal doses.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Anti-Hist Allergy is indicated in adults and children 6 years and above:

- for the relief of nasal and ocular symptoms of seasonal and perennial allergic rhinitis.
- for the relief of symptoms of chronic idiopathic urticaria.

4.2 Posology and method of administration

Posology

Adults and adolescents over 12 years of age

10 mg once daily (1 tablet).

Elderly

Data do not suggest that the dose needs to be reduced in elderly subjects provided that the renal function is normal.

Renal impairment

There are no data to document the efficacy/safety ratio in patients with renal impairment. Since cetirizine is mainly excreted via renal route (see section 5.2), in cases no alternative treatment can be used, the dosing intervals must be individualized according to renal function.

Refer to the following table and adjust the dose as indicated. To use this dosing table, an estimate of the patient's creatinine clearance (CL_{cr}) in ml/min is needed. The CL_{cr} (ml/min) may be estimated from serum creatinine (mg/dl) determination using the following formula:

$$\text{CL}_{\text{cr}} = \frac{[140 - \text{age (years)}] \times \text{weight (kg)}}{72 \times \text{serum creatinine (mg/dl)}} \quad (\times 0.85 \text{ for women})$$

Dosing adjustments for adult patients with impaired renal function

Group	Creatinine clearance (ml/min)	Dosage and frequency
Normal	> 80	10mg once daily
Mild	50-79	10mg once daily
Moderate	30-49	5mg once daily
Severe	<30	5mg once every 2 days
End-stage renal disease undergoing dialysis	<10	contra-indicated

In paediatric patients suffering from renal impairment, the dose will have to be adjusted on an individual basis taking into account the renal clearance of the patient and their body weight.

Hepatic impairment

No dose adjustment is needed in patients with solely hepatic impairment.

Simultaneous hepatic and renal impairment

Dose adjustment is recommended (see Patients with moderate to severe renal impairment above).

Paediatric population

Children aged from 6 to 12 years: 5 mg twice daily (a half tablet twice daily).

The tablet formulation should not be used in children under 6 years of age as it does not allow the necessary dose adjustments.

Method of administration

The tablets need to be swallowed with a glass of liquid.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Hypersensitivity to hydroxyzine or to any piperazine derivatives.

Patients with severe renal impairment at less than 10 ml/min creatinine clearance.

4.4 Special warnings and precautions for use

At therapeutic doses, no clinically significant interactions have been demonstrated with alcohol (for a blood alcohol level of 0.5 g/L). Nevertheless, precaution is recommended if alcohol is taken concomitantly.

Caution should be taken in patients with predisposition factors of urinary retention (e.g. spinal cord lesion, prostatic hyperplasia) as cetirizine may increase the risk of urinary retention.

Caution in epileptic patients and patients at risk of convulsions is recommended.

Pruritus and/or urticaria may occur when cetirizine is stopped, even if those symptoms were not present before treatment initiation. In some cases, the symptoms may be intense and may require treatment to be restarted. The symptoms should resolve when the treatment is restarted.

Paediatric population

The use of the film-coated tablet formulation is not recommended in children aged less than 6 years since this formulation does not allow for appropriate dose adaptation.

Allergy skin tests are inhibited by antihistamines and a wash-out period (of 3 days) is required before performing them.

This medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

This medicinal product contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interactions

Due to the pharmacokinetic, pharmacodynamic and tolerance profile of cetirizine, no interactions are expected with this antihistamine. Actually, neither pharmacodynamic nor significant pharmacokinetic interaction was reported in drug-drug interactions studies performed, notably with pseudoephedrine or theophylline (400 mg/day).

The extent of absorption of cetirizine is not reduced with food, although the rate of absorption is decreased.

In sensitive patients, the concurrent use of alcohol or other CNS depressants may cause additional reductions in alertness and impairment of performance, although cetirizine does not potentiate the effect of alcohol (0.5 g/L blood levels).

4.6 Fertility, pregnancy and lactation

Pregnancy

For cetirizine very rare clinical data on exposed pregnancies are available. Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/fetal development, parturition or postnatal development. Caution should be exercised when prescribing to pregnant women.

Breast-feeding

Cetirizine is excreted in human milk at concentrations representing 0.25 to 0.90 those measured in plasma, depending on sampling time after administration. Therefore, caution should be exercised when prescribing cetirizine to lactating women.

Fertility

Limited data is available on human fertility but no safety concern has been identified. Animal data show no safety concern for human reproduction.

4.7 Effects on ability to drive and use machines

Objective measurements of driving ability, sleep latency and assembly line performance have not demonstrated any clinically relevant effects at the recommended dose of 10 mg.

Patients intending to drive, engaging in potentially hazardous activities or operating machinery should not exceed the recommended dose and should take their response to the medicinal product into account. In these sensitive patients, concurrent use with alcohol or other CNS depressants may cause additional reductions in alertness and impairment of performance.

4.8 Undesirable effects

Clinical studies have shown that cetirizine at the recommended dosage has minor adverse effects on the CNS, including somnolence, fatigue, dizziness and headache. In some cases, paradoxical CNS stimulation has been reported.

Although cetirizine is a selective antagonist of peripheral H₁-receptors and is relatively free of anticholinergic activity, isolated cases of micturition difficulty, eye accommodation disorders and dry mouth have been reported.

Instances of abnormal hepatic function with elevated hepatic enzymes accompanied by elevated bilirubin have been reported. Mostly this resolves upon discontinuation of the treatment with cetirizine dihydrochloride.

a) Clinical trials

Adult Population

Double blind controlled clinical or pharmacoclinical trials comparing cetirizine to placebo or other antihistamines at the recommended dosage (10 mg daily for cetirizine), of which quantified safety data are available, included more than 3200 subjects exposed to cetirizine.

From this pooling, the following adverse events were reported for cetirizine 10 mg in the placebo-controlled trials at rates of 1.0 % or greater:

Adverse event (WHO-ART)	Cetirizine 10 mg (n= 3260)	Placebo (n = 3061)
<i>Body as a whole – general disorders</i> Fatigue	1.63%	0.95%
<i>Central and peripheral nervous system disorders</i> Dizziness Headache	1.10% 7.42%	0.98% 8.07%
<i>Gastro-intestinal system disorders</i> Abdominal pain Dry mouth Nausea	0.98% 2.09% 1.07%	1.08% 0.82% 1.14%
<i>Psychiatric disorders</i> Somnolence	9.63%	5.00%
<i>Respiratory system disorders</i> Pharyngitis	1.29%	1.34%

Although statistically more common than under placebo, somnolence was mild to moderate in the majority of cases. Objective tests as demonstrated by other studies have demonstrated that usual daily activities are unaffected at the recommended daily dose in healthy young volunteers.

Paediatric Population

Adverse drug reactions at rates of 1 % or greater in children aged from 6 months to 12 years, included in placebo-controlled clinical or pharmacoclinical trials are:

Adverse drug reactions (WHO-ART)	Cetirizine 10 mg (n= 1656)	Placebo (n = 1294)
<i>Gastro-intestinal system disorders</i> Diarrhoea	1.0%	0.6%
<i>Psychiatric disorders</i> Somnolence	1.8%	1.4%
<i>Respiratory system disorders</i> Rhinitis	1.4%	1.1%
<i>Body as a whole – general disorders</i> Fatigue	1.0%	0.3%

b) Post-marketing experience

In addition to the adverse effects reported during clinical studies and listed above, isolated cases of the following undesirable effects have been reported in post-marketing experience.

Undesirable effects are described according to MedDRA System Organ Class and by estimated frequency based on post-marketing experience.

Frequencies are defined as follows: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from available data).

Blood and lymphatic disorders

Very rare: thrombocytopenia

Immune system disorders

Rare: hypersensitivity

Very rare: anaphylactic shock

Metabolism and nutrition disorders

Not known: increased appetite

Psychiatric disorders

Uncommon: agitation

Rare: aggression, confusion, depression, hallucination, insomnia

Very rare: tics

Not known: suicidal ideation, nightmare

Nervous system disorders

Uncommon: paraesthesia

Rare: convulsions

Very rare: dysgeusia, syncope, tremor, dystonia, dyskinesia

Not known: amnesia, memory impairment

Eye disorders

Very rare: accommodation disorder, blurred vision, oculogyration

Ear and labyrinth disorders

Not known: vertigo

Cardiac disorders

Rare: tachycardia

Gastro-intestinal disorders

Uncommon: diarrhoea

Hepatobiliary disorders

Rare: hepatic function abnormal (increased transaminases, alkaline phosphatase, γ -GT and bilirubin)

Not known: hepatitis

Skin and subcutaneous tissue disorders

Uncommon: pruritus, rash

Rare: urticaria

Very rare: angioneurotic oedema, fixed drug eruption

Not known: acute generalized exanthematous pustulosis

Musculoskeletal and connective tissue disorders

Not known: arthralgia

Renal and urinary disorders

Very rare: dysuria, enuresis

Not known: urinary retention

General disorders and administration site conditions

Uncommon: asthenia, malaise

Rare: oedema

Investigations

Rare: weight increased

Description of selected adverse reactions

After discontinuation of cetirizine, pruritus (intense itching) and/or urticaria have been reported.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRC Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie.

4.9 Overdose

a) Symptoms

Symptoms observed after an overdose of cetirizine are mainly associated with CNS effects or with effects that could suggest an anticholinergic effect.

Adverse events reported after an intake of at least 5 times the recommended daily dose are: confusion, diarrhoea, dizziness, fatigue, headache, malaise, mydriasis, pruritus, restlessness, sedation, somnolence, stupor, tachycardia, tremor, and urinary retention.

b) Management

There is no known specific antidote to cetirizine.

Should overdose occur, symptomatic or supportive treatment is recommended. Gastric lavage should be considered following ingestion of a short occurrence.

Cetirizine is not effectively removed by dialysis.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Piperazine derivatives, ATC code: R06AE07

Mechanism of action

Cetirizine, a human metabolite of hydroxyzine, is a potent and selective antagonist of peripheral H₁-receptors. *In vitro* receptor binding studies have shown no measurable affinity for other than H₁-receptors.

Pharmacodynamic effects

In addition to its anti-H₁ effect, cetirizine was shown to display anti-allergic activities: at a dose of 10 mg once or twice daily, it inhibits the late phase recruitment of eosinophils, in the skin and conjunctiva of atopic subjects submitted to allergen challenge.

Clinical efficacy and safety

Studies in healthy volunteers show that cetirizine, at doses of 5 and 10 mg strongly inhibits the wheal and flare reactions induced by very high concentrations of histamine into the skin, but the correlation with efficacy is not established.

In a six-week, placebo-controlled study of 186 patients with allergic rhinitis and concomitant mild to moderate asthma, cetirizine 10 mg once daily improved rhinitis symptoms and did not alter pulmonary function. This study supports the safety of administering cetirizine to allergic patients with mild to moderate asthma.

In a placebo-controlled study, cetirizine given at the high daily dose of 60 mg for seven days did not cause statistically significant prolongation of QT interval.

At the recommended dosage, cetirizine has demonstrated that it improves the quality of life of patients with perennial and seasonal allergic rhinitis.

Paediatric population

In a 35-day study in children aged 5 to 12, no tolerance to the antihistaminic effect (suppression of wheal and flare) of cetirizine was found.

When a treatment with cetirizine is stopped after repeated administration, the skin recovers its normal reactivity to histamine within 3 days.

5.2 Pharmacokinetic properties

Absorption

The steady - state peak plasma concentrations is approximately 300 ng/ml and is achieved within 1.0 ± 0.5 h.

The distribution of pharmacokinetic parameters such as peak plasma concentration (C_{max}) and area under curve (AUC), is unimodal in human volunteers.

The extent of absorption of cetirizine is not reduced with food, although the rate of absorption is decreased. The extent of bioavailability is similar when cetirizine is given as solutions, capsules or tablets.

Distribution

The apparent volume of distribution is 0.50 l/kg. Plasma protein binding of cetirizine is 93 ± 0.3 %. Cetirizine does not modify the protein binding of warfarin.

Biotransformation

Cetirizine does not undergo extensive first pass metabolism.

Elimination

About two third of the dose are excreted unchanged in urine. The terminal half-life is approximately 10 hours. No accumulation is observed for cetirizine following daily doses of 10 mg for 10 days.

Linearity/non-linearity

Cetirizine exhibits linear kinetics over the range of 5 to 60 mg.

Special populations

Elderly: Following a single 10 mg oral dose, half-life increased by about 50 % and clearance decreased by 40 % in 16 elderly subjects compared to the normal subjects. The decrease in Cetirizine clearance in these elderly volunteers appeared to be related to their decreased renal function.

Paediatric population: The half-life of cetirizine was about 6 hours in children of 6-12 years and 5 hours in children 2-6 years. In infants and toddlers aged 6 to 24 months, it is reduced to 3.1 hours

Renal impairment: The pharmacokinetics of the drug were similar in patients with mild impairment (creatinine clearance higher than 40 ml/min) and healthy volunteers. Patients with moderate renal impairment had a 3-fold increase in half-life and 70 % decrease in clearance compared to healthy volunteers.

Patients on hemodialysis (creatinine clearance less than 7 ml/min) given a single oral 10 mg dose of cetirizine had a 3-fold increase in half-life and a 70 % decrease in clearance compared to normals.

Cetirizine was poorly cleared by haemodialysis. Dosing adjustment is necessary in patients with moderate or severe renal impairment (see section 4.2).

Hepatic impairment : Patients with chronic liver diseases (hepatocellular, cholestatic, and biliary cirrhosis) given 10 or 20 mg of cetirizine as a single dose had a 50 % increase in half-life along with a 40 % decrease in clearance compared to healthy subjects.

Dosing adjustment is only necessary in patients with hepatic impairment if concomitant renal impairment is present.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Lactose monohydrate
Microcrystalline Cellulose
Maize Starch
Colloidal Anhydrous Silica
Magnesium Stearate
Talc

Film-coat:
Hypromellose
Lactose monohydrate
Titanium dioxide (E171)
Macrogol 4000
Sodium Citrate

6.2 Incompatibilities

Not applicable

6.3 Shelf life

3 years.

6.4 Special precautions for storage

This medicinal product does not require any special precautions for storage.

6.5 Nature and contents of container

Blister strips composed of transparent PVC and aluminium foil.
Pack sizes: 7, 10, 14, 21, 28, 30 tablets

Not all pack sizes may be marketed

6.6 Special precautions for disposal

No special requirements.

7 MARKETING AUTHORISATION HOLDER

Clonmel Healthcare Ltd
Clonmel
Co. Tipperary
Ireland

8 MARKETING AUTHORISATION NUMBER

PA0126/267/001
13 August 2019

CRN008YTZ

Page 8 of 9

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of First Authorisation: 23rd June 2006

Date of Last Renewal: 23rd June 2011

10 DATE OF REVISION OF THE TEXT

August 2019