

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Keral 25 mg oral solution in sachet

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each sachet of oral solution contains: dexketoprofen 25 mg as dexketoprofen trometamol.
Excipients with known effect: 2 g of sucrose and 20 mg of methyl parahydroxybenzoate (E 218).

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Oral solution in sachet.
Slightly coloured solution with lemon odour and sweet lemon-citrus flavour.

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Short term symptomatic treatment of acute pain of mild to moderate intensity, such as acute musculo-skeletal pain, dysmenorrhoea, dental pain.
This medicinal product is indicated in adult patients.

4.2 Posology and method of administration

Posology

The lowest effective dose should be used for the shortest duration necessary to relieve symptoms (see section 4.4).

Adults:

According to the nature and severity of pain, the recommended dosage is generally 25 mg every 8 hours. The total daily dose should not exceed 75 mg.

This medicinal product is only intended for short term use and the treatment must be limited to the symptomatic period.

Elderly:

In elderly it is recommended to start the therapy at the lower end of the dosage range (50 mg total daily dose). The dosage may be increased to that recommended for the general population only after good general tolerance has been ascertained. Due to the possible adverse effect profile (see section 4.4), elderly individuals should receive particularly close monitoring.

Hepatic impairment:

Patients with mild to moderate hepatic dysfunction should start therapy at reduced doses (50 mg total daily dose) and be closely monitored. Dexketoprofen should not be used in patients with severe hepatic dysfunction.

Renal impairment:

The initial dosage should be reduced to 50 mg total daily dose in patients with mildly impaired renal function (creatinine clearance 60 – 89 ml / min) (see section 4.4). Dexketoprofen should not be used in patients with moderate to severe renal dysfunction (creatinine clearance \leq 59 ml / min) (see section 4.3).

Paediatric population:

Dexketoprofen has not been studied in children and adolescent. Therefore the safety and efficacy in children and adolescents have not been established and the product should not be used in children and adolescent.

Method of administration

Oral use. The oral solution may be taken directly from the sachet or after stirring the whole content in a glass of water. Once the sachet opens, the entire content should be taken.

Concomitant administration with food delays the absorption rate of the drug (see Pharmacokinetic Properties), thus in case of acute pain it is recommended that administration is at least 15 minutes before meals.

4.3 Contraindications

- patients hypersensitive to the active substance, to any other NSAID, or to any of the excipients listed in section 6.1.
- patients in whom substances with a similar action (e.g. acetylsalicylic acid, or other NSAIDs) precipitate attacks of asthma, bronchospasm, acute rhinitis, or cause nasal polyps, urticaria or angioedema.
- known photoallergic or phototoxic reactions during treatment with ketoprofen or fibrates
- patients with history of gastrointestinal bleeding or perforation related to previous NSAIDs therapy.
- patients with active peptic ulcer/gastrointestinal haemorrhage or any history of gastrointestinal bleeding, ulceration or perforation.
- patients with chronic dyspepsia.
- patients who have other active bleedings or bleeding disorders.
- patients with Crohn's disease or ulcerative colitis.
- patients with severe heart failure.
- patients with moderate to severe renal dysfunction (creatinine clearance ≤ 59 ml/min).
- patients with severely impaired hepatic function (Child-Pugh score 10 - 15).
- patients with haemorrhagic diathesis and other coagulation disorders.
- patients with severe dehydration (caused by vomiting, diarrhoea or insufficient fluid intake).
- during the third trimester of pregnancy and lactation period (see section 4.6).

4.4 Special warnings and precautions for use

Administer with caution in patients with a history of allergic conditions.

The concomitant use of dexketoprofen with other NSAIDs including cyclooxygenase-2 selective inhibitors should be avoided.

Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms (see section 4.2, and GI and cardiovascular risks below).

Gastrointestinal safety

Gastrointestinal bleeding, ulceration or perforation which can be fatal, have been reported with all NSAIDs at anytime during treatment, with or without warning symptoms or a previous history of serious gastrointestinal events. When gastrointestinal bleeding or ulceration occurs in patients receiving dexketoprofen, the treatment should be withdrawn.

The risk of gastrointestinal bleeding, ulceration or perforation is higher with increasing NSAID doses, in patients with a history of ulcer, particularly if complicated with haemorrhage or perforation (see section 4.3), and in the elderly.

Use in elderly: The elderly have an increased frequency of adverse reactions to NSAIDs especially gastrointestinal bleeding and perforation which may be fatal (see section 4.2). These patients should commence treatment on the lowest dose available.

As with all NSAIDs, any history of oesophagitis, gastritis and/or peptic ulcer must be sought in order to ensure their total cure before starting treatment with dexketoprofen. Patients with gastrointestinal symptoms or history of gastrointestinal disease should be monitored for digestive disturbances, especially gastrointestinal bleeding.

NSAIDs should be given with care to patients with a history of gastrointestinal disease (ulcerative colitis, Crohn's disease) as their condition may be exacerbated (see section 4.8).

Combination therapy with protective agents (e.g. misoprostol or proton pump inhibitors) should be considered for these patients, and also for patients requiring concomitant low dose acetylsalicylic acid, or other drugs likely to increase gastrointestinal risk (see below and section 4.5).

Patients with a history of gastrointestinal toxicity, particularly when elderly, should report any unusual abdominal symptoms (especially gastrointestinal bleeding) particularly in the initial stages of treatment.

Caution should be advised in patients receiving concomitant medications which could increase the risk of ulceration or bleeding, such as oral corticosteroids, anticoagulants such as warfarin, selective serotonin-reuptake inhibitors or anti-platelet agents such as acetylsalicylic acid (see section 4.5).

Renal Safety

Caution should be exercised in patients with impairment of renal functions. In these patients, the use of NSAIDs may result in deterioration of renal function, fluid retention and oedema. Caution is also required in patients receiving diuretic therapy or those who could develop hypovolaemia as there is an increased risk of nephrotoxicity.

Adequate fluid intake should be ensured during treatment to prevent dehydration and possibly associated increased renal toxicity.

As with all NSAIDs, it can increase plasma urea nitrogen and creatinine. As with other inhibitors of prostaglandin synthesis, it can be associated with adverse effects on the renal system which can lead to glomerular nephritis, interstitial nephritis, renal papillary necrosis, nephrotic syndrome and acute renal failure.

Elderly patients are more likely to be suffering from impaired renal function (see section 4.2).

Liver Safety

Caution should be exercised in patients with impairment of hepatic functions.

As with other NSAIDs, it can cause transient small increases in some liver parameters, and also significant increases in SGOT and SGPT. In case of a relevant increase in such parameters, therapy must be discontinued.

Elderly patients are more likely to be suffering from impaired hepatic function (see section 4.2).

Cardiovascular and cerebrovascular safety

Appropriate monitoring and advice are required for patients with history of hypertension and/or mild to moderate heart failure. Special caution should be exercised in patients with a history of cardiac disease, in particular those with previous episodes of heart failure as there is an increased risk of triggering heart failure, since fluid retention and oedema have been reported in association with NSAIDs therapy.

Clinical trial and epidemiological data suggest that use of some NSAIDs (particularly at high doses and in long term treatment) may be associated with a small increased risk of arterial thrombotic events (e.g myocardial infarction or stroke). There are insufficient data to exclude such a risk for dexketoprofen.

Consequently, patients with uncontrolled hypertension, congestive heart failure, established ischaemic heart disease, peripheral arterial disease, and/or cerebrovascular disease should only be treated with dexketoprofen after careful consideration. Similar consideration should be made before initiating longer-term treatment of the patients with risk factors for cardiovascular disease (e.g. hypertension, hyperlipidaemia, diabetes mellitus, smoking).

All non-selective NSAIDs can inhibit platelet aggregation and prolong bleeding time via inhibition of prostaglandin synthesis. Therefore, the use of dexketoprofen in patients who are receiving other therapy that interferes with haemostasis, such as warfarin or other coumarins or heparins is not recommended (see section 4.5).

Elderly patients are more likely to be suffering from impaired cardiovascular function (see section 4.2).

Skin reactions

Serious skin reactions, some of them fatal, including exfoliative dermatitis, Stevens-Johnson syndrome, and toxic epidermal necrolysis, have been reported very rarely in association with the use of NSAIDs (see section 4.8). Patients appear to be at highest risk of these reactions early in the course of therapy, the onset of the reaction occurring in the majority of cases within the first month of treatment. Keral should be discontinued at the first appearance of skin rash, mucosal lesions, or any other sign of hypersensitivity.

Masking of symptoms of underlying infections

Dexketoprofen can mask symptoms of infection, which may lead to delayed initiation of appropriate treatment and thereby worsening the outcome of the infection. This has been observed in bacterial community acquired pneumonia and bacterial complications to varicella. When this medicine is administered for pain relief in relation to infection, monitoring of infection is advised. In non-hospital settings, the patient should consult a doctor if symptoms persist or worsen.

Other information

Particular caution is required in patients with:

- congenital disorder of porphyrin metabolism (e.g. acute intermittent porphyria)
- dehydration
- directly after major surgery

If the physician considers long-term dexketoprofen therapy to be necessary, hepatic and renal function and the blood count should be regularly checked.

Severe acute hypersensitivity reactions (anaphylactic shock, for example) have been observed on very rare occasions. Treatment must be discontinued at the first signs of severe hypersensitivity reactions following intake of dexketoprofen. Depending on the symptoms, any medically required procedures must be initiated by specialist healthcare professionals.

Patients with asthma combined with chronic rhinitis, chronic sinusitis, and/or nasal polyposis have a higher risk of allergy to acetylsalicylic acid and/or NSAIDs than the rest of the population. Administration of this medicinal product can cause asthma attacks or bronchospasm, particularly in subjects allergic to acetylsalicylic acid or NSAIDs (see section 4.3).

Exceptionally, varicella can be at the origin of serious cutaneous and soft tissues infectious complications. To date, the contributing role of NSAIDs in the worsening of these infections cannot be ruled out. Thus, it is advisable to avoid use of dexketoprofen in case of varicella.

Dexketoprofen should be administered with caution to patients suffering from haematopoietic disorders, systemic lupus erythematosus or mixed connective tissue disease.

Through concomitant intake of alcohol, active substance-related undesirable effects, particularly those that concern the gastrointestinal tract or the central nervous system, may be increased on use of NSAIDs.

This medicine may cause allergic reactions (possibly delayed) as it contains methyl parahydroxybenzoate (E 218).

This medicinal product contains sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine. This should be taken into account in patients with diabetes mellitus.

This medicine contains less than 1 mmol sodium (23 mg) per sachet, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interactions

The following interactions apply to non-steroidal anti-inflammatory drugs (NSAIDs) in general:

Inadvisable combinations:

- Other NSAIDs, (including cyclooxygenase-2 selective inhibitors) and high doses of salicylates (≥ 3 g/day): administration of several NSAIDs together may increase the risk of gastrointestinal ulcers and bleeding, via a synergistic effect.
- Anticoagulants: NSAIDs may enhance the effects of anti-coagulants, such as warfarin (see section 4.4), due to the high plasma protein binding of dexketoprofen and the inhibition of platelet function and damage to the gastroduodenal mucosa. If the combination cannot be avoided, close clinical observation and monitoring of laboratory values should be carried out.
- Heparins: increased risk of haemorrhage (due to the inhibition of platelet function and damage to the gastroduodenal mucosa). If the combination cannot be avoided, close clinical observation and monitoring of laboratory values should be carried out.
- Corticosteroids: there is an increased risk of gastrointestinal ulceration or bleeding (see section 4.4).
- Lithium (described with several NSAIDs): NSAIDs increase blood lithium levels, which may reach toxic values (decreased renal excretion of lithium). This parameter therefore requires monitoring during the initiation, adjustment and withdrawal of treatment with dexketoprofen.
- Methotrexate, used at high doses of 15 mg/week or more: increased haematological toxicity of methotrexate via a decrease in its renal clearance by NSAIDs in general.
- Hydantoines and sulphonamides: the toxic effects of these substances may be increased. Combinations requiring precautions:
- Diuretics, ACE inhibitors, antibacterial aminoglycosides and angiotensin II receptor antagonists: Dexketoprofen may reduce the effect of diuretics and antihypertensive drugs. In some patients with compromised renal function (e.g. dehydrated patients or elderly with compromised renal function), the coadministration of agents that inhibit cyclo-oxygenase and ACE inhibitor or angiotensin II receptor antagonists or antibacterial aminoglycosides may result in further deterioration of renal function, which is usually reversible. In case of combined prescription of dexketoprofen and a diuretic, it is essential to ensure that the patient is adequately hydrated and to monitor renal function at the start of the treatment and periodically thereafter. Co-administration of dexketoprofen and potassium-sparing diuretics can lead to hyperkalaemia. Monitoring of blood potassium concentrations is required (see section 4.4).
- Methotrexate, used at low doses, less than 15 mg/week: increased haematological toxicity of methotrexate via a decrease in its renal clearance by antiinflammatory agents in general. Weekly monitoring of blood count during the first weeks of the combination. Increased surveillance in the presence of even mildly impaired renal function, as well as in the elderly.
- Pentoxifylline: increased risk of bleeding. Increase clinical monitoring and check bleeding time more often.

- Zidovudine: risk of increased red cell line toxicity via action on reticulocytes, with severe anaemia occurring one week after the NSAID is started. Check complete blood count and reticulocyte count one to two weeks after starting treatment with the NSAID.
- Sulfonylureas: NSAIDs can increase the hypoglycaemic effect of sulfonylureas by displacement from plasma protein binding sites. Combinations needing to be taken into account:
- Beta-blockers: treatment with a NSAID may decrease their antihypertensive effect via inhibition of prostaglandin synthesis.
- Cyclosporin and tacrolimus: nephrotoxicity may be enhanced by NSAIDs via renal prostaglandin mediated effects. During combination therapy, renal function has to be measured.
- Thrombolytics: increased risk of bleeding.
- Anti-platelet agents and selective serotonin reuptake inhibitors (SSRIs): increased risk of ulcer or gastrointestinal bleeding (see section 4.4).
- Probenecid: plasma concentrations of dexketoprofen may be increased; this interaction can be due to an inhibitory mechanism at the site of renal tubular secretion and of glucuronconjugation and requires adjustment of the dose of dexketoprofen.
- Cardiac glycosides: NSAIDs may increase plasma glycoside concentration.
- Mifepristone: there is a theoretical risk that prostaglandin synthetase inhibitors may alter the efficacy of mifepristone. Limited evidence suggests that co-administration of NSAIDs on the day of prostaglandin administration does not adversely influence the effects of mifepristone or the prostaglandin on cervical ripening or uterine contractility and does not reduce the clinical efficacy of medical termination of pregnancy.
- Quinolone antibiotics: animal data indicate that high doses of quinolones in combination with NSAIDs can increase the risk of developing convulsions.
- Tenofovir: concomitant use with NSAID can increase plasma urea nitrogen and creatinine, renal function should be monitored in order to control a potential synergic influence on renal function.
- Deferasirox: concomitant use with NSAIDs can increase the risk of gastrointestinal toxicity. Close clinical monitoring is required when deferasirox is combined with these substances.
- Pemetrexed: concomitant use with NSAIDs may decrease pemetrexed elimination, therefore caution should be made when administering higher doses of NSAIDs. In patients with mild to moderate renal insufficiency (creatinine clearance from 45 to 79 ml/min), the concomitant administration of pemetrexed with NSAIDs doses should be avoided for 2 days before and 2 days following pemetrexed administration.

4.6 Fertility, pregnancy and lactation

Dexketoprofen use is contraindicated during third trimester of pregnancy and lactation (see section 4.3).

Pregnancy

Inhibition of prostaglandin synthesis may adversely affect the pregnancy and/or the embryo/foetal development. Data from epidemiological studies raise concern about an increased risk of miscarriage and of cardiac malformation and gastroschisis after use of a prostaglandin synthesis inhibitor in early pregnancy. The absolute risk for cardiovascular malformation was increased from less than 1%, up to approximately 1.5%. The risk is believed to increase with dose and duration of therapy. In animals, administration of a prostaglandin synthesis inhibitor has been shown to result in increased pre- and post-implantation loss and embryo-foetal lethality.

In addition, increased incidences of various malformations including cardiovascular, have been reported in animals given a prostaglandin synthesis inhibitor during the organogenetic period (see section 5.3). During the first and second trimester of pregnancy, dexketoprofen trometamol should not be given unless clearly necessary. If dexketoprofen is used by a woman attempting to conceive, or during the first and second trimester of pregnancy, the dose should be kept as low and duration of treatment as short as possible.

During the third trimester of pregnancy, all prostaglandin synthesis inhibitors may expose:

- the fetus to:

- cardiopulmonary toxicity (with premature closure of the ductus arteriosus and pulmonary hypertension);
- renal dysfunction, which may progress to renal failure with oligo-hydroamniosis;
- the mother and the neonate, at the end of pregnancy, to:
- possible prolongation of bleeding time, an anti-aggregating effect which may occur even at very low doses;
- inhibition of uterine contractions resulting in delayed or prolonged labour.

Breast-feeding

It is not known whether dexketoprofen is excreted in human milk. Its use is contraindicated during breast-feeding (see section 4.3).

Fertility

As with other NSAIDs, the use of dexketoprofen may impair female fertility and is not recommended in women attempting to conceive. In women who have difficulties conceiving or who are undergoing investigation of infertility, withdrawal of dexketoprofen should be considered.

4.7 Effects on ability to drive and use machines

This medicinal product may cause undesirable effects such as dizziness, visual disturbances or drowsiness. The ability to react and the ability to take part actively in road traffic and to operate machines may be impaired in these cases

4.8 Undesirable effects

The adverse events reported as at least possibly related with dexketoprofen in clinical trials (tablet formulation), as well as the adverse reactions reported after the marketing of Keral oral solution in sachet are tabulated below, classified by system organ class and ordered by frequency:

Because C_{max} plasma levels of dexketoprofen for the oral solution formulation are higher than those reported for the tablet formulation, a potentially increased risk for adverse (gastrointestinal) events cannot be excluded.

SYSTEM ORGAN CLASS	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Very rare (<1/10,000)
Blood and lymphatic system disorders	---	---	---	Neutropenia, thrombocytopenia
Immune system disorders	---	---	Laryngeal oedema	Anaphylactic reaction, including anaphylactic shock
Metabolism and nutrition disorders	---	---	Anorexia	---
Psychiatric disorders	---	Insomnia, anxiety	---	---
Nervous system disorders	---	Headache, dizziness, somnolence	Paraesthesia, syncope	---
Eye disorders	---	---	---	Blurred vision
Ear and labyrinth disorders	---	Vertigo	---	Tinnitus
Cardiac disorders	---	Palpitations	---	Tachycardia
Vascular disorders	---	Flushing	Hypertension	Hypotension
Respiratory, thoracic and mediastinal disorders	---	---	Bradypnoea	Bronchospasm, dyspnoea
Gastrointestinal disorders	Nausea and/or vomiting, abdominal pain, diarrhoea, dyspepsia	Gastritis, constipation, dry mouth, flatulence	Peptic ulcer, peptic ulcer haemorrhage or peptic ulcer perforation (see section 4.4)	Pancreatitis
Hepatobiliary disorders	---	---	Hepatocellular injury	

Skin and subcutaneous tissue disorders	---	Rash	Urticaria, acne, sweating increased	Stevens Johnson syndrome, toxic epidermal necrolysis (Lyell's syndrome), angioedema, facial oedema, photosensitivity reaction, pruritus
Musculoskeletal and connective tissue disorders	---	---	Back pain	---
Renal and urinary disorders	---	---	Acute renal failure, Polyuria	Nephritis or nephrotic syndrome
Reproductive system and breast disorders	---	---	Menstrual disorder, prostatic disorder	---
General disorders and administration site conditions	---	Fatigue, pain, asthenia, rigors, malaise	Peripheral oedema	---
Investigations	---	---	Liver function test abnormal	---

The most commonly-observed adverse events are gastrointestinal in nature. Peptic ulcers, perforation or gastrointestinal bleeding, sometimes fatal, particularly in the elderly, may occur (see section 4.4 Special warnings and precautions for use). Nausea, vomiting, diarrhoea, flatulence, constipation, dyspepsia, abdominal pain, melaena, haematemesis, ulcerative stomatitis, exacerbation of colitis and Crohn's disease (see section 4.4) have been reported following administration. Less frequently, gastritis has been observed. Oedema, hypertension and cardiac failure have been reported in association with NSAIDs treatment.

As with other NSAIDs the following undesirable effects may appear aseptic meningitis, which might predominantly occur in patients with systemic lupus erythematosus or mixed connective tissue disease; haematological reactions (purpura, aplastic and haemolytic anaemia, and rarely agranulocytosis and medullar hypoplasia).

Bullous reactions including Stevens Johnson Syndrome and Toxic Epidermal Necrolysis (very rare).

Clinical trial and epidemiological data suggest that use of some NSAIDs (particularly at high doses and in long term treatment) may be associated with a small increased risk of arterial thrombotic events (e.g myocardial infarction or stroke; see section 4.4).

Reporting of suspected adverse reactions.

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme. Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

4.9 Overdose

The symptomatology following overdose is not known. Similar medicinal products have produced gastrointestinal (vomiting, anorexia, abdominal pain) and neurological (somnia, vertigo, disorientation, headache) disorders.

In case of accidental or excessive intake, immediately institute symptomatic therapy according to the patient's clinical condition. Activated charcoal should be administered if more than 5 mg/kg has been ingested by an adult or a child within an hour.

Dexketoprofen trometamol may be removed by dialysis.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: propionic acid derivatives
ATC code: M01AE17.

Dexketoprofen trometamol is the tromethamine salt of S-(+)-2-(3-benzoylphenyl)propionic acid, an analgesic, anti-inflammatory and antipyretic drug, which belongs to the non-steroidal anti-inflammatory group of drugs (M01AE).

Mechanism of action

The mechanism of action of non-steroidal antiinflammatory drugs is related to the reduction of prostaglandin synthesis by the inhibition of cyclooxygenase pathway. Specifically, there is an inhibition of the transformation of arachidonic acid into cyclic endoperoxides, PGG₂ and PGH₂, which produce prostaglandins PGE₁, PGE₂, PGF_{2a} and PGD₂ and also prostacyclin PGI₂ and thromboxanes (TxA₂ and TxB₂). Furthermore, the inhibition of the synthesis of prostaglandins could affect other inflammation mediators such as kinins, causing an indirect action which would be additional to the direct action.

Pharmacodynamic effects

Dexketoprofen has been demonstrated to be an inhibitor for COX-1 and COX-2 activities in experimental animals and humans.

Clinical efficacy and safety

Clinical studies performed on several pain models demonstrated effective analgesic activity of dexketoprofen. The onset of the analgesic activity was obtained in some studies at 30 minutes post-administration. The analgesic effect persists for 4 to 6 hours.

5.2 Pharmacokinetic properties

Two bioequivalence studies on healthy volunteers were carried out to compare dexketoprofen 25mg oral solution in sachet taken with and without water versus the film-coated tablets.

The comparison between dexketoprofen oral tablet and oral solution taken with water indicated that the two formulations were bioequivalent in terms of total exposure (AUC). Peak concentrations (C_{max}) were approximately 20% higher for the oral solution in comparison to the tablet.

The comparison between dexketoprofen tablet and oral solution taken without water indicated that the two formulations were bioequivalent in terms of both AUC and C_{max} .

Absorption

Dexketoprofen is rapidly absorbed after oral administration, showing maximum plasma concentrations achieved after 15-20 minutes (range 10 - 75 minutes) when administered as oral solution.

When administered concomitantly with food, the AUC does not change, however the C_{max} of dexketoprofen decreases and its absorption rate is delayed (increased t_{max}).

Distribution

The distribution half-life and elimination half-life values of dexketoprofen are 0.35 and 1.65 hours, respectively. As with other drugs with a high plasma protein binding (99%), its volume of distribution has a mean value below 0.25 l/kg. In multiple-dose pharmacokinetic studies, it was observed that the AUC after the last administration is not different from that obtained following a single dose, indicating that no drug accumulation occurs.

Biotransformation and elimination

After administration of dexketoprofen only the S-(+) enantiomer is obtained in urine, demonstrating that no conversion to the R-(-) enantiomer occurs in humans.

The main elimination route for dexketoprofen is glucuronide conjugation followed by renal excretion.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, genotoxicity and immunopharmacology. The chronic toxicity studies carried out in mice and monkeys gave a No Observed Adverse Effect Level (NOAEL) at doses 2 fold higher than maximum recommended human dose. In monkey, at higher doses, the main adverse effect observed were blood in faeces, decreased body weight gain and, at the highest dose, erosive gastrointestinal lesions. These effects appeared at doses determining a drug exposure 14-18 fold higher than that at the maximum recommended human dose.

There are no studies on the carcinogenic potential in animals.

As it has been recognised for the whole pharmacological class of NSAIDs, dexketoprofen may cause changes of embryo-foetal survival in animal models, both indirectly, through the gastrointestinal toxicity on the pregnant mothers, and directly upon the development of the foetus.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Ammonium glycyrrhizate
Neohesperidin-dihydrochalcone
Methyl parahydroxybenzoate (E 218)
Saccharin sodium
Sucrose
Macrogol 400
Lemon aroma
Povidone K-90
Anhydrous disodium phosphate
Sodium dihydrogen phosphate dihydrate
Purified water

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

Single dose sachets made of laminated polyester/ aluminium/ low density polyethylene foil. Each sachet contains 10 ml oral solution.

Each pack contains 2, 4, 10 or 20 single-dose sachets.

Not all pack size may be marketed.

6.6 Special precautions for disposal and other handling

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

Menarini International Operations Luxembourg S.A.
1, Avenue de la Gare
1611 Luxembourg
Luxembourg

8 MARKETING AUTHORISATION NUMBER

PA0865/002/006

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 12th May 2017

Date of last renewal: 15th February 2021

10 DATE OF REVISION OF THE TEXT

August 2021

31 August 2021

CRN00C45M

Page 9 of 9