

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Cyclimorph 10 solution for injection

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Cyclimorph 10 Injection contains morphine tartrate 10 mg and cyclizine tartrate 50 mg (equivalent to 39.01 mg cyclizine) in each 1 ml ampoule.

Excipient with known effect:

Sodium metabisulphite (E223) (1 mg /1 ml ampoule).

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Solution for Injection (Injection)

A clear very slightly coloured solution. pH 4.3 - 5.0

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Cyclimorph Injection is indicated in adults and children for the relief of moderate to severe pain in all suitable medical and surgical conditions (see section 4.3 and 4.4) in which reduction of the nausea and vomiting associated with the administration of morphine is required.

4.2 Posology and method of administration

Posology

Adults and children over 12 years:

The usual dose is 10-20 mg morphine tartrate, given subcutaneously, intramuscularly or intravenously. Additional doses may not be given more frequently than 4-hourly.

Not more than 3 doses (representing 150 mg cyclizine tartrate: i.e. 3 ml of Cyclimorph 10Injection) should be given in any 24-hour period.

Elderly:

Morphine doses should be reduced in elderly patients and titrated to provide optimal pain relief with minimal side effects since:

- Increased duration of pain relief from a standard dose of morphine has been reported in elderly patients.
- A review of pharmacokinetic studies has suggested that morphine clearance decreases and half-life increases in older patients.
- The elderly may be particularly sensitive to the adverse effects of morphine.

Paediatric population

Children 6-12 years: 5-10 mg morphine tartrate as a maximum single dose.

Children 1-5 years: 2.5-5 mg morphine tartrate as a maximum single dose.

Method of administration

Subcutaneous, intramuscular or intravenous injection.

When used intravenously, Cyclimorph Injection should be injected slowly into the bloodstream, over a period of 4 to 5 minutes with the patient in the recumbent position, with only minimal withdrawal of blood into the syringe.

Discontinuation of therapy

An abstinence syndrome may be precipitated if opioid administration is suddenly discontinued. Therefore the dose should be gradually reduced prior to discontinuation.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Respiratory depression or obstructive airways disease. Patients with excessive bronchial secretions should not be given Cyclimorph Injection as morphine diminishes the cough response.

An attack of bronchial asthma or in heart failure secondary to chronic lung disease.

Head injury or raised intra-cranial pressure.

Acute alcohol intoxication. The antiemetic properties of cyclizine may increase the toxicity of alcohol.

Concomitant therapy with monoamine oxidase inhibitors or within 14 days of stopping such treatment.

Ulcerative colitis, since such preparations may precipitate toxic dilation or spasm of the colon.
Paralytic ileus and delayed gastric emptying.

Biliary and renal tract spasm and in patients immediately after operative interventions in the biliary tract.

Renal impairment:

Severe and prolonged respiratory depression may occur in patients with renal impairment given morphine; this is attributed to the accumulation of the active metabolite morphine-6-glucuronide. Therefore, Cyclimorph Injection should not be administered to patients with moderate or severe renal impairment (glomerular filtration rate <20 ml/min).

Hepatic impairment:

As with other opioid analgesic containing preparations Cyclimorph Injection should not be administered to patients with severe hepatic impairment as it may precipitate coma.

4.4 Special warnings and precautions for use

Dependence and withdrawal (abstinence) syndrome

Use of opioid analgesics may be associated with the development of physical and/or psychological dependence or tolerance. The risk increases with the time the drug is used, and with higher doses. Symptoms can be minimised with adjustments of dose or dosage form, and gradual withdrawal of morphine. For individual symptoms, see section 4.8.

Cyclimorph Injection has an abuse potential similar to other strong agonist opioids, and should be used with particular caution in patients with a history of alcohol or drug abuse.

Cyclimorph Injection should be used with caution in the debilitated since they may be more sensitive to the respiratory depressant effects.

Cyclimorph Injection should be used with caution in the presence of the following: convulsive disorders, delirium tremens, severe cor pulmonale, hypothyroidism, adrenocortical insufficiency, hypopituitarism, myxoedema, prostatic hypertrophy, shock, diabetes mellitus, myasthenia gravis, hypotension and hypovolaemia, pancreatitis. Obstructive bowel disorders and inflammatory bowel disorders.

Morphine may lower the seizure threshold in patients with a history of epilepsy.

Extreme caution should be exercised when administering Cyclimorph Injection to patients with pheochromocytoma, since aggravated hypertension has been reported in association with diamorphine.

Cyclizine may cause a fall in cardiac output associated with increases in heart rate, mean arterial pressure and pulmonary wedge pressure. Cyclimorph Injection should therefore be used with caution in patients with severe heart failure.

Acute chest syndrome (ACS) in patients with sickle cell disease (SCD)

Due to a possible association between ACS and morphine use in SCD patients treated with Cyclimorph during a vaso-occlusive crisis, close monitoring for ACS symptoms is warranted.

Cyclizine should be avoided in patients with porphyria. Therefore, use of Cyclimorph Injection should also be avoided in these patients.

Case reports of paralysis have been received in patients using intravenous cyclizine. Some of the patients mentioned in these reports had an underlying neuromuscular disorder, thus intravenous cyclizine should be used with caution in all patients in general, and patients with underlying neuromuscular disorders in particular.

In common with other opioids, morphine may produce orthostatic hypotension and drowsiness in ambulatory patients. Sedation of short duration has been reported in patients receiving intravenous cyclizine. The CNS depressant effects of Cyclimorph Injection may be enhanced by combination with other centrally acting agents (see section 4.5). Patients should therefore be cautioned against activities requiring vigilance including driving vehicles and operating machinery.

Adrenal insufficiency

Opioid analgesics may cause reversible adrenal insufficiency requiring monitoring and glucocorticoid replacement therapy. Symptoms of adrenal insufficiency may include e.g. nausea, vomiting, loss of appetite, fatigue, weakness, dizziness, or low blood pressure.

Because cyclizine has anticholinergic activity it may precipitate incipient glaucoma. It should be used with caution and appropriate monitoring in patients with glaucoma and also in obstructive disease of the gastrointestinal tract.

Risk from concomitant use of sedative medicines such as benzodiazepines or related drugs:

Concomitant use of Cyclimorph 10 Solution for Injection and sedative medicines such as benzodiazepines or related drugs may result in sedation, respiratory depression, coma and death. Because of these risks, concomitant prescribing with these sedative medicines should be reserved for patients for whom alternative treatment options are not possible. If a decision is made to prescribe Cyclimorph 10 Solution for Injection concomitantly with sedative medicines, the lowest effective dose should be used, and the duration of treatment should be as short as possible.

The patients should be followed closely for signs and symptoms of respiratory depression and sedation. In this respect, it is strongly recommended to inform patients and their caregivers to be aware of these symptoms (see section 4.5).

Oral P2Y12 inhibitor antiplatelet therapy

Within the first day of concomitant P2Y12 inhibitor and morphine treatment, reduced efficacy of P2Y12 inhibitor treatment has been observed (see section 4.5).

Decreased Sex Hormones and increased prolactin

Long-term use of opioid analgesics may be associated with decreased sex hormone levels and increased prolactin. Symptoms include decreased libido, impotence or amenorrhea.

Hyperalgesia that does not respond to a further dose increase of morphine may occur in particular in high doses. A morphine dose reduction or change in opioid may be required.

Plasma concentrations of morphine may be reduced by rifampicin. The analgesic effect of morphine should be monitored and doses of morphine adjusted during and after treatment with rifampicin.

Cyclimorph 10 mg Injection contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per 1ml, that is to say essentially 'sodium-free'.

This medicine contains sodium metabisulphite which may rarely cause severe hypersensitivity reactions and bronchospasm.

4.5 Interaction with other medicinal products and other forms of interactions

The central nervous system depressant effects of Cyclimorph Injection may be enhanced by other centrally-acting agents such as phenothiazines, hypnotics, tranquillisers, anaesthetics, neuroleptics, alcohol and muscle relaxants.

Cyclizine enhances the soporific effect of pethidine.

Because of its anticholinergic activity cyclizine may enhance the side effects of other anticholinergic drugs, and may have an additive antimuscarinic action with other antimuscarinic drugs, such as atropine and some antidepressants (both tricyclics and MAOI's).

Mixed agonist/ antagonist opioid analgesics: Mixed agonist/antagonist opioid analgesics (e.g. pentazocine, nalbu phine, and buprenorphine) can reduce the analgesic effect of morphine by competitive blocking of the receptor. Therefore, these drugs should not be administered to patients who have received or are receiving a course of therapy with a pure opioid agonist analgesic.

Muscle relaxants: Morphine may enhance the neuromuscular blocking action of skeletal muscle relaxants.

The action of morphine may in turn affect the activities of other compounds, for example its gastrointestinal effects may delay absorption as with mexilitine or may be counteractive as with metoclopramide.

Monoamine oxidase inhibitors (MAOI's) may prolong and enhance the respiratory depressant effects of morphine. Opioids and MAOI's used together may cause fatal hypotension and coma (see section 4.3).

Cimetidine inhibits the metabolism of morphine.

The analgesic effect of opioids tends to be enhanced by co-administration of dexamphetamine, hydroxyzine and some phenothiazines although respiratory depression may also be enhanced by the latter combination.

Morphine may reduce the efficacy of diuretics by inducing the release of antidiuretic hormone. Morphine may also lead to acute retention of urine by causing spasm of the sphincter of the bladder, particularly in men with prostatism.

A delayed and decreased exposure to oral P2Y12 inhibitor antiplatelet therapy has been observed in patients with acute coronary syndrome treated with morphine. This interaction may be related to reduced gastrointestinal motility and apply to other opioids. The clinical relevance is unknown, but data indicate the potential for reduced P2Y12 inhibitor efficacy in patients co-administered morphine and a P2Y12 inhibitor (see section 4.4). In patients with acute coronary syndrome, in whom morphine cannot be withheld and fast P2Y12 inhibition is deemed crucial, the use of a parenteral P2Y12 inhibitor may be considered.

Propranolol has been reported to enhance the lethality of toxic doses of opioids in animals. Although the significance of this finding is not known for man, caution should be exercised when these drugs are administered concurrently.

In vitro data suggest that St. John's Wort (*Hypericum perforatum*) may induce cytochrome P450 3A4. There is a theoretical possibility therefore, that plasma levels of morphine tartrate may be decreased during concomitant administration and increased upon withdrawal of St. John's Wort.

Although there are no pharmacokinetic data available for concomitant use of ritonavir with morphine, ritonavir induces the hepatic enzymes responsible for the glucuronidation of morphine, and may possibly decrease plasma concentrations of morphine.

Dopaminergic: hyperpyrexia and CNS toxicity reported with selegiline.

Sedative medicines such as benzodiazepines or related drugs:

The concomitant use of opioids with sedative medicines such as benzodiazepines or related drugs increases the risk of sedation, respiratory depression, coma and death because of additive CNS depressant effect. The dose and duration of concomitant use should be limited (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy

There is no evidence on the safety of the combination in human pregnancy nor is there evidence from animal work that the constituents are free from hazard. However, limited data from epidemiological studies of cyclizine and morphine in human pregnancies have found no evidence of teratogenicity. In the absence of definitive human data with the combination the use of Cyclimorph Injection in pregnancy is not advised.

Newborns whose mothers received opioid analgesics during pregnancy should be monitored for signs of neonatal withdrawal (abstinence) syndrome. Treatment may include an opioid and supportive care.

Administration of morphine during labour may cause respiratory depression in the newborn infant.

Breast-feeding

Cyclizine is excreted in human milk; however, the amount has not been quantified.

Morphine can significantly suppress lactation. Morphine is excreted in human milk, but the amount is generally considered to be less than 1% of any dose.

Fertility

In a study involving prolonged administration of cyclizine to male and female rats, there was no evidence of impaired fertility after continuous treatment for 90-100 days at dose levels of approximately 15 and 25 mg/kg/day.

Effects of morphine exposure on sexual maturation of male rats, their reproductive capacity and the development of their progeny have been examined. Results indicated that exposure during adolescence led to pronounced inhibition of several indices of sexual maturation (e.g. hormone levels, reduced gonad weights), smaller litters and selective gender specific effects on endocrine function in the offspring.

Animal studies have shown that morphine may reduce fertility (see 5.3 preclinical safety data).

A disruption in ovulation and amenorrhoea can occur in women given morphine.

4.7 Effects on ability to drive and use machines

In common with other opioids, morphine may produce orthostatic hypotension and drowsiness in ambulatory patients. Sedation of short duration has been reported in patients receiving intravenous cyclizine. The CNS depressant effects of Cyclimorph Injection may be enhanced by combination with other centrally acting agents (see section 4.5). Patients should therefore be cautioned against activities requiring vigilance including driving vehicles and operating machinery.

4.8 Undesirable effects

Adverse reactions are listed below by system organ class and frequency. Frequencies are defined as: Very common: ($\geq 1/10$); Common ($\geq 1/100$ to $< 1/10$); Uncommon ($\geq 1/1,000$ to $< 1/100$); Rare ($\geq 1/10,000$ to $< 1/1,000$); Very rare ($< 1/10,000$); not known (cannot be estimated from the available data)

The following undesirable effects have been reported with a frequency of Not known:

Adverse reactions attributable to morphine include:

System Organ Class	Frequency	Adverse reactions
Blood and lymphatic system disorders	Not known	Thrombocytopenia
Immune system disorders	Not known	Anaphylactic shock, anaphylactoid reactions
Psychiatric disorders	Not known	Confusional state, dysphoria, restlessness, agitation, insomnia, euphoria, hallucinations, mood altered, dependence

Nervous system disorders	Not known	Intracranial pressure increased, headache, convulsions, paraesthesia, hyperesthesia/ allodynia, dizziness, syncope, coma, somnolence. A case of psychomotor hyperactivity following intravenous administration of morphine during induction of anaesthesia has been reported, allodynia, hyperalgesia (see section 4.4), hyperhidrosis
Eye disorders	Not known	Miosis, visual disturbance
Ear and labyrinth disorders	Not known	Vertigo
Cardiac disorders	Not known	Bradycardia, palpitations, tachycardia, heart failure
Vascular disorders	Not known	Orthostatic hypotension, facial flushing, hypertension, circulatory failure
Respiratory, thoracic and mediastinal disorders	Not known	Respiratory depression, bronchospasm, pulmonary oedema, respiratory failure
Gastrointestinal disorders	Not known	Constipation, nausea, vomiting, abdominal pain, anorexia, dry mouth, dyspepsia, narcotic bowel syndrome, ileus, taste perversion
Hepatobiliary disorders	Not known	Biliary colic, exacerbation of pancreatitis, increased hepatic enzymes
Skin and subcutaneous tissue disorders	Not known	Urticaria, hyperhidrosis, rash
Musculoskeletal and connective tissue disorders	Not known	Muscle spasm, hypertonia, myoclonus
Renal and urinary disorders	Not known	Dysuria, renal colic, urinary retention, ureteric spasm
Reproductive system and breast disorders	Not known	Morphine has a depressant effect on gonadal hormone secretion which can result in a reduction of testosterone leading to regression of secondary sexual characteristics in men on long-term therapy, amenorrhea, erectile dysfunction
General Disorders	Not known	Drug withdrawal (abstinence) syndrome, dysphoric mood, anxiety

Adverse reactions attributable to cyclizine include:

System Organ Class	Frequency	Adverse reactions
Blood and lymphatic system disorders	Not known	Agranulocytosis
Immune system disorders	Not known	Hypersensitivity, anaphylactic reaction, hypersensitivity hepatitis has occurred
Psychiatric disorders	Not known	Restlessness or agitation, nervousness, euphoria, insomnia, auditory and visual hallucinations (particularly when dosage recommendations have been exceeded), disorientation
Nervous system disorders	Not known	Case reports of paralysis have been received in patients using intravenous cyclizine. Some of the patients mentioned in these reports had an underlying neuromuscular disorder. Thus, intravenous cyclizine should be used with caution in all patients in general, and patients with underlying neuromuscular disorders in particular. Headache, somnolence, incoordination, sedation, dyskinesia, dystonia, extrapyramidal disorder, tremor, convulsions, dizziness, depressed level of consciousness, speech disorder, paraesthesia, generalised chorea
Eye disorders	Not known	Blurred vision, oculogyric crisis
Ear and labyrinth disorders	Not known	Tinnitus
Cardiac disorders	Not known	Tachycardia, palpitations, arrhythmias
Vascular disorders	Not known	Hypertension and hypotension
Respiratory, thoracic and mediastinal disorders	Not known	Nasal dryness, dry throat, bronchospasm and apnoea
Gastrointestinal disorders	Not known	Dryness of mouth, constipation
Hepatobiliary disorders	Not known	Hepatic function abnormal, jaundice cholestatic, hepatitis cholestatic, hepatitis
Skin and subcutaneous tissue disorders	Not known	Urticaria, pruritus, rash, angioedema, dermatitis allergic, fixed

		drug eruption
Musculoskeletal and connective tissue disorders	Not known	Twitching, muscle spasms
Renal and urinary disorders	Not known	Urinary retention

Rapid IV administration of cyclizine can lead to symptoms similar to overdose.

Adverse effects related to Injection formulation

System Organ Class	Adverse reactions
Ear and labyrinth disorders	Vertigo
Cardiac disorders	Bradycardia, palpitations
Vascular disorders	Thrombophlebitis
Gastrointestinal disorders	Constipation
Skin and subcutaneous tissue disorders	Urticaria, erythema, pruritus
General disorder and administration site condition	Injection site reaction, asthenia, malaise, hypothermia, pain

Cyclimorph IV Injection has demonstrated significant incidence of single cough or paroxysm of coughing immediately after its administration.

Drug dependence and withdrawal (abstinence) syndrome

Use of opioid analgesics may be associated with the development of physical and/or psychological dependence or tolerance. An abstinence syndrome may be precipitated when opioid administration is suddenly discontinued or opioid antagonists administered, or can sometimes be experienced between doses. For management, see 4.4.

Physiological withdrawal symptoms include: Body aches, tremors, restless legs syndrome, diarrhoea, abdominal colic, nausea, flu-like symptoms, tachycardia and mydriasis. Psychological symptoms include dysphoric mood, anxiety and irritability. In drug dependence, "drug craving" is often involved.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the HPRA Pharmacovigilance.

Website: www.hpra.ie.

4.9 Overdose

Symptoms

The signs of overdose with Cyclimorph Injection are those pathognomic of opioid poisoning i.e. respiratory depression, bradycardia, pin point pupils, hypotension, circulatory failure and deepening coma. Mydriasis may replace miosis as asphyxia intervenes. Opioid overdose can result in death. Death may occur from respiratory failure. Pneumonia aspiration.

Drowsiness, floppiness, miosis and apnoea are signs of opioid overdose in children as are convulsions.

Rhabdomyolysis progressing to renal failure has been reported in opioid overdose.

Signs and symptoms of acute toxicity from cyclizine arise from peripheral anticholinergic effects and effects on the central nervous system.

Peripheral anticholinergic symptoms include, dry mouth, nose and throat, blurred vision, tachycardia and urinary retention.

Central nervous system effects include drowsiness, dizziness, incoordination, ataxia, weakness, hyperexcitability, disorientation, impaired judgement, hallucinations, hyperkinesia, extrapyramidal motor disturbances, convulsions, hyperpyrexia and respiratory depression.

Management

It is imperative to maintain and support respiration and circulation.

The specific opioid antagonist naloxone is the treatment of choice for the reversal of coma and restoration of spontaneous respiration. The literature should be consulted for details of appropriate dosage.

The use of a specific opioid antagonist in patients tolerant to morphine may produce withdrawal symptoms.

Convulsions should be controlled with parenteral anticonvulsant therapy.

Patients should be monitored closely for at least 48 hours in case of relapse.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: piperazine derivatives.

ATC code: R06AE53

Mechanism of Action of cyclizine

Cyclizine is a histamine H₁ receptor antagonist of the piperazine class. It possesses anticholinergic and antiemetic properties. The exact mechanism by which cyclizine can prevent or suppress both nausea and vomiting from various causes is unknown.

Pharmacodynamic effects of cyclizine

Cyclizine increases lower oesophageal sphincter tone and reduces the sensitivity of the labyrinthine apparatus.

It may inhibit the part of the midbrain known collectively as the emetic centre.

Mechanism of action of Morphine

Morphine is a competitive agonist at the μ -opioid receptor and is a potent analgesic. It is thought that activity at the μ^1 -receptor subtype may mediate the analgesic and euphoric actions of morphine whilst activity at the μ^2 -receptor subtype may mediate respiratory depression and inhibition of gut motility.

Pharmacodynamic effects of Morphine

An action at the κ -opioid receptor may mediate spinal analgesia.

Clinical efficacy and safety

Cyclizine produces its anti-emetic effect within two hours and lasts approximately four hours.

5.2 Pharmacokinetic properties

Absorption of cyclizine

In a healthy adult volunteer, the administration of a single oral dose of 50 mg cyclizine resulted in a peak plasma concentration of approximately 70 ng/ml, occurring at about 2 hours after administration. Urine collected over 24 hours contained less than 1% of the total dose administered. In a separate study in one healthy adult volunteer the plasma elimination half-life of cyclizine was approximately 20 hours.

Biotransformation of cyclizine

Cyclizine is metabolised to its N-dimethylated derivative norcyclizine, which has little antihistaminic (H₁) activity compared to cyclizine.

Biotransformation of morphine

Morphine is extensively metabolised by hepatic biotransformation. In addition, the kidney has been shown to have the capacity to form morphine glucuronides. The major metabolite is morphine-3-glucuronide (approximately 45% of a dose). Morphine-6-glucuronide is a minor metabolite (approx. 5% of the dose) but is highly active. Although renal excretion is a minor route of elimination for unchanged morphine, it constitutes the major mechanism of elimination of conjugated morphine metabolites including the active morphine-6-glucuronide.

Distribution of morphine

Morphine is bound to plasma proteins only to the extent of 25-35% and therefore functions that change the extent of protein binding will have only a minor impact on its pharmacodynamic effects.

Elimination of morphine

The mean elimination half-life for morphine in blood and plasma is 2.7h (range 1.2-4.9h) and 2.95h (range 0.8-5h) respectively.

Interference with laboratory tests

Morphine can react with Folin-Ciocalteu reagent in the Lowry method of protein estimation.

Morphine can also interfere with the determination of urinary 17-ketosteroids due to chemical structure effects in the Zimmerman procedure.

5.3 Preclinical safety data

Mutagenicity

Cyclizine was not mutagenic in an Ames test (at a dose level of 100 µg/plate), with or without metabolic activation.

No bacterial mutagenicity studies with morphine have been reported. A review of the literature has indicated that morphine was negative in gene mutation assays in *Drosophila melanogaster*, but was positive in a mammalian spermatocyte test. The results of another study by the same authors has indicated that morphine causes chromosomal aberrations, in germ cells of male mice when given at dose levels of 10, 20, 40 or 60 mg/kg bodyweight for 3 consecutive days.

Carcinogenicity

No long-term studies have been conducted in animals to determine whether cyclizine or morphine are potentially carcinogenic.

Teratogenicity

Some animal studies indicate that cyclizine may be teratogenic at dose levels up to 25 times the clinical dose level. In another study, cyclizine was negative at oral dose levels up to 65 mg/kg in rats and 75 mg/kg in rabbits.

Morphine was not teratogenic in rats when dosed for up to 15 days at 70 mg/kg/day. Morphine given subcutaneously to mice at very high doses (200, 300 or 400 mg/kg/day) on days 8 or 9 of gestation, resulted in a few cases of exencephaly and axial skeletal fusions. The hypoxic effects of such high doses could account for the defects seen.

Lower doses of morphine (40, 4.0 or 0.4 mg/ml) given to mice as a continuous i.v. infusion (at a dose volume of 0.3 ml/kg) between days 7 and 10 of gestation, caused soft tissue and skeletal malformations as shown in previous studies.

Fertility

In male rats, reduced fertility and chromosomal damage in gametes have been reported.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tartaric acid
Sodium metabisulphite (E223)
Water for injections

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products. Physicochemical incompatibility (formation of precipitates) has been demonstrated between solutions of morphine sulphate and 5- fluorouracil.

6.3 Shelf life

Unopened: 3 years

Once Opened: From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user.

6.4 Special precautions for storage

Do not store above 25°C. Keep the ampoules in the outer carton in order to protect from light.
Do not freeze.

6.5 Nature and contents of container

Ampoules comply with the requirements of the European Pharmacopoeia for Type I neutral glass. Cyclimorph Injection is supplied in boxes of 5 x 1ml ampoules.

6.6 Special precautions for disposal and other handling

No special requirements for disposal.

For single use only. Discard any remaining solution.

7 MARKETING AUTHORISATION HOLDER

Amdipharm Limited
Temple Chambers
3 Burlington Road
Dublin 4
Ireland

8 MARKETING AUTHORISATION NUMBER

PA1142/003/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 01 April 1978

Date of last renewal: 01 April 2008

10 DATE OF REVISION OF THE TEXT

January 2021