

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Caspofungin 50mg powder for concentrate for solution for infusion

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each vial contains 50 mg caspofungin (as acetate). After reconstitution each ml concentrate for solution for infusion contains 5.2 mg caspofungin.

### Excipient with known effect

Sodium less than 1 mmol (23 mg) per vial.

For the full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Powder for concentrate for solution for infusion

White to off-white compact powder.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indications

- Treatment of invasive candidiasis in adult or paediatric patients.
- Treatment of invasive aspergillosis in adult or paediatric patients who are refractory to or intolerant of amphotericin B lipid formulations of amphotericin B and/or itraconazole. Refractoriness is defined as progression of infection or failure to improve after a minimum of 7 days of prior therapeutic doses of effective antifungal therapy.
- Empirical therapy for presumed fungal infections (such as *Candida* or *Aspergillus*) in febrile, neutropenic adult or paediatric patients.

### 4.2 Posology and method of administration

Caspofungin should be initiated by a physician experienced in the management of invasive fungal infections.

#### Posology

##### *Adult patients*

A single 70 mg loading dose should be administered on Day 1, followed by 50 mg daily thereafter. In patients weighing more than 80 kg, after the initial 70 mg loading dose, caspofungin 70 mg daily is recommended (see section 5.2). No dosage adjustment is necessary based on gender or race (see section 5.2).

##### *Paediatric patients (12 months to 17 years)*

In paediatric patients (12 months to 17 years of age), dosing should be based on the patient's body surface area (see Instructions for Use in Paediatric Patients, Mosteller<sup>1</sup> Formula). For all indications, a single 70-mg/m<sup>2</sup> loading dose (not to exceed an actual dose of 70 mg) should be administered on Day 1, followed by 50 mg/m<sup>2</sup> daily thereafter (not to exceed an actual dose of 70 mg daily). If the 50 mg/m<sup>2</sup> daily dose is well tolerated but does not provide an adequate clinical response, the daily dose can be increased to 70 mg/m<sup>2</sup> daily (not to exceed an actual daily dose of 70 mg).

The safety and efficacy of caspofungin have not been sufficiently studied in clinical trials involving neonates and infants below 12 months of age. Caution is advised when treating this age group. Limited data suggest that caspofungin at 25 mg/m<sup>2</sup> daily in neonates and infants (less than 3 months of age) and 50 mg/m<sup>2</sup> daily in young children (3 to 11 months of age) can be considered (see section 5.2).

**Duration of treatment**

Duration of empirical therapy should be based on the patient's clinical response. Therapy should be continued until up to 72 hours after resolution of neutropaenia (ANC  $\geq$  500). Patients found to have a fungal infection should be treated for a minimum of 14 days and treatment should continue for at least 7 days after both neutropaenia and clinical symptoms are resolved.

Duration of treatment of invasive candidiasis should be based upon the patient's clinical and microbiological response. After signs and symptoms of invasive candidiasis have improved and cultures have become negative, a switch to oral antifungal therapy may be considered. In general, antifungal therapy should continue for at least 14 days after the last positive culture.

Duration of treatment of invasive aspergillosis is determined on a case by case basis and should be based upon the severity of the patient's underlying disease, recovery from immunosuppression, and clinical response. In general, treatment should continue for at least 7 days after resolution of symptoms.

The safety information on treatment durations longer than 4 weeks is limited. However, available data suggest that caspofungin continues to be well tolerated with longer courses of therapy (up to 162 days in adult patients and up to 87 days in paediatric patients).

**Special populations***Elderly patients*

In elderly patients (65 years of age or more), the area under the curve (AUC) is increased by approximately 30 %. However, no systematic dosage adjustment is required. There is limited treatment experience in patients 65 years of age and older (see section 5.2).

*Renal impairment*

No dosage adjustment is necessary based on renal impairment (see section 5.2).

*Hepatic impairment*

For adult patients with mild hepatic impairment (Child-Pugh score 5 to 6), no dosage adjustment is needed. For adult patients with moderate hepatic impairment (Child-Pugh score 7 to 9), caspofungin 35 mg daily is recommended based upon pharmacokinetic data. An initial 70 mg loading dose should be administered on Day 1. There is no clinical experience in adult patients with severe hepatic impairment (Child-Pugh score greater than 9) and in paediatric patients with any degree of hepatic impairment (see section 4.4).

*Co-administration with inducers of metabolic enzymes*

Limited data suggest that an increase in the daily dose of caspofungin to 70 mg, following the 70 mg loading dose, should be considered when co-administering caspofungin in adult patients with certain inducers of metabolic enzymes (see section 4.5). When caspofungin is co-administered to paediatric patients (12 months to 17 years of age) with these same inducers of metabolic enzymes (see section 4.5), a caspofungin dose of 70-mg/m<sup>2</sup> daily (not to exceed an actual daily dose of 70 mg) should be considered.

Method of administration

After reconstitution and dilution, the solution should be administered by slow intravenous infusion over approximately 1 hour. For instructions on reconstitution and dilution of the medicinal product, see section 6.6.

Caspofungin is also available in vials with 70 mg caspofungin.

Caspofungin should be given as a single daily infusion.

<sup>1</sup> Mosteller RD. Simplified Calculation of Body Surface Area. N Engl J Med. 22 Oct 1987; N317(17): p.1098 (letter).

**4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

**4.4 Special warnings and precautions for use**

Anaphylaxis has been reported during administration of caspofungin. If this occurs, caspofungin should be discontinued and appropriate treatment administered. Possibly histamine-mediated adverse reactions, including rash, facial swelling, angioedema, pruritus, sensation of warmth, or bronchospasm have been reported and may require discontinuation and/or administration of appropriate treatment.

Limited data suggest that less common non- *Candida* yeasts and non- *Aspergillus* moulds are not covered by caspofungin. The efficacy of caspofungin against these fungal pathogens has not been established.

Concomitant use of caspofungin with cyclosporin has been evaluated in healthy adult volunteers and in adult patients. Some healthy adult volunteers who received two 3 mg/kg doses of cyclosporin with caspofungin showed transient increases in alanine transaminase (ALT) and aspartate transaminase (AST) of less than or equal to 3-fold the upper limit of normal (ULN) that resolved with discontinuation of the treatment. In a retrospective study of 40 patients treated during marketed use with caspofungin and cyclosporin for 1 to 290 days (median 17.5 days), no serious hepatic adverse reactions were noted. These data suggest that caspofungin can be used in patients receiving cyclosporin when the potential benefit outweighs the potential risk. Close monitoring of liver enzymes should be considered if caspofungin and cyclosporin are used concomitantly.

In adult patients with mild and moderate hepatic impairment, the AUC is increased about 20 % and 75 %, respectively. A reduction of the daily dose to 35 mg is recommended for adults with moderate hepatic impairment. There is no clinical experience in adults with severe hepatic impairment or in paediatric patients with any degree of hepatic impairment. A higher exposure than in moderate hepatic impairment is expected and caspofungin should be used with caution in these patients (see sections 4.2 and 5.2).

Laboratory abnormalities in liver function tests have been seen in healthy volunteers and adult and paediatric patients treated with caspofungin. In some adult and paediatric patients with serious underlying conditions who were receiving multiple concomitant medications with caspofungin, cases of clinically significant hepatic dysfunction, hepatitis and hepatic failure have been reported; a causal relationship to caspofungin has not been established. Patients who develop abnormal liver function tests during caspofungin therapy should be monitored for evidence of worsening hepatic function and the risk/benefit of continuing caspofungin therapy should be re-evaluated.

Cases of Stevens-Johnson Syndrome (SJS) and toxic epidermal necrolysis (TEN) have been reported after post-marketing use of caspofungin. Caution should apply in patients with history of allergic skin reaction (see section 4.8).

This medicinal product contains less than 1 mmol sodium (23 mg) per vial, i.e. essentially "sodium-free".

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Studies *in vitro* show that caspofungin is not an inhibitor of any enzyme in the cytochrome P450 (CYP) system. In clinical studies, caspofungin did not induce the CYP3A4 metabolism of other substances. Caspofungin is not a substrate for P-glycoprotein and is a poor substrate for cytochrome P450 enzymes. However, caspofungin has been shown to interact with other medicinal products in pharmacological and clinical studies (see below).

In two clinical studies performed in healthy adult subjects, cyclosporin A (one 4 mg/kg dose or two 3 mg/kg doses 12 hours apart) increased the AUC of caspofungin by approximately 35 %. These AUC increases are probably due to reduced uptake of caspofungin by the liver. Caspofungin did not increase the plasma levels of cyclosporin. There were transient increases in liver ALT and AST of less than or equal to 3-fold the upper limit of normal (ULN) when caspofungin and cyclosporin were co-administered, that resolved with discontinuation of the medicinal products. In a retrospective study of 40 patients treated during marketed use with caspofungin and cyclosporin for 1 to 290 days (median 17.5 days), no serious hepatic adverse reactions were noted (see section 4.4). Close monitoring of liver enzymes should be considered if the two medicinal products are used concomitantly.

Caspofungin reduced the trough concentration of tacrolimus by 26 % in healthy adult volunteers. For patients receiving both therapies, standard monitoring of tacrolimus blood concentrations and appropriate tacrolimus dosage adjustments are mandatory.

Clinical studies in healthy adult volunteers show that the pharmacokinetics of caspofungin are not altered to a clinically relevant extent by itraconazole, amphotericin B, mycophenolate, nelfinavir or tacrolimus. Caspofungin did not influence the pharmacokinetics of amphotericin B, itraconazole, rifampicin or mycophenolate mofetil. Although safety data are limited it

appears that no special precautions are needed when amphotericin B, itraconazole, nelfinavir or mycophenolate mofetil are co-administered with caspofungin.

Rifampicin caused a 60 % increase in AUC and 170 % increase in trough concentration of caspofungin on the first day of co-administration when both medicinal products were initiated together in healthy adult volunteers. Caspofungin trough levels gradually decreased upon repeated administration. After two weeks' administration rifampicin had limited effect on AUC, but trough levels were 30 % lower than in adult subjects who received caspofungin alone. The mechanism of interaction could possibly be due to an initial inhibition and subsequent induction of transport proteins. A similar effect could be expected for other medicinal products that induce metabolic enzymes. Limited data from population pharmacokinetics studies indicate that concomitant use of caspofungin with the inducers efavirenz, nevirapine, rifampicin, dexamethasone, phenytoin or carbamazepine may result in a decrease in caspofungin AUC. When co-administering inducers of metabolic enzymes, an increase in the daily dose of caspofungin to 70 mg, following the 70 mg loading dose, should be considered in adult patients (see section 4.2).

All adult drug-drug interaction studies described above were conducted at a 50 or 70 mg daily caspofungin dose. The interaction of higher doses of caspofungin with other medicinal products has not been formally studied.

In paediatric patients, results from regression analyses of pharmacokinetic data suggest that co-administration of dexamethasone with caspofungin may result in clinically meaningful reductions in caspofungin trough concentrations. This finding may indicate that paediatric patients will have similar reductions with inducers as seen in adults. When caspofungin is co-administered to paediatric patients (12 months to 17 years of age) with inducers of drug clearance, such as rifampicin, efavirenz, nevirapine, phenytoin, dexamethasone or carbamazepine, a caspofungin dose of 70 mg/m<sup>2</sup> daily (not to exceed an actual daily dose of 70 mg) should be considered.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

There are no or limited data from the use of caspofungin in pregnant women. Caspofungin should not be used during pregnancy unless clearly necessary. Animal studies have shown developmental toxicity (see section 5.3). Caspofungin has been shown to cross the placental barrier in animal studies.

##### Breastfeeding

It is unknown whether caspofungin is excreted in human milk. Available pharmacodynamic/toxicological data in animals have shown excretion of caspofungin in milk. Women receiving caspofungin should not breastfeed.

##### Fertility

For caspofungin, there were no effects on fertility in studies conducted in male and female rats (see section 5.3). There are no clinical data for caspofungin to assess its impact on fertility.

#### **4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive and use machines have been performed.

#### **4.8 Undesirable effects**

Hypersensitivity reactions (anaphylaxis and possibly histamine-mediated adverse reactions) have been reported (see section 4.4).

Also reported in patients with invasive aspergillosis were pulmonary oedema, adult respiratory distress syndrome (ARDS), and radiographic infiltrates.

##### Adult patients

In clinical studies, 1,865 adult individuals received single or multiple doses of caspofungin: 564 febrile neutropenic patients (empirical therapy study), 382 patients with invasive candidiasis, 228 patients with invasive aspergillosis, 297 patients with localised *Candida* infections, and 394 individuals enrolled in Phase I studies. In the empirical therapy study patients had received chemotherapy for malignancy or had undergone hematopoietic stem-cell transplantation (including 39 allogeneic

transplantations). In the studies involving patients with documented *Candida* infections, the majority of the patients with invasive *Candida* infections had serious underlying medical conditions (e.g., haematologic or other malignancy, recent major surgery, HIV) requiring multiple concomitant medications. Patients in the non-comparative *Aspergillus* study often had serious predisposing medical conditions (e.g., bone marrow or peripheral stem cell transplants, haematologic malignancy, solid tumours or organ transplants) requiring multiple concomitant medications.

Phlebitis was a commonly reported local injection-site adverse reaction in all patient populations. Other local reactions included erythema, pain/tenderness, itching, discharge and a burning sensation.

Reported clinical and laboratory abnormalities among all adults treated with caspofungin (total 1,780) were typically mild and rarely led to discontinuation.

#### Tabulated list of adverse reactions

The following adverse reactions were reported during clinical studies and/or post-marketing use:

<b>System Organ Class</b>	<b>Common (≥ 1/100 to &lt; 1/10)</b>	<b>Uncommon (≥ 1/1,000 to &lt; 1/100)</b>	<b>Not known (cannot be estimated from available data)</b>
Blood and lymphatic system disorders	haemoglobin decreased, haematocrit decreased, white blood cell count decreased	anaemia, thrombocytopenia, coagulopathy, leukopenia, eosinophil count increased, platelet count decreased, platelet count increased, lymphocyte count decreased, white blood cell count increased, neutrophil count decreased	
Metabolism and nutrition disorders	hypokalaemia	fluid overload, hypomagnesaemia, anorexia, electrolyte imbalance, hyperglycaemia, hypocalcaemia, metabolic acidosis	
Psychiatric disorders		anxiety, disorientation, insomnia	
Nervous system disorders	headache	dizziness, dysgeusia, paraesthesia, somnolence, tremor, hypoaesthesia	
Eye disorders		ocular icterus, vision blurred, eyelid oedema, lacrimation increased	
Cardiac disorders		palpitations, tachycardia, arrhythmia, atrial fibrillation, cardiac failure congestive	
Vascular disorders	phlebitis	thrombophlebitis, flushing, hot flush, hypertension, hypotension	
Respiratory, thoracic and mediastinal disorders	dyspnoea	nasal congestion, pharyngolaryngeal pain, tachypnoea, bronchospasm, cough, dyspnoea paroxysmal nocturnal, hypoxia, rales, wheezing	
Gastrointestinal disorders	nausea, diarrhoea, vomiting	abdominal pain, abdominal pain upper, dry mouth, dyspepsia, stomach discomfort, abdominal distension, ascites, constipation, dysphagia, flatulence	
Hepatobiliary disorders	elevated liver values (alanine aminotransferase, aspartate aminotransferase, blood alkaline phosphatase, bilirubin conjugated, blood bilirubin)	cholestasis, hepatomegaly, hyperbilirubinaemia, jaundice, hepatic function abnormal, hepatotoxicity, liver disorder, gamma-glutamyltransferase increased	
Skin and subcutaneous tissue disorders	rash, pruritus, erythema, hyperhidrosis	erythema multiforme, rash macular, rash maculo-papular, rash pruritic, urticaria, dermatitis allergic, pruritus generalised, rash erythematous, rash generalised, rash morbilliform, skin lesion	Toxic epidermal necrolysis and Stevens-Johnson syndrome (see section 4.4)
Musculoskeletal and	arthralgia	back pain, pain in extremity, bone pain, muscular	

connective tissue disorders		weakness, myalgia	
Renal and urinary disorders		renal failure, renal failure acute	
General disorders and administration site conditions	pyrexia, chills, infusion-site pruritus	pain, catheter site pain, fatigue, feeling cold, feeling hot, infusion site erythema, infusion site induration, infusion site pain, infusion site swelling, injection site phlebitis, oedema peripheral, tenderness, chest discomfort, chest pain, face oedema, feeling of body temperature change, induration, infusion site extravasation, infusion site irritation, infusion site phlebitis, infusion site rash, infusion site urticaria, injection site erythema, injection site oedema, injection site pain, injection site swelling, malaise, oedema	
Investigations	blood potassium decreased, blood albumin decreased	blood creatinine increased, red blood cells urine positive, protein total decreased, protein urine present, prothrombin time prolonged, prothrombin time shortened, blood sodium decreased, blood sodium increased, blood calcium decreased, blood calcium increased, blood chloride decreased, blood glucose increased, blood magnesium decreased, blood phosphorus decreased, blood phosphorus increased, blood urea increased, activated partial thromboplastin time prolonged, blood bicarbonate decreased, blood chloride increased, blood potassium increased, blood pressure increased, blood uric acid decreased, blood urine present, breath sounds abnormal, carbon dioxide decreased, immunosuppressant drug level increased, international normalised ratio increased, urinary casts, white blood cells urine positive, and pH urine increased.	

Caspofungin has also been evaluated at 150 mg daily (for up to 51 days) in 100 adult patients (see section 5.1). The study compared caspofungin at 50 mg daily (following a 70-mg loading dose on Day 1) versus 150 mg daily in the treatment of invasive candidiasis. In this group of patients, the safety of caspofungin at this higher dose appeared generally similar to patients receiving the 50 mg daily dose of caspofungin. The proportion of patients with a serious drug-related adverse reaction or a drug-related adverse reaction leading to caspofungin discontinuation was comparable in the 2 treatment groups.

#### Paediatric Patients

Data from 5 clinical studies completed in 171 paediatric patients suggest that the overall incidence of clinical adverse experiences (26.3 %; 95 % CI: -19.9, 33.6) is not worse than reported for adults treated with caspofungin (43.1 %; 95 % CI: -40.0, 46.2). However, paediatric patients probably have a different adverse event profile compared to adult patients. The most common drug-related clinical adverse experiences reported in paediatric patients treated with caspofungin were pyrexia (11.7 %), rash (4.7 %) and headache (2.9 %).

#### Tabulated list of adverse reactions

The following adverse reactions were reported:

<b>System Organ Class</b>	<b>Very common (≥ 1/10)</b>	<b>Common (≥ 1/100 to &lt; 1/10)</b>
Blood and lymphatic system disorders		eosinophil count increased
Nervous system disorders		headache
Cardiac disorders		tachycardia
Vascular disorders		flushing, hypotension
Hepatobiliary disorders		elevated liver enzyme levels (AST, ALT)
Skin and subcutaneous tissue disorders		rash, pruritus

General disorders and administration site conditions	fever	chills, catheter site pain
Investigations		decreased potassium, hypomagnesemia, increased glucose, decreased phosphorus, and increased phosphorus

#### Reporting of suspected adverse reactions

Reporting of suspected adverse reactions Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: [www.hpra.ie](http://www.hpra.ie); E-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie).

#### 4.9 Overdose

Inadvertent administration of up to 400 mg of caspofungin in one day has been reported. These occurrences did not result in clinically important adverse reactions. Caspofungin is not dialysable.

### 5 PHARMACOLOGICAL PROPERTIES

#### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: antimycotics for systemic use, ATC code: J02AX04 Mechanism of action

Caspofungin acetate is a semi-synthetic lipopeptide (echinocandin) compound synthesised from a fermentation product of *Glarea lozoyensis*. Caspofungin acetate inhibits the synthesis of beta (1,3)-D-glucan, an essential component of the cell wall of many filamentous fungi and yeast. Beta (1,3)-D-glucan is not present in mammalian cells.

Fungicidal activity with caspofungin has been demonstrated against *Candida* yeasts. Studies *in vitro* and *in vivo* demonstrate that exposure of *Aspergillus* to caspofungin results in lysis and death of hyphal apical tips and branch points where cell growth and division occur.

#### Pharmacodynamic effects

Caspofungin has *in vitro* activity against *Aspergillus* species (*Aspergillus fumigatus* [n = 75], *Aspergillus flavus* [n = 111], *Aspergillus niger* [n = 31], *Aspergillus nidulans* [n = 8], *Aspergillus terreus* [n = 52], and *Aspergillus candidus* [n = 3]). Caspofungin also has *in vitro* activity against *Candida* species (*Candida albicans* [n = 1,032], *Candida dubliniensis* [n = 100], *Candida glabrata* [n = 151], *Candida guilliermondii* [n = 67], *Candida kefyr* [n = 62], *Candida krusei* [n = 147], *Candida lipolytica* [n = 20], *Candida lusitanae* [n = 80], *Candida parapsilosis* [n = 215], *Candida rugosa* [n = 1] and *Candida tropicalis* [n = 258]), including isolates with multiple resistance transport mutations and those with acquired or intrinsic resistance to fluconazole, amphotericin B and 5-flucytosine. Susceptibility testing was performed according to a modification of both the Clinical and Laboratory Standards Institute (CLSI, formerly known as the National Committee for Clinical Laboratory Standards [NCCLS]) method M38-A2 (for *Aspergillus* species) and method M27-A3 (for

*Candida* species). Standardised techniques for susceptibility testing have been established for yeasts by EUCAST. EUCAST breakpoints have not yet been established for caspofungin, due to significant inter-laboratory variation in MIC ranges for caspofungin. In lieu of breakpoints, *Candida* isolates that are susceptible to anidulafungin as well as micafungin should be considered susceptible to caspofungin. Similarly, *C. parapsilosis* isolates intermediate to anidulafungin and micafungin can be regarded intermediate to caspofungin.

#### Mechanism of resistance

Isolates of *Candida* with reduced susceptibility to caspofungin have been identified in a small number of patients during treatment (MICs for caspofungin > 2 mg/L (4- to 30-fold MIC increases) have been reported using standardized MIC testing techniques approved by the CLSI). The mechanism of resistance identified is FKS1 and/or FKS2 (for *C. glabrata*) gene mutations. These cases have been associated with poor clinical outcomes.

Development of *in vitro* resistance to caspofungin by *Aspergillus* species has been identified. In limited clinical experience, resistance to caspofungin in patients with invasive aspergillosis has been observed. The mechanism of resistance has not been

established. The incidence of resistance to caspofungin by various clinical isolates of *Aspergillus* is rare. Caspofungin resistance in *Candida* has been observed but the incidence may differ by species or region.

#### Clinicalefficacyandsafety

##### *InvasiveCandidiasisinAdultPatients*

Two hundred thirty-nine patients were enrolled in an initial study to compare caspofungin and amphotericin B for the treatment of invasive candidiasis. Twenty-four patients had neutropaenia. The most frequent diagnoses were bloodstream infections (candidaemia) (77 %, n = 186) and *Candidaperitonitis* (8 %, n = 19); patients with *Candida* endocarditis, osteomyelitis or meningitis were excluded from this study. Caspofungin 50 mg once daily was administered following a 70-mg loading dose, while amphotericin B was administered at 0.6 to 0.7 mg/kg/day to non-neutropenic patients or 0.7 to 1.0 mg/kg/day to neutropenic patients. The mean duration of intravenous therapy was 11.9 days, with a range of 1 to 28 days. A favourable response required both symptom resolution and microbiological clearance of the *Candida* infection. Two hundred twenty-four patients were included in the primary efficacy analysis (MITT analysis) of response at the end of IV study therapy; favourable response rates for the treatment of invasive candidiasis were comparable for caspofungin (73 % [80/109]) and amphotericin B (62 % [71/115]) [% difference 12.7 (95.6 % CI: -0.7, 26.0)]. Among patients with candidaemia, favourable response rates at the end of IV study therapy were comparable for caspofungin (72 % [66/92]) and amphotericin B (63 % [59/94]) in the primary efficacy analysis (MITT analysis) [% difference 10.0 (95.0 % CI: -4.5, 24.5)]. Data in patients with non-blood sites of infection were more limited. Favourable response rates in neutropenic patients were 7/14 (50 %) in the caspofungin group and 4/10 (40 %) in the amphotericin B group. These limited data are supported by the outcome of the empirical therapy study.

In a second study, patients with invasive candidiasis received daily doses of caspofungin at 50 mg/day (following a 70-mg loading dose on Day 1) or caspofungin at 150 mg/day (see section 4.8). In this study, the caspofungin dose was administered over 2 hours (instead of the routine 1-hour administration). The study excluded patients with suspected *Candida* endocarditis, meningitis or osteomyelitis. As this was a primary therapy study, patients who were refractory to prior antifungal agents were also excluded. The number of neutropenic patients enrolled in this study was also limited (8.0 %). Efficacy was a secondary endpoint in this study. Patients who met the entry criteria and received one or more doses of caspofungin study therapy were included in the efficacy analysis. The favourable overall response rates at the end of caspofungin therapy were similar in the 2 treatment groups: 72 % (73/102) and 78 % (74/95) for the caspofungin 50-mg and 150-mg treatment groups, respectively (difference 6.3 % [95 % CI: -5.9, 18.4]).

##### *InvasiveAspergillosisinAdultPatients*

Sixty-nine adult patients (age 18-80) with invasive aspergillosis were enrolled in an open-label, non-comparative study to evaluate the safety, tolerability and efficacy of caspofungin. Patients had to be either refractory to (disease progression or failure to improve with other antifungal therapies given

for at least 7 days) (84 % of the enrolled patients) or intolerant of (16 % of enrolled patients) other standard antifungal therapies. Most patients had underlying conditions (haematologic malignancy [n = 24], allogeneic bone marrow transplant or stem cell transplant [n = 18], organ transplant [n = 8], solid tumour [n = 3] or other conditions [n = 10]). Stringent definitions, modelled after the Mycoses Study Group Criteria, were used for diagnosis of invasive aspergillosis and for response to therapy (favourable response required clinically significant improvement in radiographs as well as in signs and symptoms). The mean duration of therapy was 33.7 days, with a range of 1 to 162 days. An independent expert panel determined that 41 % (26/63) of patients receiving at least one dose of caspofungin had a favourable response. For those patients who received more than 7 days of therapy with caspofungin, 50 % (26/52) had a favourable response. The favourable response rates for patients who were either refractory to or intolerant of previous therapies were 36 % (19/53) and 70 % (7/10), respectively. Although the doses of prior antifungal therapies in 5 patients enrolled as refractory were lower than those often administered for invasive aspergillosis, the favourable response rate during therapy with caspofungin was similar in these patients to that seen in the remaining refractory patients (2/5 versus 17/48, respectively). The response rates among patients with pulmonary disease and extrapulmonary disease were 47 % (21/45) and 28 % (5/18), respectively. Among patients with extrapulmonary disease, 2 of 8 patients who also had definite, probable, or possible CNS involvement had a favourable response.

##### *EmpiricalTherapyinFebrile,NeutropenicAdultPatients*

A total of 1,111 patients with persistent fever and neutropaenia were enrolled in a clinical study and treated with either caspofungin 50 mg once daily following a 70-mg loading dose or liposomal amphotericin B 3.0 mg/kg/day. Eligible patients had received chemotherapy for malignancy or had undergone hematopoietic stem cell transplantation, and presented with neutropaenia (< 500 cells/mm<sup>3</sup> for 96 hours) and fever (> 38.0°C) not responding to ≥ 96 hours of parenteral antibacterial therapy.



Patients were to be treated until up to 72 hours after resolution of neutropaenia, with a maximum duration of 28 days. However, patients found to have a documented fungal infection could be treated longer. If the drug was well tolerated but the patient's fever persisted and clinical condition deteriorated after 5 days of therapy, the dosage of study drug could be increased to 70 mg/day of caspofungin (13.3 % of patients treated) or to 5.0 mg/kg/day of liposomal amphotericin B (14.3 % of patients treated). There were 1,095 patients included in the primary Modified Intention-To-Treat (MITT) efficacy analysis of overall favourable response; caspofungin (33.9 %) was as effective as liposomal amphotericin B (33.7 %) [% difference 0.2 (95.2% CI: -5.6, 6.0)]. An overall favourable response required meeting each of 5 criteria:

1. successful treatment of any baseline fungal infection (caspofungin 51.9 % [14/27], liposomal amphotericin B 25.9 % [7/27]),
2. no breakthrough fungal infections during administration of study drug or within 7 days after completion of treatment (caspofungin 94.8 % [527/556], liposomal amphotericin B 95.5 % [515/539]),
3. survival for 7 days after completion of study therapy (caspofungin 92.6 % [515/556], liposomal amphotericin B 89.2% [481/539]),
4. no discontinuation from the study drug because of drug-related toxicity or lack of efficacy (caspofungin 89.7 % [499/556], liposomal amphotericin B 85.5 % [461/539]), and
5. resolution of fever during the period of neutropaenia (caspofungin 41.2 % [229/556], liposomal amphotericin B 41.4 % [223/539]).

Response rates to caspofungin and liposomal amphotericin B for baseline infections caused by *Aspergillus* species were, respectively, 41.7 % (5/12) and 8.3 % (1/12), and by *Candida* species were 66.7 % (8/12) and 41.7 % (5/12). Patients in the caspofungin group experienced breakthrough infections due to the following uncommon yeasts and moulds: *Trichosporon* species (1), *Fusarium* species (1), *Mucor* species (1), and *Rhizopus* species (1).

#### *Paediatric population*

The safety and efficacy of caspofungin was evaluated in paediatric patients 3 months to 17 years of age in two prospective, multicentre clinical trials. The study design, diagnostic criteria and criteria for efficacy assessment were similar to the corresponding studies in adult patients (see section 5.1).

The first study, which enrolled 82 patients between 2 to 17 years of age, was a randomized, double-blind study comparing caspofungin (50 mg/m<sup>2</sup> IV once daily following a 70-mg/m<sup>2</sup> loading dose on Day 1 [not to exceed 70 mg daily]) to liposomal amphotericin B (3 mg/kg IV daily) in a 2:1 treatment fashion (56 on caspofungin, 26 on liposomal amphotericin B) as empirical therapy in paediatric patients with persistent fever and neutropenia. The overall success rates in the MITT analysis results, adjusted by risk strata, were as follows: 46.6 % (26/56) for caspofungin and 32.2 % (8/25) for liposomal amphotericin B.

The second study was a prospective, open-label, non-comparative study estimating the safety and efficacy of caspofungin in paediatric patients (ages 6 months to 17 years) with invasive candidiasis, oesophageal candidiasis and invasive aspergillosis (as salvage therapy). Forty-nine patients were enrolled and received caspofungin at 50 mg/m<sup>2</sup> IV once daily following a 70-mg/m<sup>2</sup> loading dose on Day 1 (not to exceed 70 mg daily), of whom 48 were included in the MITT analysis. Of these, 37 had invasive candidiasis, 10 had invasive aspergillosis and 1 patient had oesophageal candidiasis. The favourable response rate, by indication, at the end of caspofungin therapy was as follows in the MITT analysis: 81 % (30/37) in invasive candidiasis, 50 % (5/10) in invasive aspergillosis and 100 % (1/1) in oesophageal candidiasis.

In a double-blind, randomized (2:1) comparator-controlled study safety, tolerability and efficacy of caspofungin (2 mg/kg/d intravenously, infused over 2 hours) vs amphotericin B deoxycholate (1 mg/kg/d) was evaluated in neonates and infants less than 3 months of age with (culture-confirmed) invasive candidiasis. Due to poor enrolment, the study was terminated early and only 51 patients were randomized. The proportion of patients with fungal-free survival at 2 weeks post-therapy in the caspofungin treatment group (71.0 %) was similar to that seen in the amphotericin B deoxycholate treatment group (68.8 %). Based on this study, no posology recommendations for neonates and infants can be made.

## **5.2 Pharmacokinetic properties**

### Distribution

Caspofungin is extensively bound to albumin. The unbound fraction of caspofungin in plasma varies from 3.5 % in healthy volunteers to 7.6 % in patients with invasive candidiasis. Distribution plays the prominent role in caspofungin plasma pharmacokinetics and is the rate-controlling step in both the alpha- and beta-disposition phases. The distribution into tissues

peaked at 1.5 to 2 days after dosing when 92 % of the dose was distributed into tissues. It is likely that only a small fraction of the caspofungin taken up into tissues later returns to plasma as parent compound. Therefore, elimination occurs in the absence of a distribution equilibrium, and a true estimate of the volume of distribution of caspofungin is currently impossible to obtain.

#### Biotransformation

Caspofungin undergoes spontaneous degradation to an open ring compound. Further metabolism involves peptide hydrolysis and N-acetylation. Two intermediate products, formed during the degradation of caspofungin to this open ring compound, form covalent adducts to plasma proteins resulting in a low-level, irreversible binding to plasma proteins.

*In vitro* studies show that caspofungin is not an inhibitor of cytochrome P450 enzymes 1A2, 2A6, 2C9, 2C19, 2D6 or 3A4. In clinical studies, caspofungin did not induce or inhibit the CYP3A4 metabolism of other medicinal products. Caspofungin is not a substrate for P-glycoprotein and is a poor substrate for cytochrome P450 enzymes.

#### Elimination

The elimination of caspofungin from plasma is slow with a clearance of 10-12 ml/min. Plasma concentrations of caspofungin decline in a polyphasic manner following single 1-hour intravenous infusions. A short alpha-phase occurs immediately post-infusion, followed by a beta-phase with a half-life of 9 to 11 hours. An additional gamma-phase also occurs with a half-life of 45 hours. Distribution, rather than excretion or biotransformation, is the dominant mechanism influencing plasma clearance.

Approximately 75 % of a radioactive dose was recovered during 27 days: 41 % in urine and 34 % in faeces. There is little excretion or biotransformation of caspofungin during the first 30 hours after administration. Excretion is slow and the terminal half-life of radioactivity was 12 to 15 days. A small amount of caspofungin is excreted unchanged in urine (approximately 1.4 % of dose).

Caspofungin displays moderate non-linear pharmacokinetics with increased accumulation as the dose is increased, and a dose dependency in the time to reach steady state upon multiple-dose administration.

#### Special populations

Increased caspofungin exposure was seen in adult patients with renal impairment and mild liver impairment, in female subjects and in the elderly. Generally the increase was modest and not large enough to warrant dosage adjustment. In adult patients with moderate liver impairment or in higher weight patients, a dosage adjustment may be necessary (see below).

#### *Weight*

Weight was found to influence caspofungin pharmacokinetics in the population pharmacokinetic analysis in adult candidiasis patients. The plasma concentrations decrease with increasing weight. The average exposure in an adult patient weighing 80 kg was predicted to be about 23 % lower than in an adult patient weighing 60 kg (see section 4.2).

#### *Hepatic impairment*

In adult patients with mild and moderate hepatic impairment, the AUC is increased about 20 and 75 %, respectively. There is no clinical experience in adult patients with severe hepatic impairment and in paediatric patients with any degree of hepatic impairment. In a multiple-dose study, a dose reduction of the daily dose to 35 mg in adult patients with moderate hepatic impairment has been shown to provide an AUC similar to that obtained in adult subjects with normal hepatic function receiving the standard regimen (see section 4.2).

#### *Renal impairment*

In a clinical study of single 70 mg doses, caspofungin pharmacokinetics were similar in adult volunteers with mild renal impairment (creatinine clearance 50 to 80 ml/min) and control subjects. Moderate (creatinine clearance 31 to 49 ml/min), advanced (creatinine clearance 5 to 30 ml/min) and end-stage (creatinine clearance <10 ml/min and dialysis dependent) renal impairment moderately increased caspofungin plasma concentrations after single-dose administration (range: 30 to 49 % for AUC). However, in adult patients with invasive candidiasis, oesophageal candidiasis or invasive aspergillosis who received multiple daily doses of caspofungin 50 mg, there was no significant effect of mild to advanced renal impairment on caspofungin concentrations. No dosage adjustment is necessary for patients with renal impairment. Caspofungin is not dialysable, thus supplementary dosing is not required following haemodialysis.

#### *Gender*

Caspofungin plasma concentrations were on average 17 - 38 % higher in women than in men.

#### *Elderly*

A modest increase in AUC (28 %) and  $C_{24h}$  (32 %) was observed in elderly male subjects compared with young male subjects. In patients who were treated empirically or who had invasive candidiasis, a similar modest effect of age was seen in older patients relative to younger patients.

#### Race

Patient pharmacokinetic data indicated that no clinically significant differences in the pharmacokinetics of caspofungin were seen among Caucasians, Blacks, Hispanics and Mestizos.

#### Paediatric Patients

In adolescents (ages 12 to 17 years) receiving caspofungin at 50 mg/m<sup>2</sup> daily (maximum 70 mg daily), the caspofungin plasma AUC<sub>0-24 hr</sub> was generally comparable to that seen in adults receiving caspofungin at 50 mg daily. All adolescents received doses > 50 mg daily, and, in fact, 6 of 8 received the maximum dose of 70 mg/day. The caspofungin plasma concentrations in these adolescents were reduced relative to adults receiving 70 mg daily, the dose most often administered to adolescents.

In children (ages 2 to 11 years) receiving caspofungin at 50 mg/m<sup>2</sup> daily (maximum 70 mg daily), the caspofungin plasma AUC<sub>0-24 hr</sub> after multiple doses was comparable to that seen in adults receiving caspofungin at 50 mg/day.

In young children and toddlers (ages 12 to 23 months) receiving caspofungin at 50 mg/m<sup>2</sup> daily (maximum 70 mg daily), the caspofungin plasma AUC<sub>0-24 hr</sub> after multiple doses was comparable to that seen in adults receiving caspofungin at 50 mg daily and to that in older children (2 to 11 years of age) receiving the 50 mg/m<sup>2</sup> daily dose.

Overall, the available pharmacokinetic, efficacy and safety data are limited in patients 3 to 10 months of age. Pharmacokinetic data from one 10-month old child receiving the 50 mg/m<sup>2</sup> daily dose indicated an AUC<sub>0-24 hr</sub> within the same range as that observed in older children and adults at the 50 mg/m<sup>2</sup> and the 50 mg dose, respectively, while in one 6-month old child receiving the 50 mg/m<sup>2</sup> dose, the AUC<sub>0-24 hr</sub> was somewhat higher.

In neonates and infants (< 3 months) receiving caspofungin at 25 mg/m<sup>2</sup> daily (corresponding mean daily dose of 2.1 mg/kg), caspofungin peak concentration ( $C_{1 hr}$ ) and caspofungin trough concentration ( $C_{24 hr}$ ) after multiple doses were comparable to that seen in adults receiving caspofungin at 50 mg daily. On Day 1,  $C_{1 hr}$  was comparable and  $C_{24 hr}$  modestly elevated (36 %) in these neonates and infants relative to adults. However, variability was seen in both  $C_{1 hr}$  (Day 4 geometric mean 11.73 µg/mL, range 2.63 to 22.05 µg/mL) and  $C_{24 hr}$  (Day 4 geometric mean 3.55 µg/mL, range 0.13 to 7.17 µg/mL). AUC<sub>0-24</sub> measurements were not performed in this study due to the sparse plasma sampling. Of note, the efficacy and safety of caspofungin have not been adequately studied in prospective clinical trials involving neonates and infants under 3 months of age.

### 5.3 Preclinical safety data

Repeated dose toxicity studies in rats and monkeys using doses up to 7-8 mg/kg given intravenously showed injection site reactions in rats and monkeys, signs of histamine release in rats and evidence of adverse effects directed at the liver in monkeys. Developmental toxicity studies in rats showed that caspofungin caused decreases in foetal body weights and an increase in the incidence of incomplete ossification of vertebra, sternebra and skull bone at doses of 5 mg/kg that were coupled to adverse maternal effects such as signs of histamine release in pregnant rats. An increase in the incidence of cervical ribs was also noted. Caspofungin was negative in *in vitro* assays for potential genotoxicity as well as in the *in vivo* mouse bone marrow chromosomal test. No long-term studies in animals have been performed to evaluate the carcinogenic potential. For caspofungin, there were no effects on fertility in studies conducted in male and female rats up to 5 mg/kg/day.

## 6 PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Sucrose  
Mannitol  
Glacial acetic acid  
Sodium hydroxide 3.9 % (to adjust the pH)

### 6.2 Incompatibilities

Do not mix with diluents containing glucose, as Caspofungin is not stable in diluents containing glucose. In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products. <?xml:namespace prefix = o ns = urn:schemas-microsoft-com:office:office />

### 6.3 Shelf life

2 years

The following in-use storage times of the reconstituted concentrate for solution for infusion and the diluted solution for infusion are not additive.

#### *Reconstituted concentrate for solution for infusion*

Chemical and physical in-use stability has been demonstrated for 24 hours at  $\leq 25$  °C. From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8 °C, unless reconstitution has taken place in controlled and validated aseptic conditions.

Do not freeze the reconstituted concentrate for solution for infusion.

#### *Diluted solution for infusion*

Chemical and physical in-use stability has been demonstrated for 24 hours at  $\leq 25$  °C and for 48 hours at 2 to 8 °C. From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8 °C, unless dilution has taken place in controlled and validated aseptic conditions.

Do not freeze the reconstituted diluted solution for infusion.

### 6.4 Special precautions for storage

Store in a refrigerator (2 °C - 8 °C).

For storage conditions after reconstitution and dilution of the medicinal product, see section 6.3.

### 6.5 Nature and contents of container

10 ml type I glass vial with a grey bromobutyl rubber stopper and a flip-off seal with plastic cap. Supplied in packs of 1 vial.

### 6.6 Special precautions for disposal and other handling

#### **INSTRUCTIONS FOR USE IN ADULT PATIENTS**

##### **Step 1 Reconstitution of conventional vials**

To reconstitute the powder, bring the vial to room temperature and aseptically add 10.5 ml of water for injection.

The white to off-white compact lyophilised powder will dissolve completely. Mix gently until a clear solution is obtained. Reconstituted solutions should be visually inspected for particulate matter or discolouration.

The concentrations of the reconstituted vials will be 5.2 mg/ml.

##### **Step 2 Addition of reconstituted Caspofungin to patient infusion solution**

Diluents for the final solution for infusion are: sodium chloride solution for injection 9 mg/ml (0.9 %), or lactated Ringer's solution.

The solution for infusion is prepared by aseptically adding the appropriate amount of reconstituted concentrate for solution for infusion (as shown in the table below) to a 250 ml infusion bag or bottle.

Reduced volume infusions in 100 ml may be used, when medically necessary, for 50 mg or 35 mg daily doses.

Visually inspect the infusion solution for particulate matter or discolouration. Do not use if the solution is cloudy or has precipitated.

## PREPARATION OF THE SOLUTION FOR INFUSION IN ADULTS

DOSE*	Volume of reconstituted Caspofungin for transfer to intravenous bag or bottle	Standard preparation final concentration (reconstituted Caspofungin added to 250 ml diluent)	Reduced volume infusion final concentration (reconstituted Caspofungin added to 100 ml diluent)
50 mg	10 ml	0.20 mg/ml	-
50 mg at reduced volume	10 ml	-	0.47 mg/ml
35 mg for moderate hepatic impairment (from one 50 mg vial)	7 ml	0.14 mg/ml	-
35 mg for moderate hepatic impairment (from one 50 mg vial) at reduced volume	7 ml	-	0.34 mg/ml

\* 10.5 ml should be used for reconstitution of all vials

## INSTRUCTIONS FOR USE IN PAEDIATRIC PATIENTS

### Calculation of Body Surface Area (BSA) for paediatric dosing

Before preparation of infusion, calculate the body surface area (BSA) of the patient using the following formula (Mosteller Formula):

### Preparation of the 70 mg/m<sup>2</sup> infusion for paediatric patients >3 months of age (using a 50-mg vial)

1. Determine the actual loading dose to be used in the paediatric patient by using the patient's BSA (as calculated above) and the following equation:

$$\text{BSA (m}^2\text{)} \times 70 \text{ mg/m}^2 = \text{Loading Dose}$$

The maximum loading dose on Day 1 should not exceed 70 mg regardless of the patient's calculated dose.

2. Equilibrate the refrigerated vial of Caspofungin to room temperature.
3. Aseptically add 10.5 ml of water for injection.<sup>a</sup> This will give a final caspofungin concentration in the vial of 5.2 mg/ml.
4. Remove the volume of medicinal product equal to the calculated loading dose (step 1) from the vial. Aseptically transfer this volume (ml)<sup>b</sup> of reconstituted Caspofungin to an IV bag (or bottle) containing 250 ml of sodium chloride solution for injection 9 mg/ml (0.9 %), or lactated Ringer's solution. Alternatively, the volume (ml)<sup>b</sup> of reconstituted Caspofungin can be added to a reduced volume of sodium chloride solution for injection 9 mg/ml (0.9 %) or lactated Ringer's solution, not to exceed a final concentration of 0.5 mg/ml.

### Preparation of the 50 mg/m<sup>2</sup> infusion for paediatric patients >3 months of age (using a 50-mg vial)

1. Determine the actual daily maintenance dose to be used in the paediatric patient by using the patient's BSA (as calculated above) and the following equation:

$$\text{BSA (m}^2\text{)} \times 50 \text{ mg/m}^2 = \text{Daily Maintenance Dose}$$

The daily maintenance dose should not exceed 70 mg regardless of the patient's calculated dose.

2. Equilibrate the refrigerated vial of Caspofungin to room temperature.
3. Aseptically add 10.5 ml of water for injection.<sup>a</sup> This will give a final caspofungin concentration in the vial of 5.2 mg/ml.
4. Remove the volume of medicinal product equal to the calculated daily maintenance dose (step 1) from the vial. Aseptically transfer this volume (ml)<sup>b</sup> of reconstituted caspofungin to an IV bag (or bottle) containing 250 ml of sodium chloride solution for injection 9 mg/ml (0.9 %), or lactated Ringer's solution. Alternatively, the volume (ml)<sup>b</sup> of reconstituted Caspofungin can be added to a reduced volume of sodium chloride solution for injection 9 mg/ml (0.9 %) or lactated Ringer's solution, not to exceed a final concentration of 0.5 mg/ml.

### Preparation notes:

- a. The white to off-white cake will dissolve completely. Mix gently until a clear solution is obtained.
- b. Caspofungin is formulated to provide the full labelled vial dose (50 mg) when 10 ml is withdrawn from the vial.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

**7 MARKETING AUTHORISATION HOLDER**

Flynn Pharma Limited  
5th Floor  
40 Mespil Road  
Dublin 4  
D04 C2N4  
Ireland

**8 MARKETING AUTHORISATION NUMBER**

PA1226/013/001

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 13<sup>th</sup> January 2017

Date of last renewal: 25<sup>th</sup> November 2021

**10 DATE OF REVISION OF THE TEXT**

February 2023