

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Co-amoxiclav Bluefish 875 mg/125 mg film- coated tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film coated tablet contains 1006.25 mg amoxicillin trihydrate equivalent to 875 mg amoxicillin and potassium clavulanate equivalent to 125 mg - clavulanic acid.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Film coated tablet.

A white, elongated capsule shaped film coated tablets with breakline on both sides..

The scoreline is only to facilitate breaking for ease of swallowing and not to divide in to equal doses.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Amoxicillin-clavulanic acid is indicated for the treatment of the following infections in adults and children (see sections 4.2, 4.4 and 5.1):

- Acute bacterial sinusitis (adequately diagnosed)
- Acute otitis media
- Acute exacerbations of chronic bronchitis (adequately diagnosed)
- Community-acquired pneumonia
- Cystitis
- Pyelonephritis
- Skin and soft tissue infections in particular cellulitis, animal bites and severe dental abscess with spreading cellulitis;
- Bone and joint infections, in particular osteomyelitis

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Posology

Dosages are expressed throughout in terms of amoxicillin-/clavulanic acid content except when doses are stated in terms of an individual component.

The dose of amoxicillin-/clavulanic acid that is selected to treat an individual infection should take into account:

- The expected pathogens and their likely susceptibility to antibacterial agents (see section 4.4)
- The severity and the site of the infection
- The age, weight and renal function of the patient as shown below.

The use of alternative presentations of amoxicillin-/clavulanic acid (e.g. those that provide higher doses of amoxicillin and/or different ratios of amoxicillin to clavulanic acid) should be considered as necessary (see sections 4.4 and 5.1).

For adults and children ≥ 40 kg, this formulation of amoxicillin-/clavulanic acid provides a total daily dose of 1750 mg amoxicillin/ 250 mg clavulanic acid with twice daily dosing and 2625 mg amoxicillin/375 mg clavulanic acid with three times daily dosing, when administered as recommended below. For children < 40 kg, this formulation of amoxicillin-/clavulanic acid provides a maximum daily dose of 1000-2800 mg amoxicillin/143-400 mg clavulanic acid, when administered as recommended below. If it is considered that a higher daily dose of amoxicillin is required, it is recommended that another preparation of amoxicillin-/clavulanic acid is selected in order to avoid administration of unnecessarily high daily doses of clavulanic acid (see sections 4.4 and 5.1).

The duration of therapy should be determined by the response of the patient. Some infections (e.g. osteomyelitis) require longer periods of treatment. Treatment should not be extended beyond 14 days without review (see section 4.4 regarding prolonged therapy).

Adults and children \geq 40 kg

Recommended doses:

- standard dose: (for all indications) 875 mg/125 mg two times a day;
- higher dose - (particularly for infections such as otitis media, sinusitis, lower respiratory tract infections and urinary tract infections): 875 mg/125 mg three times a day.

Children < 40 kg

Children may be treated with amoxicillin-/clavulanic acid tablets, suspensions or paediatric sachets.

Recommended doses:

- 25 mg/3.6 mg/kg/day to 45 mg/6.4 mg/kg/day given as two divided doses;
- up to 70 mg/10 mg/kg/day given as two divided doses may be considered for some infections (such as otitis media, sinusitis and lower respiratory tract infections).

No clinical data are available for amoxicillin-/clavulanic acid 7:1 formulations regarding doses higher than 45 mg/6.4 mg per kg per day in children under 2 years

There are no clinical data for amoxicillin-/clavulanic acid 7:1 formulations for patients under 2 months of age. Dosing recommendations in this population therefore cannot be made.

Elderly

No dose adjustment is considered necessary.

Renal impairment

No dose adjustment is required in patients with creatinine clearance (CrCl) greater than 30 ml/min.

In patients with creatinine clearance less than 30 ml/min, the use of amoxicillin-/clavulanic acid presentations with an amoxicillin to clavulanic acid ratio of 7:1 is not recommended, as no recommendations for dose adjustments are available.

Hepatic impairment

Dose with caution and monitor hepatic function at regular intervals (see sections 4.3 and 4.4).

Method of administration

Amoxicillin/clavulanic acid tablet is for oral use.

Administer at the start of a meal to minimise potential gastrointestinal intolerance and optimise absorption of amoxicillin/clavulanic acid.

4.3 Contraindications

Hypersensitivity to the active substances, to any of the penicillins or to any of the excipients.

History of a severe immediate hypersensitivity reaction (e.g. anaphylaxis) to another beta-lactam agent (e.g. a cephalosporin, carbapenem or monobactam).

History of jaundice/hepatic impairment due to amoxicillin/clavulanic acid (see section 4.8).

4.4 Special warnings and precautions for use

Before initiating therapy with amoxicillin/clavulanic acid, careful enquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins or other beta-lactam agents (see sections 4.3 and 4.8).

Serious and occasionally fatal hypersensitivity (including anaphylactoid and severe cutaneous adverse reactions) reactions have been reported in patients on penicillin therapy. Hypersensitivity reactions can also progress to Kounis syndrome, a serious allergic reaction that can result in myocardial infarction (see section 4.8). These reactions are more likely to occur in individuals

with a history of penicillin hypersensitivity and in atopic individuals. If an allergic reaction occurs, amoxicillin/clavulanic acid therapy must be discontinued and appropriate alternative therapy instituted.

Convulsions may occur in patients with impaired renal function or in those receiving high doses (see 4.8).

Amoxicillin/clavulanic acid should be avoided if infectious mononucleosis is suspected since the occurrence of a morbilliform rash has been associated with this condition following the use of amoxicillin.

Concomitant use of allopurinol during treatment with amoxicillin can increase the likelihood of allergic skin reactions.

Prolonged use may also occasionally result in overgrowth of non-susceptible organisms.

The occurrence at the treatment initiation of a feverish generalised erythema associated with pustula may be a symptom of acute generalised exanthemous pustulosis (AGEP) (see Section 4.8). This reaction requires amoxicillin/clavulanic acid discontinuation and contra-indicates any subsequent administration of amoxicillin.

Amoxicillin/clavulanic acid should be used with caution in patients with evidence of hepatic impairment (see sections 4.2, 4.3 and 4.8).

Hepatic events have been reported predominantly in males and elderly patients and may be associated with prolonged treatment. These events have been very rarely reported in children. In all populations, signs and symptoms usually occur during or shortly after treatment but in some cases may not become apparent until several weeks after treatment has ceased. These are usually reversible. Hepatic events may be severe and in extremely rare circumstances, deaths have been reported. These have almost always occurred in patients with serious underlying disease or taking concomitant medications known to have the potential for hepatic effects (see section 4.8).

Antibiotic-associated colitis has been reported with nearly all antibacterial agents including amoxicillin and may range in severity from mild to life threatening (see section 4.8). Therefore, it is important to consider this diagnosis in patients who present with diarrhoea during or subsequent to the administration of any antibiotics. Should antibiotic-associated colitis occur, amoxicillin/clavulanic acid should immediately be discontinued, a physician be consulted and an appropriate therapy initiated. Anti-peristaltic drugs are contra-indicated in this situation.

Periodic assessment of organ system functions, including renal, hepatic and haematopoietic function is advisable during prolonged therapy.

Drug-induced enterocolitis syndrome (DIES) has been reported mainly in children receiving amoxicillin/clavulanate (see section 4.8). DIES is an allergic reaction with the leading symptom of protracted vomiting (1-4 hours after drug intake) in the absence of allergic skin or respiratory symptoms. Further symptoms could comprise abdominal pain, diarrhoea, hypotension or leucocytosis with neutrophilia. There have been severe cases including progression to shock.

Prolongation of prothrombin time has been reported rarely in patients receiving amoxicillin/clavulanic acid. Appropriate monitoring should be undertaken when anticoagulants are prescribed concomitantly. Adjustments in the dose of oral anticoagulants may be necessary to maintain the desired level of anticoagulation (see section 4.5 and 4.8).

In patients with renal impairment, dosage should be adjusted according to the degree of impairment (see section 4.2).

In patients with reduced urine output, crystalluria (including acute renal injury) has been observed very rarely, predominantly with parenteral therapy. During the administration of high doses of amoxicillin, it is advisable to maintain adequate fluid intake and urinary output in order to reduce the possibility of amoxicillin crystalluria. In patients with bladder catheters, a regular check of patency should be maintained (see section 4.8 and 4.9).

During treatment with amoxicillin, enzymatic glucose oxidase methods should be used whenever testing for the presence of glucose in urine because false positive results may occur with non-enzymatic methods.

The presence of clavulanic acid in tablet may cause a non-specific binding of IgG and albumin by red cell membranes leading to a false positive Coombs test.

There have been reports of positive test results using the Bio-Rad Laboratories Platelia *Aspergillus* EIA test in patients receiving amoxicillin/clavulanic acid who were subsequently found to be free of *Aspergillus* infection. Cross-reactions with

non-*Aspergillus* polysaccharides and polyfuranoses with Bio-Rad Laboratories Platelia *Aspergillus* EIA test have been reported. Therefore, positive test results in patients receiving amoxicillin/clavulanic acid should be interpreted cautiously and confirmed by other diagnostic methods.

This medicine contains less than 1 mmol sodium (23 mg) per 875mg/125mg co-amoxiclav tablet, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Oral anticoagulants

Oral anticoagulants and penicillin antibiotics have been widely used in practice without reports of interaction. However, in the literature there are cases of increased international normalised ratio in patients maintained on acenocoumarol or warfarin and prescribed a course of amoxicillin. If co-administration is necessary, the prothrombin time or international normalised ratio should be carefully monitored with the addition or withdrawal of amoxicillin. Moreover, adjustments in the dose of oral anticoagulants may be necessary (see sections 4.4 and 4.8).

Methotrexate

Penicillins may reduce the excretion of methotrexate causing a potential increase in toxicity.

Probenecid

Concomitant use of probenecid is not recommended. Probenecid decreases the renal tubular secretion of amoxicillin. Concomitant use of probenecid may result in increased and prolonged blood levels of amoxicillin, but not of clavulanic acid.

4.6 Fertility, pregnancy and lactation

Pregnancy

Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development (see section 5.3). Limited data on the use of amoxicillin/clavulanic acid during pregnancy in humans do not indicate an increased risk of congenital malformations. In a single study in women with preterm, premature rupture of the foetal membrane, it was reported that prophylactic treatment with amoxicillin-clavulanic acid may be associated with an increased risk of necrotising enterocolitis in neonates. Use should be avoided during pregnancy, unless considered essential by the physician.

Brest-feeding

Both substances are excreted into breast milk (nothing is known of the effects of clavulanic acid on the breast-fed infant). Consequently, diarrhoea and fungus infection of the mucous membranes are possible in the breast-fed infant, so that breast-feeding might have to be discontinued. Amoxicillin/clavulanic acid should only be used during breast-feeding after benefit/risk assessment by the physician in charge.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. However, undesirable effects may occur (e.g. allergic reactions, dizziness, convulsions), which may influence the ability to drive and use machines (see section 4.8).

4.8 Undesirable effects

The most commonly reported adverse drug reactions (ADRs) are diarrhoea, nausea and vomiting.

The ADRs derived from clinical studies and post-marketing surveillance with amoxicillin/clavulanic acid, sorted by MedDRA System Organ Class are listed below.

The following terminologies have been used in order to classify the occurrence of undesirable effects:

Very common ($\geq 1/10$)

Common ($\geq 1/100$ to $<1/10$)

Uncommon ($\geq 1/1000$ to $<1/100$)

Rare ($\geq 1/10,000$ to $<1/1000$)

Very rare ($<1/10,000$)

Not known (cannot be estimated from the available data).

Infections and infestations

Common: Mucocutaneous candidiasis

Not known : Overgrowth of non-susceptible organisms

Blood and lymphatic system disorders

Rare: Reversible leucopenia (including neutropenia) and thrombocytopenia

Not known: Reversible agranulocytosis and haemolytic anaemia. Prolongation of bleeding time and prothrombin time¹

Immune system disorders¹⁰

Not known: Angioneurotic oedema, anaphylaxis, serum sickness-like syndrome, hypersensitivity vasculitis.

Nervous system disorders

Uncommon: Dizziness, headache

Not known: Reversible hyperactivity, Aseptic meningitis and convulsions.

Cardiac disorders

Not known: Kounis syndrome

Gastrointestinal disorders

Very common: Diarrhoea

Common: Nausea³, vomiting

Uncommon: Indigestion

Not known: Antibiotic-associated colitis⁴, black hairy tongue, Drug-induced enterocolitis syndrome, Pancreatitis acute.

Hepatobiliary disorders

Uncommon: Rise in AST and/or ALT⁵ Not known: Hepatitis⁶ and cholestatic jaundice⁶.

Skin and subcutaneous tissue disorders⁷

Uncommon: Skin rash, pruritus, urticaria

Rare: Erythema multiforme

Not known: Stevens-Johnson syndrome, toxic epidermal necrolysis, bullous exfoliative-dermatitis, Linear IgA disease, acute generalised exanthemous pustulosis (AGEP)⁹

Drug reaction with eosinophilia and systemic symptoms (DRESS).

Renal and urinary disorders

Not known: Interstitial nephritis, crystalluria⁸ (including acute renal injury)

¹ See section 4.4

² See section 4.4

³ Nausea is more often associated with higher oral doses. If gastrointestinal reactions are evident, they may be reduced by taking amoxicillin-/clavulanic acid at the start of a meal.

⁴ Including pseudomembranous colitis and haemorrhagic colitis (see section 4.4)

⁵ A moderate rise in AST and/or ALT has been noted in patients treated with beta-lactam class antibiotics, but the significance of these findings is unknown.

⁶ These events have been noted with other penicillins and cephalosporins (see section 4.4).

⁷ If any hypersensitivity dermatitis reaction occurs, treatment should be discontinued (see section 4.4).

⁸ See section 4.9

⁹ See section 4.3

¹⁰ See section 4.4

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions directly via HPRA Pharmacovigilance, Website: www.hpra.ie.

4.9 Overdose

Symptoms and signs of overdose

Gastrointestinal symptoms and disturbance of the fluid and electrolyte balances may be evident. Amoxicillin crystalluria, in some cases leading to renal failure, has been observed (see section 4.4).

Amoxicillin has been reported to precipitate in bladder catheters, predominantly after intravenous administration of large doses. A regular check of patency should be maintained (see section 4.4).

Convulsions may occur in patients with impaired renal function or in those receiving high doses.

Treatment of intoxication

Gastrointestinal symptoms may be treated symptomatically, with attention to the water/electrolyte balance.

Amoxicillin/clavulanic acid can be removed from the circulation by haemodialysis.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: β -lactam antibacterials, combination of penicillin and beta-lactamase inhibitor ATC code: JOICR02

Mode of action

Amoxicillin is a semisynthetic penicillin (beta-lactam antibiotic) that inhibits one or more enzymes (often referred to as penicillin-binding proteins, PBPs) in the biosynthetic pathway of bacterial peptidoglycan, which is an integral structural component of the bacterial cell wall. Inhibition of peptidoglycan synthesis leads to weakening of the cell wall that is usually followed by cell lysis and death.

Amoxicillin is susceptible to degradation by beta-lactamases produced by resistant bacteria and therefore the spectrum of activity of amoxicillin alone does not include organisms which produce these enzymes.

Clavulanic acid is a beta-lactam agent structurally related to penicillins. It inactivates some beta-lactamase enzymes thereby preventing inactivation of amoxicillin. Clavulanic acid alone does not exert a clinically useful antibacterial effect.

Pharmacokinetic/Pharmacodynamic Relationship

The time above the minimum inhibitory concentration (T>MIC) is considered to be the major determinant of efficacy for amoxicillin.

Mechanisms of resistance

The two main mechanisms of resistance to amoxicillin/clavulanic acid are :

- Inactivation by those bacterial beta-lactamases that are not themselves inhibited by clavulanic acid, including class B, C and D.
- Alteration of PBPs, which reduce the affinity of the antibacterial agent for the target.

Impermeability of bacteria or efflux pump mechanisms may cause or contribute to bacterial resistance, particularly in Gram-negative bacteria.

Breakpoints

MIC breakpoints for amoxicillin/clavulanic acid are those of the European Committee on Antimicrobial Susceptibility Testing (EUCAST):

Organism	Susceptibility Breakpoints (microgram/ml)		
	Susceptible	Intermediate	Resistant
<i>Haemophilus influenzae</i> ¹	≤ 1	-	> 1
<i>Moraxella catarrhalis</i> ¹	≤ 1	-	> 1
<i>Staphylococcus aureus</i> ²	≤ 2	-	> 2
Coagulase-negative staphylococci ²	≤ 0.25		> 0.25
<i>Enterococcus</i> ¹	≤ 4	8	> 8
<i>Streptococcus</i> A, B, C, G ⁵	≤ 0.25	-	> 0.25
<i>Streptococcus pneumoniae</i> ³	≤ 0.5	1-2	> 2
Enterobacteriaceae ^{1,4}	-	-	> 8
Gram-negative Anaerobes ¹	≤ 4	8	> 8

Gram-positive Anaerobes ¹	≤ 4	8	> 8
Non-species related breakpoints ¹	≤ 2	4-8	> 8
¹ The reported values are for Amoxicillin concentrations. For susceptibility testing purposes, the concentration of Clavulanic acid is fixed at 2 mg/l. ² The reported values are Oxacillin concentrations. ³ Breakpoint values in the table are based on Ampicillin breakpoints. ⁴ The resistant breakpoint of R>8 mg/l ensures that all isolates with resistance mechanisms are reported resistant. ⁵ Breakpoint values in the table are based on Benzylpenicillin breakpoints			

The prevalence of resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of the agent in at least some types of infections is questionable.

Commonly susceptible species

Aerobic Gram-positive micro-organisms:

Enterococcus faecalis
Gardnerella vaginalis
Staphylococcus aureus (methicillin-susceptible)
Streptococcus agalactiae
*Streptococcus pneumoniae*¹
Streptococcus pyogenes and other beta-haemolytic streptococci
Streptococcus viridans group

Aerobic Gram-negative micro-organisms:

Capnocytophaga spp.
Eikenella corrodens
*Haemophilus influenzae*²
Moraxella catarrhalis
Pasteurella multocida

Anaerobic micro-organisms:

Bacteroides fragilis
Fusobacterium nucleatum
Prevotella spp.

Species for which acquired resistance may be a problem:

Aerobic Gram-positive micro-organisms:

*Enterococcus faecium*⁵

Aerobic Gram-negative micro-organisms:

Escherichia coli
Klebsiella oxytoca
Klebsiella pneumoniae
Proteus mirabilis
Proteus vulgaris

Inherently resistant organisms:

Aerobic Gram-positive micro-organisms:

Acinetobacter sp.
Citrobacter freundii
Enterobacter sp.
Legionella pneumophila
Morganella morganii
Providencia spp.
Pseudomonas sp.
Serratia sp.
Stenotrophomonas maltophilia

Other micro-organisms:

Chlamydophila pneumoniae

Chlamydomphila psittaci
Coxiella burnetti
Mycoplasma pneumoniae

§ Natural intermediate susceptibility in the absence of acquired mechanism of resistance.

£ All methicillin-resistant staphylococci are resistant to amoxicillin/clavulanic acid

¹ *Streptococcus pneumoniae* that are resistant to penicillin should not be treated with this presentation of amoxicillin/clavulanic acid (see sections 4.2 and 4.4).

² Strains with decreased susceptibility have been reported in some countries in the EU with a frequency higher than 10%.

5.2 Pharmacokinetic properties

Absorption

Amoxicillin and clavulanic acid are fully dissociated in aqueous solution at physiological pH. Both components are rapidly and well absorbed by the oral route of administration. Absorption of amoxicillin-clavulanic acid is optimized when taken at the start of a meal. Following oral administration, amoxicillin and clavulanic acid are approximately 70% bioavailable. The plasma profiles of both components are similar and the time to peak plasma concentration (T_{max}) in each case is approximately one hour.

The pharmacokinetic results for a study, in which amoxicillin/clavulanic acid (875 mg/125 mg tablets given twice daily) was administered in the fasting state to groups of healthy volunteers are presented below.

Mean (± SD) pharmacokinetic parameters					
Active substance(s) administered	Dose	C _{max}	T _{max} *	AUC (0-24h)	T 1/2
	(mg)	(microgram/ml)	(h)	(microgram.h/ml)	(h)
Amoxicillin					
AMX/CA 875 mg/125 mg	875	11.64 ± 2.78	1.50 (1.0-2.5)	53.52 ± 12.31	1.19 ± 0.21
Clavulanic acid					
AMX/CA 875 mg/125 mg	125	2.18 ± 0.99	1.25 (1.0-2.0)	10.16 ± 3.04	0.96 ± 0.12
AMX – amoxicillin, CA – clavulanic acid					
* Median (range)					

Amoxicillin and clavulanic acid serum concentrations achieved with amoxicillin/clavulanic acid are similar to those produced by the oral administration of equivalent doses of amoxicillin or clavulanic acid alone.

Distribution

About 25% of total plasma clavulanic acid and 18% of total plasma amoxicillin is bound to protein. The apparent volume of distribution is around 0.3-0.4 l/kg for amoxicillin and around 0.2 l/kg for clavulanic acid.

Following intravenous administration, therapeutic concentrations of both amoxicillin and clavulanic acid have been found in gall bladder, abdominal tissue, skin, fat, and muscle tissues; synovial and peritoneal fluids, bile and pus.

Amoxicillin does not adequately distribute into the cerebrospinal fluid.

Neither amoxicillin nor clavulanic acid is highly protein bound, studies show that about 25% for clavulanic acid and 18% for amoxicillin of total plasma drug content is bound to protein.

From animal studies there is no evidence for significant tissue retention of drug-derived material for either component.

Amoxicillin, like most penicillins, can be detected in breast milk. Trace quantities of clavulanic acid can also be detected in breast milk (see section 4.6).

Both amoxicillin and clavulanic acid have been shown to cross the placental barrier (see section 4.6).

Biotransformation

Amoxicillin is partly excreted in the urine as the inactive penicilloic acid in quantities equivalent to up to 10 to 25% of the initial dose. Clavulanic acid is extensively metabolized in man and eliminated in urine and faeces as carbon dioxide in expired air.

Elimination

The major route of elimination for amoxicillin is via the kidney, whereas for clavulanate it is by both renal and non-renal mechanisms.

Amoxicillin/clavulanic acid has a mean elimination half-life of approximately one hour and a mean total clearance of approximately 25 l/h in healthy subjects. Approximately 60 to 70% of the amoxicillin and approximately 40 to 65% of the clavulanic acid are excreted unchanged in urine during the first 6 h after administration of a single 250/125 mg or a single 500/125 mg tablet. Various studies have found the urinary excretion to be 50-85% for amoxicillin and between 27-60% for clavulanic acid over a 24 hour period. In the case of clavulanic acid, the largest amount of drug is excreted during the first 2 hours after administration.

Concomitant use of probenecid delays amoxicillin excretion but does not delay renal excretion of clavulanic acid (see section 4.5).

Age

The elimination half-life of amoxicillin is similar for children aged around 3 months to 2 years and older children and adults. For very young children (including preterm newborns) in the first week of life the interval of administration should not exceed twice daily administration due to immaturity of the renal pathway of elimination. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

Gender

Following oral administration of amoxicillin/clavulanic acid to healthy males and female subjects, gender has no significant impact on the pharmacokinetics of either amoxicillin or clavulanic acid.

Renal impairment

The total serum clearance of amoxicillin/clavulanic acid decreases proportionately with decreasing renal function. The reduction in drug clearance is more pronounced for amoxicillin than for clavulanic acid, as a higher proportion of amoxicillin is excreted via the renal route. Doses in renal impairment must therefore prevent undue accumulation of amoxicillin while maintaining adequate levels of clavulanic acid (see section 4.2).

Hepatic impairment

Hepatically impaired patients should be dosed with caution and hepatic function monitored at regular intervals.

5.3 Preclinical safety data

Nonclinical data reveal no special hazard for humans based on studies of safety pharmacology, genotoxicity and toxicity to reproduction.

Repeat dose toxicity studies performed in dogs with amoxicillin/clavulanic acid demonstrate gastric irritancy and vomiting, and discoloured tongue.

Carcinogenicity studies have not been conducted with amoxicillin/clavulanic acid or its components.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Core tablet-

Microcrystalline Cellulose (AVICEL Ph 102)

Croscarmellose sodium

Purified Talc

Magnesium Stearate

Coating -

Hypromellose (E-15)

Ethylcellulose

Propylene Glycol

Hydroxy propyl cellulose

Titanium dioxide

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

24 months

6.4 Special precautions for storage

Do not store above 25°C.

6.5 Nature and contents of container

The tablets are available in blisters and are sealed in aluminium pouches.
7, 10, 12, 14, 20, 21, 24 & 30 film-coated tablets in PVC/Al blister packs.
Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

No special requirements.

7 MARKETING AUTHORISATION HOLDER

Bluefish Pharmaceuticals AB
P.O. Box 49013
100 28 Stockholm
Sweden

8 MARKETING AUTHORISATION NUMBER

PA1436/002/003

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 6th November 2009

Date of last renewal: 16th May 2014

10 DATE OF REVISION OF THE TEXT

April 2023