

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Emazole Control 20 mg gastro-resistant tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each gastro-resistant tablet contains 20 mg esomeprazole (as magnesium dihydrate).

Excipient(s) with known effect:

Each 20 mg gastro-resistant tablet contains 12.90 mg – 14.76 mg sucrose and 0.81 mg glucose.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Gastro-resistant tablet

20 mg:

Light pink oval film-coated tablet.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Emazole Control 20 mg gastro-resistant tablets are indicated for:

The short-term treatment of reflux symptoms (e.g. heartburn and acid regurgitation) in adults.

4.2 Posology and method of administration

Posology

The recommended dose is 20 mg esomeprazole (one tablet) per day.

It might be necessary to take the tablets for 2-3 consecutive days to achieve improvement of symptoms. The duration of treatment is up to 2 weeks. Once complete relief of symptoms has occurred, treatment should be discontinued.

If no symptom relief is obtained within 2 weeks of continuous treatment, the patient should be instructed to consult a doctor.

Special populations

Patients with impaired renal function

Dose adjustment is not required in patients with impaired renal function. Due to limited experience in patients with severe renal insufficiency, such patients should be treated with caution (see section 5.2).

Patients with impaired hepatic function

Dose adjustment is not required in patients with mild to moderate liver impairment. For patients with severe liver impairment, a maximum dose of 20 mg esomeprazole should not be exceeded (see section 5.2).

Elderly (≥65 years old)

Dose adjustment is not required in the elderly.

Paediatric population

There is no relevant use of Emazole Control in the paediatric population below 18 years of age in the indication: "short-term treatment of reflux symptoms (e.g., heartburn and acid regurgitation)".

Method of administration

The tablets should be swallowed whole with liquid. The tablets should not be chewed or crushed.

For patients who have difficulty in swallowing, the tablets can also be dispersed in half a glass of non-carbonated water. No other liquids should be used as the enteric coating may be dissolved. Stir until the tablets disintegrate and drink the liquid with the pellets immediately or within 15 minutes. Rinse the glass with half a glass of water and drink. The pellets must not be chewed or crushed.

For patients who cannot swallow, the tablets can be dispersed in non-carbonated water and administered through a gastric tube. It is important that the appropriateness of the selected syringe and tube is carefully tested. For preparation and administration instructions see section 6.6.

4.3 Contraindications

Hypersensitivity to the active substance, to substituted benzimidazoles or to any of the excipients listed in section 6.1. Esomeprazole must not be used concomitantly with nelfinavir (see section 4.5).

4.4 Special warnings and precautions for use

General

Patients should be instructed to consult a doctor:

In the presence of any alarm symptom (e.g. significant unintentional weight loss, recurrent vomiting, dysphagia, haematemesis or melaena) and when gastric ulcer is suspected or present, malignancy should be excluded, as treatment with esomeprazole may alleviate symptoms and delay diagnosis.

They have had previous gastric ulcer or gastrointestinal surgery.

- They have been on continuous symptomatic treatment of indigestion or heartburn for 4 or more weeks.
- They have jaundice or severe liver disease.
- They are aged over 55 years with new or recently changed symptoms.

Patients with long-term recurrent symptoms of indigestion or heartburn should see their doctor at regular intervals. Patients over 55 years taking any non-prescription indigestion or heartburn remedy on a daily basis should inform their pharmacist or doctor.

Patients should not take Emazole Control as a long term preventive medicinal product.

Treatment with proton pump inhibitors (PPIs) may lead to a slightly increased risk of gastrointestinal infections such as Salmonella and Campylobacter and in hospitalised patients, also possibly Clostridium difficile (see section 5.1).

Patients should consult their doctor before taking this medicinal product if they are due to have an endoscopy or urea breath test.

Absorption of vitamin B12

Esomeprazole, as all acid-blocking medicines, may reduce the absorption of vitamin B12 (cyanocobalamin) due to hypo- or achlorhydria. This should be considered in patients with reduced body stores or risk factors for reduced vitamin B12 absorption on long-term therapy.

Hypomagnesaemia

Severe hypomagnesaemia has been reported in patients treated with PPIs like esomeprazole for at least three months, and in most cases for a year. Serious manifestations of hypomagnesaemia such as fatigue, tetany, delirium, convulsions, dizziness and ventricular arrhythmia can occur but they may begin insidiously and be overlooked. In most affected patients, hypomagnesaemia improved after magnesium replacement and discontinuation of the PPI.

For patients expected to be on prolonged treatment or who take PPIs with digoxin or drugs that may cause hypomagnesaemia (e.g., diuretics), health care professionals should consider measuring magnesium levels before starting PPI treatment and periodically during treatment

Risk of fracture

Proton pump inhibitors, especially if used in high doses and over long durations (>1 year), may modestly increase the risk of hip, wrist and spine fracture, predominantly in the elderly or in presence of other recognised risk factors. Observational studies suggest that proton pump inhibitors may increase the overall risk of fracture by 10–40%. Some of this increase may be due to other risk factors. Patients at risk of osteoporosis should receive care according to current clinical guidelines and they should have an adequate intake of vitamin D and calcium.

Subacute cutaneous lupus erythematosus (SCLE)

Proton pump inhibitors are associated with very infrequent cases of SCLE. If lesions occur, especially in sun-exposed areas of the skin, and if accompanied by arthralgia, the patient should seek medical help promptly and the health care professional should consider stopping Emazole Control. SCLE after previous treatment with a proton pump inhibitor may increase the risk of SCLE with other proton pump inhibitors.

Combination with other medicinal products

Co-administration of esomeprazole with atazanavir is not recommended (see section 4.5). If the combination of atazanavir with a proton pump inhibitor is judged unavoidable, close clinical monitoring is recommended in combination with an increase in the dose of atazanavir to 400 mg with 100 mg of ritonavir; esomeprazole 20 mg should not be exceeded.

Esomeprazole is a CYP2C19 inhibitor. When starting or ending treatment with esomeprazole, the potential for interactions with drugs metabolised through CYP2C19 should be considered. An interaction is observed between clopidogrel and esomeprazole (see section 4.5). The clinical relevance of this interaction is uncertain. As a precaution, concomitant use of esomeprazole and clopidogrel should be discouraged.

Interference with laboratory tests

Increased Chromogranin A (CgA) level may interfere with investigations for neuroendocrine tumours. To avoid this interference, Emazole Control treatment should be stopped for at least 5 days before CgA measurements (see section 5.1). If CgA and gastrin levels have not returned to reference range after initial measurement, measurements should be repeated 14 days after cessation of proton pump inhibitor treatment.

Excipients with known effect

This medicinal product contains glucose and sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Effects of esomeprazole on the pharmacokinetics of other medicinal products

Medicinal products with pH dependent absorption

Gastric acid suppression during treatment with esomeprazole and other PPIs might decrease or increase the absorption of medicinal products with a gastric pH dependent absorption. As with other medicinal products that decrease intragastric acidity, the absorption of medicinal products such as ketoconazole, itraconazole and erlotinib can decrease and the absorption of digoxin can increase during treatment with esomeprazole. Concomitant treatment with omeprazole (20 mg daily) and digoxin in healthy subjects increased the bioavailability of digoxin by 10% (up to 30% in two out of ten subjects). Digoxin toxicity has been rarely reported. However, caution should be exercised when esomeprazole is given at high doses in elderly patients. Therapeutic drug monitoring of digoxin should then be reinforced.

Protease inhibitors

Omeprazole has been reported to interact with some protease inhibitors. The clinical importance and the mechanisms behind these reported interactions are not always known. Increased gastric pH during omeprazole treatment may change the absorption of the protease inhibitors. Other possible interaction mechanisms are via inhibition of CYP2C19. For atazanavir and nelfinavir, decreased serum levels have been reported when given together with omeprazole and concomitant administration is not recommended. Co-administration of omeprazole (40 mg once daily) with atazanavir 300 mg/ritonavir 100 mg to healthy volunteers resulted in a substantial reduction in atazanavir exposure (approximately 75% decrease in AUC, C_{max} and C_{min}). Increasing the atazanavir dose to 400 mg did not compensate for the impact of omeprazole on atazanavir exposure. The co-administration of omeprazole (20 mg once daily) with atazanavir 400 mg/ritonavir 100 mg to healthy volunteers resulted in a decrease of approximately 30% in the atazanavir exposure as compared with the exposure observed with atazanavir 300 mg/ritonavir 100 mg once daily without omeprazole 20 mg once daily. Co-administration of omeprazole (40 mg once daily)

reduced mean nelfinavir AUC, C_{max} and C_{min} by 36–39 % and mean AUC, C_{max} and C_{min} for the pharmacologically active metabolite M8 was reduced by 75-92%.

Due to the similar pharmacodynamic effects and pharmacokinetic properties of omeprazole and esomeprazole, concomitant administration with esomeprazole and atazanavir is not recommended (see section 4.4) and concomitant administration with esomeprazole and nelfinavir is contraindicated (see section 4.3.).

For saquinavir (with concomitant ritonavir), increased serum levels (80-100%) have been reported during concomitant omeprazole treatment (40 mg once daily). Treatment with omeprazole 20 mg once daily had no effect on the exposure of darunavir (with concomitant ritonavir) and amprenavir (with concomitant ritonavir). Treatment with esomeprazole 20 mg once daily had no effect on the exposure of amprenavir (with and without concomitant ritonavir). Treatment with omeprazole 40 mg once daily had no effect on the exposure of lopinavir (with concomitant ritonavir).

Medicinal products metabolised by CYP2C19

Esomeprazole inhibits CYP2C19, the major esomeprazole metabolising enzyme. Thus, when esomeprazole is combined with medicinal products metabolised by CYP2C19, such as diazepam, citalopram, imipramine, clomipramine, phenytoin, etc., the plasma concentrations of these medicinal products may be increased and a dose reduction could be needed. This should be considered especially when prescribing esomeprazole for on demand therapy.

Diazepam

Concomitant administration of 30 mg esomeprazole resulted in a 45% decrease in clearance of the CYP2C19 substrate diazepam.

Phenytoin

Concomitant administration of 40 mg esomeprazole resulted in a 13% increase in trough plasma levels of phenytoin in epileptic patients. It is recommended to monitor the plasma concentrations of phenytoin when treatment with esomeprazole is introduced or withdrawn.

Voriconazole

Omeprazole (40 mg once daily) increased voriconazole (a CYP2C19 substrate) C_{max} and AUC_{τ} by 15% and 41%, respectively.

Warfarin

Concomitant administration of 40 mg esomeprazole to warfarin-treated patients in a clinical trial showed that coagulation times were within the accepted range. However, post-marketing, a few isolated cases of elevated INR of clinical significance have been reported during concomitant treatment. Monitoring is recommended when initiating and ending concomitant esomeprazole treatment during treatment with warfarin or other coumarin derivatives.

Cilostazol

Omeprazole as well as esomeprazole act as inhibitors of CYP2C19. Omeprazole, given in doses of 40 mg to healthy subjects in a cross-over study, increased C_{max} and AUC for cilostazol by 18% and 26% respectively, and one of its active metabolites by 29% and 69% respectively.

Cisapride

In healthy volunteers, concomitant administration of 40 mg esomeprazole resulted in a 32% increase in area under the plasma concentration-time curve (AUC) and a 31% prolongation of elimination half-life ($t_{1/2}$) but no significant increase in peak plasma levels of cisapride. The slightly prolonged QTc interval observed after administration of cisapride alone, was not further prolonged when cisapride was given in combination with esomeprazole (see also section 4.4).

Clopidogrel

Results from studies in healthy subjects have shown a pharmacokinetic (PK)/ pharmacodynamic (PD) interaction between clopidogrel (300 mg loading dose/75 mg daily maintenance dose) and esomeprazole (40 mg p.o.daily) resulting in decreased exposure to the active metabolite of clopidogrel by an average of 40% and resulting in decreased maximum inhibition of (ADP induced) platelet aggregation by an average of 14%.

When clopidogrel was given together with a fixed dose combination of esomeprazole 20 mg + ASA 81 mg compared to clopidogrel alone in a study in healthy subjects there was a decreased exposure by almost 40% of the active metabolite of clopidogrel. However, the maximum levels of inhibition of (ADP induced) platelet aggregation in these subjects were the same in the clopidogrel and the clopidogrel + the combined (esomeprazole + ASA) product groups.

Inconsistent data on the clinical implications of a PK/PD interaction of esomeprazole in terms of major cardiovascular events have been reported from both observational and clinical studies. As a precaution concomitant use of clopidogrel should be discouraged.

Tacrolimus

Concomitant administration of esomeprazole has been reported to increase the serum levels of tacrolimus. A reinforced monitoring of tacrolimus concentrations as well as renal function (creatinine clearance) should be performed, and dose of tacrolimus adjusted if needed.

Methotrexate

When given together with PPIs, methotrexate levels have been reported to increase in some patients. In high-dose methotrexate administration a temporary withdrawal of esomeprazole may need to be considered.

Investigated medicinal products with no clinically relevant interaction

Amoxicillin and quinidine

Esomeprazole has been shown to have no clinically relevant effects on the pharmacokinetics of amoxicillin or quinidine.

Naproxen or rofecoxib

Studies evaluating concomitant administration of esomeprazole and either naproxen or rofecoxib did not identify any clinically relevant pharmacokinetic interactions during short-term studies.

Effects of other medicinal products on the pharmacokinetics of esomeprazole

Medicinal products which inhibit CYP2C19 and/or CYP3A4

Esomeprazole is metabolized by CYP2C19 and CYP3A4. Concomitant administration of esomeprazole and a CYP3A4 inhibitor, clarithromycin (500 mg twice daily), resulted in a doubling of the exposure (AUC) to esomeprazole. Concomitant administration of esomeprazole and a combined inhibitor of CYP2C19 and CYP3A4 may result in more than doubling of the esomeprazole exposure. The CYP2C19 and CYP 3A4 inhibitor voriconazole increased omeprazole AUC_τ by 280%. A dose adjustment of esomeprazole is not regularly required in either of these situations. However, dose adjustment should be considered in patients with severe hepatic impairment and if long-term treatment is indicated.

Medicinal products which induce CYP2C19 and/or CYP3A4

Medicinal products known to induce CYP2C19 or CYP3A4 or both (such as rifampicin and St. John's wort) may lead to decreased esomeprazole serum levels by increasing the esomeprazole metabolism.

Paediatric population

Interaction studies have only been performed in adults.

4.6 Fertility, pregnancy and lactation

Pregnancy

Clinical data on exposed pregnancies with esomeprazole are insufficient. With the racemic mixture, omeprazole, data on a larger number of exposed pregnancies from epidemiological studies indicate no malformative nor foetotoxic effect. Animal studies with esomeprazole do not indicate direct or indirect harmful effects with respect to embryonal/foetal development. Animal studies with the racemic mixture do not indicate direct or indirect harmful effects with respect to pregnancy, parturition or postnatal development. Caution should be exercised when prescribing esomeprazole to pregnant women. A moderate amount of data on pregnant women (between 300-1000 pregnancy outcomes) indicates no malformative or foeto/neonatal toxicity of esomeprazole.

Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3).

Breast-feeding

It is not known whether esomeprazole is excreted in human breast milk. There is insufficient information on the effects of esomeprazole in newborns/infants. Esomeprazole should not be used during breast-feeding.

Fertility

Animal studies with the racemic mixture omeprazole, given by oral administration do not indicate effects with respect to fertility.

4.7 Effects on ability to drive and use machines

Esomeprazole has minor influence on the ability to drive or use machines. Adverse reactions such as dizziness (uncommon) and blurred vision (rare) have been reported (see section 4.8). If affected, patients should not drive or use machines.

4.8 Undesirable effectsSummary of the safety profile

Headache, abdominal pain, diarrhoea and nausea are among those adverse reactions that have been most commonly reported in clinical trials (and also from post-marketing use). In addition, the safety profile is similar for different formulations, treatment indications, age groups and patient populations. No dose-related adverse reactions have been identified.

Tabulated list of adverse reactions

The following adverse drug reactions have been identified or suspected in the clinical trials programme for esomeprazole and post-marketing. None was found to be dose-related. The reactions are classified according to the MedDRA frequency convention:

very common $\geq 1/10$; common $\geq 1/100$ to $< 1/10$; uncommon $\geq 1/1,000$ to $< 1/100$; rare $\geq 1/10,000$ to $< 1/1,000$; very rare $< 1/10,000$; not known (frequency cannot be estimated from the available data).

	Very common	Common	Uncommon	Rare	Very rare	Not known
Blood and lymphatic system disorder				Leukopenia, thrombocytopenia	Agranulocytosis, pancytopenia	
Immune system disorders				Hypersensitivity reactions e.g. fever, angioedema and anaphylactic reaction/ shock		
Metabolism and nutrition disorders			Peripheral oedema	Hyponatraemia		Hypomagnesemia (see section 4.4); severe hypomagnesaemia can correlate with hypocalcaemia. Hypomagnasemia may also be associated with hypokalaemia.
Psychiatric disorders			Insomnia	Agitation, confusion, depression	Aggression, hallucinations	
Nervous system disorders		Headache	Dizziness paraesthesia somnolence	Taste disturbance		
Eye disorders				Blurred vision		
Ear and labyrinth disorders			Vertigo			
Respiratory, thoracic and mediastinal disorders				Broncho-spasm		

	Very common	Common	Uncommon	Rare	Very rare	Not known
Gastrointestinal disorders		Abdominal pain constipation diarrhoea flatulence nausea/ vomiting, fundic gland polyps (benign)	Dry mouth	Stomatitis, gastrointestinal candidiasis		Microscopic colitis
Hepatobiliary disorders			Increased liver enzymes	Hepatitis with or without jaundice	Hepatic failure, encephalopathy in patients with pre-existing liver disease	
Skin and subcutaneous tissue disorders			Dermatitis pruritus rash urticaria	Alopecia photosensitivity	Erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis (TEN)	Subacute cutaneous lupus erythematosus (see section 4.4)
Musculoskeletal and connective tissue disorders			Fracture of the hip, wrist or spine (see section 4.4)	Arthralgia myalgia	Muscular weakness	
Renal and urinary disorders					Interstitial nephritis; in some patients renal failure has been reported concomitantly.	
Reproductive system and breast disorders					Gynaecomastia	
General disorder and administration site conditions				Malaise increased sweating		

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRC Pharmacovigilance. Website: www.hpra.ie

4.9 Overdose

Symptoms

There is very limited experience to date with deliberate overdose. The symptoms described in connection with 280 mg were gastrointestinal symptoms and weakness. Single doses of 80 mg esomeprazole were uneventful.

Treatment

No specific antidote is known. Esomeprazole is extensively plasma protein bound and is therefore not readily dialyzable. As in any case of overdose, treatment should be symptomatic and general supportive measures should be utilised.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Drugs for peptic ulcer and gastro-oesophageal reflux disease (GORD), proton pump inhibitor
ATC code: A02B C05

Esomeprazole is the S-isomer of omeprazole and reduces gastric acid secretion through a specific targeted mechanism of action. It is a specific inhibitor of the acid pump in the parietal cell. Both the R- and S-isomer of omeprazole have similar pharmacodynamic activity.

Mechanism of action

Esomeprazole is a weak base and is concentrated and converted to the active form in the highly acidic environment of the secretory canaliculi of the parietal cell, where it inhibits the enzyme H^+K^+ -ATPase – the acid pump and inhibits both basal and stimulated acid secretion.

Pharmacodynamic effects

After oral dosing with esomeprazole 20 mg and 40 mg the onset of effect occurs within one hour. After repeated administration with 20 mg esomeprazole once daily for five days, mean peak acid output after pentagastrin stimulation is decreased 90% when measured 6-7 hours after dosing on day five.

After five days of oral dosing with 20 mg and 40 mg of esomeprazole, intragastric pH above 4 was maintained for a mean time of 13 hours and 17 hours, respectively over 24 hours in symptomatic GORD patients. The proportion of patients maintaining an intragastric pH above 4 for at least 8, 12 and 16 hours respectively were for esomeprazole 20 mg 76%, 54% and 24%. Corresponding proportions for esomeprazole 40 mg were 97%, 92% and 56%.

Using AUC as a surrogate parameter for plasma concentration, a relationship between inhibition of acid secretion and exposure has been shown.

Therapeutic effects of acid inhibition

Healing of reflux oesophagitis with esomeprazole 40 mg occurs in approximately 78% of patients after four weeks, and in 93% after eight weeks.

One week treatment with esomeprazole 20 mg twice daily and appropriate antibiotics, results in successful eradication of *H. pylori* in approximately 90% of patients.

After eradication treatment for one week there is no need for subsequent monotherapy with antisecretory medicinal products for effective ulcer healing and symptom resolution in uncomplicated duodenal ulcers.

In a randomised, double blind, placebo-controlled clinical study, patients with endoscopically confirmed peptic ulcer bleeding characterised as Forrest Ia, Ib, IIa or IIb (9%, 43%, 38% and 10% respectively) were randomised to receive esomeprazole solution for infusion (n=375) or placebo (n=389). Following endoscopic haemostasis, patients received either 80 mg esomeprazole as an intravenous infusion over 30 minutes followed by a continuous infusion of 8 mg per hour or placebo for 72 hours. After the initial 72 hour period, all patients received open-label 40 mg oral esomeprazole for 27 days for acid suppression. The occurrence of rebleeding within 3 days was 5.9% in the esomeprazole treated group compared to 10.3% for the placebo group. At 30 days post-treatment, the occurrence of rebleeding in the esomeprazole treated versus the placebo treated group was 7.7% vs 13.6%.

Other effects related to acid inhibition

During treatment with antisecretory medicinal products, serum gastrin increases in response to the decreased acid secretion. Also CgA increases due to decreased gastric acidity. The increased CgA level may interfere with investigations for neuroendocrine tumours. Available published evidence suggests that proton pump inhibitors should be discontinued between 5 days and 2 weeks prior to CgA measurements. This is to allow CgA levels that might be spuriously elevated following PPI treatment to return to reference range.

An increased number of ECL cells possibly related to the increased serum gastrin levels, have been observed in both children and adults during long term treatment with esomeprazole. The findings are considered to be of no clinical significance.

During long-term treatment with antisecretory medicinal products gastric glandular cysts have been reported to occur at a somewhat increased frequency. These changes are a physiological consequence of pronounced inhibition of acid secretion, are benign and appear to be reversible.

Decreased gastric acidity due to any means including proton pump inhibitors, increases gastric counts of bacteria normally present in the gastrointestinal tract. Treatment with proton pump inhibitors may lead to slightly increased risk of gastrointestinal infections such as *Salmonella*, *Campylobacter* and, in hospitalised patients, possibly also *Clostridium difficile*.

Clinical efficacy

In two studies with ranitidine as an active comparator, esomeprazole showed better effect in healing of gastric ulcers in patients using NSAIDs, including COX-2 selective NSAIDs.

In two studies with placebo as comparator, esomeprazole showed better effect in the prevention of gastric and duodenal ulcers in patients using NSAIDs (aged > 60 and/or with previous ulcer), including COX-2 selective NSAIDs.

Paediatric population

In a study in paediatric GORD patients (<1 to 17 years of age) receiving long-term PPI treatment, 61% of the children developed minor degrees of ECL cell hyperplasia with no known clinical significance and with no development of atrophic gastritis or carcinoid tumours.

5.2 Pharmacokinetic properties

Absorption

Esomeprazole is acid labile and is administered orally as enteric-coated granules. In vivo conversion to the R-isomer is negligible. Absorption of esomeprazole is rapid, with peak plasma levels occurring approximately 1-2 hours after dose. The absolute bioavailability is 64% after a single dose of 40 mg and increases to 89% after repeated once-daily administration. For 20 mg esomeprazole the corresponding values are 50% and 68%, respectively.

Food intake both delays and decreases the absorption of esomeprazole although this has no significant influence on the effect of esomeprazole on intragastric acidity.

Distribution

The apparent volume of distribution at steady state in healthy subjects is approximately 0.22 l/kg body weight. Esomeprazole is 97% plasma protein bound.

Biotransformation

Esomeprazole is completely metabolised by the cytochrome P450 system (CYP). The major part of the metabolism of esomeprazole is dependent on the polymorphic CYP2C19, responsible for the formation of the hydroxy- and desmethyl metabolites of esomeprazole. The remaining part is dependent on another specific isoform, CYP3A4, responsible for the formation of esomeprazole sulphone, the main metabolite in plasma.

Elimination

The parameters below reflect mainly the pharmacokinetics in individuals with a functional CYP2C19 enzyme, extensive metabolisers.

Total plasma clearance is about 17 l/h after a single dose and about 9 l/h after repeated administration. The plasma elimination half-life is about 1.3 hours after repeated once-daily dosing. Esomeprazole is completely eliminated from plasma between doses with no tendency for accumulation during once-daily administration.

The major metabolites of esomeprazole have no effect on gastric acid secretion. Almost 80% of an oral dose of esomeprazole is excreted as metabolites in the urine, the remainder in the faeces. Less than 1% of the parent drug is found in urine.

Linearity/non-linearity

The pharmacokinetics of esomeprazole has been studied in doses up to 40 mg twice daily. The area under the plasma concentration-time curve increases with repeated administration of esomeprazole. This increase is dose-dependent and results in a more than dose proportional increase in AUC after repeated administration. This time- and dose-dependency is due to a decrease of first pass metabolism and systemic clearance probably caused by an inhibition of the CYP2C19 enzyme by esomeprazole and/or its sulphone metabolite.

Special patient populations

Poor metabolisers

Approximately $2.9 \pm 1.5\%$ of the population lack a functional CYP2C19 enzyme and are called poor metabolisers. In these individuals the metabolism of esomeprazole is probably mainly catalysed by CYP3A4. After repeated once-daily administration of 40 mg esomeprazole, the mean area under the plasma concentration-time curve was approximately 100% higher in poor metabolisers than in subjects having a functional CYP2C19 enzyme (extensive metabolisers). Mean peak plasma concentrations were increased by about 60%. These findings have no implications for the posology of esomeprazole.

Elderly

The metabolism of esomeprazole is not significantly changed in elderly subjects (71-80 years of age).

Gender

Following a single dose of 40 mg esomeprazole the mean area under the plasma concentration-time curve is approximately 30% higher in females than in males. No gender difference is seen after repeated once-daily administration. These findings have no implications for the posology of esomeprazole.

Hepatic impairment

The metabolism of esomeprazole in patients with mild to moderate liver dysfunction may be impaired. The metabolic rate is decreased in patients with severe liver dysfunction resulting in a doubling of the area under the plasma concentration-time curve of esomeprazole. Therefore, a maximum of 20 mg should not be exceeded in patients with severe dysfunction. Esomeprazole or its major metabolites do not show any tendency to accumulate with once-daily dosing.

Renal impairment

No studies have been performed in patients with decreased renal function. Since the kidney is responsible for the excretion of the metabolites of esomeprazole but not for the elimination of the parent compound, the metabolism of esomeprazole is not expected to be changed in patients with impaired renal function.

Paediatric population

Adolescents 12-18 years:

Following repeated dose administration of 20 mg and 40 mg esomeprazole, the total exposure (AUC) and the time to reach maximum plasma drug concentration (t_{max}) in 12 to 18 year-olds was similar to that in adults for both esomeprazole doses.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction and development. Adverse reactions not observed in clinical studies, but seen in animals at exposure levels similar to clinical exposure levels and with possible relevance to clinical use were as follows:

Carcinogenicity studies in the rat with the racemic mixture have shown gastric ECL-cell hyperplasia and carcinoids. These gastric effects in the rat are the result of sustained, pronounced hypergastrinaemia secondary to reduced production of gastric acid and are observed after long-term treatment in the rat with inhibitors of gastric acid secretion.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core

Sucrose
Maize starch
Liquid glucose
Hydroxypropylcellulose
Povidone
Talc
Titanium dioxide (E171)
Methacrylic acid-ethyl acrylate copolymer (1:1)
Glycerol monostearate
Propylene glycol
Stearic acid
Polysorbate 80
Simeticone
Microcrystalline cellulose

Macrogol 6000
Crospovidone
Silica colloidal anhydrous
Magnesium stearate

Tablet coating

Hypromellose
Macrogol 6000
Titanium dioxide (E171)
Talc
Iron oxide, red (E172)
Iron oxide, yellow (E172)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

Aclar/aluminium blisters:
18 months

Aluminium/aluminium blister:
24 months

HDPE bottles:
24 months

Shelf life after first opening: 6 months

Do not store above 30°C. Keep the container tightly closed in order to protect from moisture.

6.4 Special precautions for storage

Aluminium/aluminium blisters and aclar/aluminium blisters:
Do not store above 25°C.

HDPE bottles:

Storage conditions before opening of the HDPE bottle:

Do not store above 30°C.

For storage conditions after first opening of the HDPE bottle, see section 6.3.

6.5 Nature and contents of container

Aluminium/aluminium blisters and aclar/aluminium blisters with 7, 14 & 28 gastro-resistant tablets.

HDPE bottles with PP cap and desiccant with 7, 14 gastro-resistant tablets.

Not all pack sizes may be marketed

6.6 Special precautions for disposal and other handling

No special requirements.

7 MARKETING AUTHORISATION HOLDER

IQ Pharmatek Ltd
Gurtnafleur
Old Waterford Road
Clonmel
Co Tipperary
08 March 2024

Ireland

8 MARKETING AUTHORISATION NUMBER

PA22777/001/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 18th October 2019

Date of last renewal: 08th March 2024

10 DATE OF REVISION OF THE TEXT

March 2024