

## Summary of Product Characteristics

### 1 NAME OF THE MEDICINAL PRODUCT

Paclitaxel 6mg/ml Concentrate for Solution for Infusion

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

1ml contains 6mg of the active ingredient, paclitaxel

1 vial of 5ml contains 30mg of paclitaxel

1 vial of 16.7ml contains 100mg of paclitaxel

1 vial of 50ml contains 300mg of paclitaxel

Excipients:

1ml contains 525mg of macroglycerol ricinoleate

1 vial of 5ml contains 2.625g of macroglycerol ricinoleate

1 vial of 16.7ml contains 8.768g of macroglycerol ricinoleate

1 vial of 50ml contains 26.250g of macroglycerol ricinoleate

1ml contains 497 mg of ethanol

1 vial of 5ml contains 2.485g of ethanol

1 vial of 16.7ml contains 8.300g of ethanol

1 vial of 50 ml contains 24.850g of ethanol

For full list of excipients, see 6.1

### 3 PHARMACEUTICAL FORM

Concentrate for solution for infusion

A clear, colourless to pale yellow viscous solution

### 4 CLINICAL PARTICULARS

#### 4.1 Therapeutic Indications

**Ovarian carcinoma:** in the first-line chemotherapy of ovarian cancer, paclitaxel infusion is indicated for the treatment of patients with advanced carcinoma of the ovary or with residual disease (> 1 cm) after initial laparotomy, in combination with cisplatin. In the second-line chemotherapy of ovarian cancer, paclitaxel infusion is indicated for the treatment of metastatic carcinoma of the ovary after failure of standard, platinum containing therapy.

**Breast carcinoma:** In the adjuvant setting, paclitaxel infusion is indicated for the treatment of patients with node-positive breast carcinoma following anthracycline and cyclophosphamide (AC) therapy. Adjuvant treatment with paclitaxel infusion should be regarded as an alternative to extended AC therapy. Paclitaxel infusion is indicated for the initial treatment of locally advanced or metastatic breast cancer either in combination with an anthracycline in patients for whom anthracycline therapy is suitable, or in combination with trastuzumab, in patients who over-express HER-2 (human epidermal growth factor receptor 2) at a 3+ level as determined by immunohistochemistry and for whom an anthracycline is not suitable (see 4.4 and 5.1). As a single agent, paclitaxel infusion is indicated for the treatment of metastatic carcinoma of the breast in patients who have failed, or are not candidates for standard, anthracycline containing therapy.

**Advanced non-small cell lung carcinoma:** paclitaxel infusion, in combination with cisplatin, is indicated for the treatment of non-small cell lung carcinoma (NSCLC) in patients who are not candidates for potentially curative surgery and/or radiation therapy.

**AIDS-related Kaposi's sarcoma:** paclitaxel infusion is indicated for the treatment of patients with advanced AIDS-related Kaposi's sarcoma (KS) who have failed prior liposomal anthracycline therapy. Limited efficacy data supports this indication, a summary of the relevant studies is shown in section 5.1.

## 4.2 Posology and method of administration

All patients must be premedicated with corticosteroids, antihistamines, and H<sub>2</sub> antagonists prior to paclitaxel infusion, e.g.

Drug	Dose	Administration prior to paclitaxel infusion
dexamethasone	20 mg oral* or IV	For oral administration: approximately 12 and 6 hours or for IV administration: 30 to 60 min
diphenhydramine**	50 mg IV	30 to 60 min
cimetidine or ranitidine	300 mg IV 50 mg IV	30 to 60 min

\* 8-20 mg for KS patients

\*\* or an equivalent antihistamine e.g. chlorpheniramine

Paclitaxel infusion should be administered through an in-line filter with a microporous membrane  $\leq 0.22 \mu\text{m}$  (see 6.6).

### Ovarian carcinoma

**First-line chemotherapy of ovarian carcinoma:** although other dosage regimens are under investigation, a combination regimen of paclitaxel infusion and cisplatin is recommended. According to duration of infusion, two doses of paclitaxel infusion are recommended: paclitaxel infusion 175 mg/m<sup>2</sup> administered intravenously over 3 hours, followed by cisplatin at a dose of 75 mg/m<sup>2</sup> every three weeks or paclitaxel infusion 135 mg/m<sup>2</sup>, in a 24-hour infusion, followed by cisplatin 75 mg/m<sup>2</sup>, with a three week interval between courses (see 5.1).

**Second-line chemotherapy of ovarian carcinoma:** the recommended dose of paclitaxel infusion is 175 mg/m<sup>2</sup> administered over a period of three hours, with a three week interval between courses.

### Breast carcinoma

**Adjuvant chemotherapy in breast carcinoma:** the recommended dose of paclitaxel infusion is 175 mg/m<sup>2</sup> administered over a period of three hours every three weeks for four courses, following AC therapy.

**First-line chemotherapy of breast carcinoma:** when used in combination with doxorubicin (50 mg/m<sup>2</sup>), paclitaxel infusion should be administered 24 hours after doxorubicin. The recommended dose of paclitaxel infusion is 220 mg/m<sup>2</sup> administered intravenously over a period of three hours, with a three-week interval between courses (see 4.5 and 5.1).

When used in combination with trastuzumab, the recommended dose of paclitaxel infusion is 175 mg/m<sup>2</sup> administered intravenously over a period of 3 hours, with a 3-week interval between courses (see 5.1).

Paclitaxel infusion may be started the day following the first dose of trastuzumab or immediately after the subsequent doses of trastuzumab if the preceding dose of trastuzumab was well tolerated (for detailed trastuzumab posology see the Summary of Product Characteristics of trastuzumab).

**Second-line chemotherapy of breast carcinoma:** the recommended dose of paclitaxel infusion is 175 mg/m<sup>2</sup> administered over a period of three hours, with a three-week interval between courses.

#### **Advanced non-small cell lung carcinoma**

**Treatment of advanced NSCLC:** the recommended dose of paclitaxel infusion is 175 mg/m<sup>2</sup> administered over a period of three hours, followed by cisplatin 80 mg/m<sup>2</sup>, with a three week interval between courses.

#### **AIDS-related Kaposi's sarcoma**

**Treatment of AIDS-related KS:** the recommended dose of paclitaxel infusion is 100 mg/m<sup>2</sup> administered as a three-hour intravenous infusion every two weeks. Subsequent doses of paclitaxel infusion should be administered according to individual patient tolerance.

Paclitaxel infusion should not be readministered until the neutrophil count is 1,500/mm<sup>3</sup> ( $\geq 1,000$ /mm<sup>3</sup> for KS patients) and the platelet count is 100,000/mm<sup>3</sup> ( $\geq 75,000$ /mm<sup>3</sup> for KS patients). Patients who experience severe neutropenia (neutrophil count < 500/mm<sup>3</sup> for 7 days) or severe peripheral neuropathy should receive a dose reduction of 20% for subsequent courses (25% for KS patients) (see 4.4).

**Patients with hepatic impairment:** Inadequate data are available to recommend dosage alterations in patients with mild to moderate hepatic impairments (see 4.4 and 5.2). Patients with severe hepatic impairment should not be treated with paclitaxel.

**Paediatric use:** Paclitaxel is not recommended for use in children below 18 years due to lack of data on safety and efficacy.

### **4.3 Contraindications**

Paclitaxel is contra-indicated in patients with severe hypersensitivity to paclitaxel or to any ingredient, especially macroglycerol ricinoleate (see 4.4).

Paclitaxel is contraindicated during pregnancy and lactation (see 4.6) and should not be used in patients with baseline neutrophils < 1,500/mm<sup>3</sup> (< 1,000/mm<sup>3</sup> for KS patients).

In KS, paclitaxel is also contraindicated in patients with concurrent, serious, uncontrolled infections.

### **4.4 Special warnings and precautions for use**

Paclitaxel should be administered under the supervision of a physician experienced in the use of cancer chemotherapeutic agents. Since significant hypersensitivity reactions may occur, appropriate supportive equipment should be available.

Patients must be pre-treated with corticosteroids, antihistamines and H<sub>2</sub> antagonists (see 4.2).

Paclitaxel should be given *before* cisplatin when used in combination (see 4.5).

**Significant hypersensitivity reactions** characterised by dyspnoea and hypotension requiring treatment, angioedema and generalised urticaria have occurred in < 1% of patients receiving paclitaxel after adequate premedication. These reactions are probably histamine-mediated. In the case of severe hypersensitivity reactions, paclitaxel should be discontinued immediately, symptomatic therapy should be initiated and the patient should not be rechallenged with the drug.

**Bone marrow suppression** (primarily neutropenia) is the dose-limiting toxicity. Frequent monitoring of blood counts should be instituted. Patients should not be retreated until neutrophils recover to  $\geq 1,500/\text{mm}^3$  ( $1,000/\text{mm}^3$  for KS patients) and platelets recover to  $\geq 100,000/\text{mm}^3$  ( $\geq 75,000/\text{mm}^3$  for KS patients). In the KS clinical study, the majority of patients were receiving granulocyte colony stimulating factor (G-CSF).

**Severe cardiac conduction abnormalities** have been reported rarely with single agent paclitaxel. If patients develop significant conduction abnormalities during paclitaxel administration, appropriate therapy should be administered and continuous cardiac monitoring should be performed during subsequent therapy with paclitaxel. Hypotension, hypertension, and bradycardia have been observed during paclitaxel administration; patients are usually asymptomatic and generally do not require treatment. Frequent vital sign monitoring, particularly during the first hour of paclitaxel infusion, is recommended. Severe cardiovascular events were observed more frequently in patients with NSCLC than in those with breast or ovarian carcinoma.

A single case of heart failure related to paclitaxel was seen in the AIDS-KS clinical study.

When paclitaxel is used in combination with doxorubicin or trastuzumab for initial treatment of metastatic breast cancer, attention should be placed on the monitoring of cardiac function. When patients are candidates for treatment with paclitaxel in these combinations, they should undergo baseline cardiac assessment including history, physical examination, ECG, echocardiogram, and/or MUGA scan. Cardiac function should be further monitored during treatment (e.g. every three months). Monitoring may help to identify patients who develop cardiac dysfunction and treating physicians should carefully assess the cumulative dose ( $\text{mg}/\text{m}^2$ ) of anthracycline administered when making decisions regarding frequency of ventricular function assessment. When testing indicates deterioration in cardiac function, even asymptomatic, treating physicians should carefully assess the clinical benefits of further therapy against the potential for producing cardiac damage, including potentially irreversible damage. If further treatment is administered, monitoring of cardiac function should be more frequent (e.g. every 1-2 cycles). For more details see Summary of Product Characteristics of trastuzumab or doxorubicin.

Although the occurrence of **peripheral neuropathy** is frequent, the development of severe symptoms is rare. In severe cases, a dose reduction of 20% (25% for KS patients) for all subsequent courses of paclitaxel is recommended. In NSCLC patients and in ovarian cancer patients treated in the first-line setting, the administration of paclitaxel as a three hour infusion in combination with cisplatin, resulted in a greater incidence of severe neurotoxicity than both single agent paclitaxel and cyclophosphamide followed by cisplatin.

**Patients with hepatic impairment** may be at increased risk of toxicity, particularly grade III-IV myelosuppression. There is no evidence that the toxicity of paclitaxel is increased when given as a three-hour infusion to patients with mildly abnormal liver function. When paclitaxel is given as a longer infusion, increased myelosuppression may be seen in patients with moderate to severe hepatic impairment. Patients should be monitored closely for the development of profound myelosuppression (see 4.2). Inadequate data are available to recommend dosage alterations in patients with mild to moderate hepatic impairments (see 5.2).

No data are available for patients with severe baseline cholestasis. Patients with severe hepatic impairment should not be treated with paclitaxel.

Since paclitaxel 6mg/ml infusion contains ethanol (396 mg/ml), consideration should be given to possible CNS and other effects.

Special care should be taken to avoid intra-arterial application of paclitaxel, since in animal studies testing for local tolerance severe tissue reactions were observed after intra-arterial application.

**Pseudomembranous colitis** has been rarely reported including cases in patients who have not been concomitantly treated with antibiotics. This reaction should be considered in the differential diagnosis of cases of severe or persistent diarrhoea occurring during or shortly after treatment with paclitaxel.

Paclitaxel in combination with radiation of the lung, irrespective of their chronological order, may contribute to the development of **interstitial pneumonitis**.

In KS patients, *severe mucositis* is rare. If severe reactions occur, the paclitaxel dose should be reduced by 25%.

*Sexually active fertile female and male* patients should use effective methods of contraception during treatment and up to six months after treatment and one month after treatment for women (see section 4.6). Hormonal contraception is contraindicated in hormone receptor positive tumors.

This product contains macrogolglycerol ricinoleate which may cause severe allergic reactions.

#### 4.5 Interaction with other medicinal products and other forms of interaction

Paclitaxel clearance is not affected by cimetidine premedication.

The recommended regimen of paclitaxel administration for the first-line chemotherapy of ovarian carcinoma is for paclitaxel to be given before cisplatin.

When paclitaxel is given before cisplatin, the safety profile of paclitaxel is consistent with that reported for single-agent use. When paclitaxel was given after cisplatin, patients showed a more profound myelosuppression and an approximately 20% decrease in paclitaxel clearance. Patients treated with paclitaxel and cisplatin may have an increased risk of renal failure as compared to cisplatin alone in gynaecological cancers.

Since the elimination of doxorubicin and its active metabolites can be reduced when paclitaxel and doxorubicin are given closer in time, paclitaxel for initial treatment of metastatic breast cancer should be administered 24 hours after doxorubicin (see 5.2).

The metabolism of paclitaxel is catalysed, in part, by cytochrome P450 isoenzymes CYP2C8 and 3A4 (see 5.2). Clinical studies have demonstrated that CYP2C8-mediated metabolism of paclitaxel, to 6 $\alpha$ -hydroxypaclitaxel, is the major metabolic pathway in humans. Concurrent administration of ketoconazole, a known potent inhibitor of CYP3A4, does not inhibit the elimination of paclitaxel in patients; thus, both medicinal products may be administered together without dosage adjustment. Further data on the potential of drug interactions between paclitaxel and other CYP3A4 substrates/inhibitors are limited. Therefore, caution should be exercised when administering paclitaxel concomitantly with medicines known to inhibit (e.g. erythromycin, fluoxetine, gemfibrozil) or induce (e.g. rifampicin, carbamazepine, phenytoin, phenobarbital, efavirenz, nevirapine) either CYP2C8 or 3A4.

Studies in KS patients, who were taking multiple concomitant medications, suggest that the systemic clearance of paclitaxel was significantly lower in the presence of nelfinavir and ritonavir, but not with indinavir. Insufficient information is available on interactions with other protease inhibitors. Consequently, paclitaxel should be administered with caution in patients receiving protease inhibitors as concomitant therapy.

#### 4.6 Fertility, pregnancy and lactation

Paclitaxel has been shown to be embryotoxic and foetotoxic in rabbits, and to decrease fertility in rats.

There is no information on the use of paclitaxel in pregnant women. As with other cytotoxic drugs, paclitaxel may cause foetal harm, and is therefore contraindicated during pregnancy. Women should be advised to avoid becoming pregnant during therapy with paclitaxel, and to inform the treating physician immediately should this occur.

Pregnancy should be avoided for at least 6 months after treatment.

It is not known whether paclitaxel is excreted in human milk. Paclitaxel is contraindicated during lactation. Breastfeeding should be discontinued for the duration of paclitaxel therapy.

#### 4.7 Effects on ability to drive and use machines

Paclitaxel has not been demonstrated to interfere with this ability. However, it should be noted that the formulation contains alcohol (see 4.4 and 6.1). The ability to drive or to use machines may be decreased due to alcohol content of this medicinal product.

## 4.8 Undesirable effects

The frequency and severity of undesirable effects, unless otherwise mentioned, are generally similar between patients receiving paclitaxel for the treatment of ovarian carcinoma, breast carcinoma, or NSCLC. None of the observed toxicities were clearly influenced by age.

The most frequent significant undesirable effect was **bone marrow suppression**. Severe neutropenia ( $< 500$  cells/mm<sup>3</sup>) occurred in 28% of patients, but was not associated with febrile episodes. Only 1% of patients experienced severe neutropenia for 7 days. Thrombocytopenia was reported in 11% of patients. Three percent of patients had a platelet count nadir  $< 50,000$ /mm<sup>3</sup> at least once while on study.

Anaemia was observed in 64% of patients, but was severe (Hb  $< 5$  mmol/l) in only 6% of patients. Incidence and severity of anaemia is related to baseline haemoglobin status.

**Neurotoxicity**, mainly **peripheral neuropathy**, appeared to be more frequent and severe with a 175 mg/m<sup>2</sup> 3-hour infusion (85% neurotoxicity, 15% severe) than with a 135 mg/m<sup>2</sup> 24-hour infusion (25% peripheral neuropathy, 3% severe) when paclitaxel was combined with cisplatin. In NSCLC patients and in ovarian cancer patients treated with paclitaxel over three hours followed by cisplatin, there is an apparent increase in the incidence of severe neurotoxicity. Peripheral neuropathy can occur following the first course and can worsen with increasing exposure to paclitaxel. Peripheral neuropathy was the cause of paclitaxel discontinuation in a few cases. Sensory symptoms have usually improved or resolved within several months of paclitaxel discontinuation. Pre-existing neuropathies resulting from prior therapies are not a contraindication for paclitaxel therapy.

**Arthralgia or myalgia** affected 60% of patients and was severe in 13% of patients.

**A significant hypersensitivity reaction** with possible fatal outcome (defined as hypotension requiring therapy, angioedema, respiratory distress requiring bronchodilator therapy, or generalised urticaria) occurred in two ( $< 1\%$ ) of patients. Thirty-four percent of patients (17% of all courses) experienced minor hypersensitivity reactions. These minor reactions, mainly flushing and rash, did not require therapeutic intervention nor did they prevent continuation of paclitaxel therapy.

**Injection site reactions** during intravenous administration may lead to localised oedema, pain, erythema, and induration; on occasion, extravasation can result in cellulitis. Skin sloughing and/or peeling has been reported, sometimes related to extravasation. Skin discoloration may also occur. Recurrence of skin reactions at a site of previous extravasation following administration of paclitaxel at a different site, i.e. “recall”, has been reported rarely. A specific treatment for extravasation reactions is unknown at this time.

The table below lists undesirable effects regardless of severity associated with the administration of single agent paclitaxel administered as a three hour infusion in the metastatic setting.

The frequency of undesirable effects listed below is defined using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$ ,  $< 1/10$ ); uncommon ( $\geq 1/1,000$ ,  $< 1/100$ ); rare ( $\geq 1/10,000$ ,  $< 1/1,000$ ); very rare ( $\geq 1/10,000$ ).

### **Infections and infestations:**

*Very common:* infection (mainly urinary tract and upper respiratory tract infections), with reported cases of fatal outcome

*Uncommon:* septic shock

*Rare:* pneumonia, peritonitis, sepsis

### **Blood and the lymphatic system disorders:**

*Very common:* myelosuppression, neutropenia, anaemia, thrombocytopenia, leucopenia, bleeding

*Rare:* febrile neutropenia

*Very rare:* acute myeloid leukaemia, myelodysplastic syndrome

**Immune system disorders:**

*Very common:* minor hypersensitivity reactions (mainly flushing and rash)

*Uncommon:* significant hypersensitivity reactions requiring therapy (e.g., hypotension, angioneurotic oedema, respiratory distress, generalised urticaria, chills, back pain, chest pain, tachycardia, abdominal pain, pain in extremities, diaphoresis and hypertension)

*Rare:* anaphylactic reactions

*Very rare:* anaphylactic shock

**Metabolism and nutrition disorders:**

*Very rare:* anorexia

**Psychiatric disorders:**

*Very rare:* confusional state

**Nervous system disorders:**

*Very common:* neurotoxicity (mainly: peripheral neuropathy)

*Rare:* motor neuropathy (with resultant minor distal weakness)

*Very rare:* autonomic neuropathy (resulting in paralytic ileus and orthostatic hypotension), grand mal seizures, convulsions, encephalopathy, dizziness, headache, ataxia

**Eye disorders:**

*Very rare:* optic nerve and/or visual disturbances (scintillating scotomata), particularly in patients who have received higher doses than recommended

**Ear and labyrinth disorders:**

*Very rare:* ototoxicity, hearing loss, tinnitus, vertigo

**Cardiac disorders:**

*Common:* bradycardia

*Uncommon:* cardiomyopathy, asymptomatic ventricular tachycardia, tachycardia with bigeminy, AV block and syncope, myocardial infarction

*Very rare :* atrial fibrillation, supraventricular tachycardia

**Vascular disorders:**

*Very common:* hypotension

*Uncommon:* hypertension, thrombosis, thrombophlebitis

*Very rare:* shock

**Respiratory, thoracic and mediastinal disorders:**

*Rare:* dyspnoea, pleural effusion, interstitial pneumonia, lung fibrosis, pulmonary embolism, respiratory failure

*Very rare:* cough

**Gastrointestinal disorders:**

*Very common:* nausea, vomiting, diarrhoea, mucosal inflammation

*Rare:* bowel obstruction, bowel perforation, ischaemic colitis, pancreatitis

*Very rare:* mesenteric thrombosis, pseudomembranous colitis, oesophagitis, constipation, ascites, neutropenic colitis

#### **Hepato-biliary disorders:**

*Very rare:* hepatic necrosis, hepatic encephalopathy (both with reported cases of fatal outcome)

#### **Skin and subcutaneous tissue disorders:**

*Very common :* alopecia

*Common:* transient and mild nail and skin changes

*Rare:* pruritus, rash, erythema

*Very rare :* Stevens-Johnson syndrome, epidermal necrolysis, erythema multiforme, exfoliative dermatitis, urticaria, onycholysis (patients on therapy should wear sun protection on hands and feet)

#### **Musculoskeletal, connective tissue and bone disorders :**

*Very common:* arthralgia, myalgia

#### **General disorders and administration site conditions:**

*Common :* injection site reactions (including localised oedema, pain, erythema, induration, on occasion extravasation can result in cellulitis)

*Rare:* asthenia, pyrexia, dehydration, oedema, malaise

#### **Investigations:**

*Common :* severe elevation in AST (SGOT), severe elevation in alkaline phosphatase

*Uncommon:* severe elevation in bilirubin *Rare :* increase in blood creatinine

Breast cancer patients who received paclitaxel in the adjuvant setting following AC experienced more neurosensory toxicity, hypersensitivity reactions, arthralgia/myalgia, anaemia, infection, fever, nausea/vomiting and diarrhoea than patients who received AC alone. However, the frequency of these events was consistent with the use of single agent paclitaxel, as reported above.

#### **Combination treatment**

When administered as a three hour infusion for the first-line chemotherapy of ovarian cancer, neurotoxicity, arthralgia/myalgia, and hypersensitivity were reported as more frequent and severe by patients treated with paclitaxel followed by cisplatin than patients treated with cyclophosphamide followed by cisplatin. Myelosuppression appeared to be less frequent and severe with paclitaxel as a three hour infusion followed by cisplatin compared with cyclophosphamide followed by cisplatin.

For the first line chemotherapy of metastatic breast cancer, neutropenia, anaemia, peripheral neuropathy, arthralgia/myalgia, asthenia, fever, and diarrhoea were reported more frequently and with greater severity when paclitaxel (220 mg/m<sup>2</sup>) was administered as a three-hour infusion 24 hours following doxorubicin (50 mg/m<sup>2</sup>) when compared to standard FAC therapy (5-FU 500 mg/m<sup>2</sup>, doxorubicin 50 mg/m<sup>2</sup>, cyclophosphamide 500 mg/m<sup>2</sup>). Nausea and vomiting appeared to be less frequent and severe with the paclitaxel (220 mg/m<sup>2</sup>) / doxorubicin (50 mg/m<sup>2</sup>) regimen as compared to the standard FAC regimen. The use of corticosteroids may have contributed to the lower frequency and severity of nausea and vomiting in the paclitaxel/doxorubicin arm.

#### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any



suspected adverse reactions via:

HPRA Pharmacovigilance  
Earlsfort Terrace  
IRL - Dublin 2  
Tel: +353 1 6764971  
Fax: +353 1 6762517  
Website: [www.hpra.ie](http://www.hpra.ie)  
e-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie)

## 4.9 Overdose

There is no known antidote for paclitaxel overdosage. The primary anticipated complications of overdosage would consist of bone marrow suppression, peripheral neurotoxicity and mucositis

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group/ATC code: cystostatic agent, L01C D01.

Paclitaxel is a novel antimicrotubule agent that promotes the assembly of microtubules from tubulin dimers and stabilises microtubules by preventing depolymerization. This stability results in the inhibition of the normal dynamic reorganisation of the microtubule network that is essential for vital interphase and mitotic cellular functions. In addition, paclitaxel induces abnormal arrays or bundles of microtubules throughout the cell cycle and multiple asters of microtubules during mitosis.

In the first-line chemotherapy of ovarian carcinoma, the safety and efficacy of paclitaxel were evaluated in two major, randomised, controlled (vs. cyclophosphamide 750 mg/m<sup>2</sup> / cisplatin 75 mg/m<sup>2</sup>) trials. In the Intergroup trial (BMS CA139-209), over 650 patients with stage II<sub>b-c</sub>, III or IV primary ovarian cancer received a maximum of 9 treatment courses of paclitaxel (175 mg/m<sup>2</sup> over three hr) followed by cisplatin (75 mg/m<sup>2</sup>) or control.

The second major trial (GOG-111/BMS CA139-022) evaluated a maximum of six courses of either paclitaxel (135 mg/m<sup>2</sup> over 24 hrs) followed by cisplatin (75 mg/m<sup>2</sup>) or control in over 400 patients with stage III/IV primary ovarian cancer, with a > 1 cm residual disease after staging laparotomy, or with distant metastases. While the two different paclitaxel posologies were not compared with each other directly, in both trials patients treated with paclitaxel in combination with cisplatin had a significantly higher response rate, longer time to progression, and longer survival time when compared with standard therapy. Increased neurotoxicity, arthralgia/myalgia but reduced myelosuppression were observed in advanced ovarian cancer patients administered three-hour infusion paclitaxel/cisplatin as compared to patients who received cyclophosphamide/cisplatin.

In the adjuvant treatment of breast carcinoma, 3121 patients with node positive breast carcinoma were treated with adjuvant paclitaxel therapy or no chemotherapy following four courses of doxorubicin and cyclophosphamide (CALGB 9344, BMS CA 139-223). Median follow-up was 69 months. Overall, paclitaxel patients had a significant reduction of 18% in the risk of disease recurrence relative to patients receiving AC alone (p = 0.0014), and a significant reduction of 19% in the risk of death (p = 0.0044) relative to patients receiving AC alone. Retrospective analyses show benefit in all patient subsets. In patients with hormone receptor negative/ unknown tumours, reduction in risk of disease recurrence was 28% (95%CI: 0.59-0.86). In the patient subgroup with hormone receptor positive tumours, the risk reduction of disease recurrence was 9% (95%CI: 0.78-1.07). However, the design of the study did not investigate the effect of extended AC therapy beyond four cycles. It cannot be excluded on the basis of this study alone that the observed effects could be partly due to the difference in duration of chemotherapy between the two arms (AC 4 cycles; AC + paclitaxel eight cycles). Therefore, adjuvant treatment with paclitaxel should be regarded as an alternative to extended AC therapy.

In a second large clinical study in adjuvant node positive breast cancer with a similar design, 3060 patients were randomized to receive or not four courses of paclitaxel at a higher dose of 225 mg/m<sup>2</sup> following four courses of AC (NSABP B-28, BMS CA139-270). At a median follow-up of 64 months, Paclitaxel patients had a significant reduction of 17% in the risk of disease recurrence relative to patients who received AC alone (p = 0.006); paclitaxel treatment was associated with a reduction in the risk of death of 7% (95% CI: 0.78-1.12). All subset analyses favoured the paclitaxel arm. In this study patients with hormone receptor positive tumour had a reduction in the risk of disease recurrence of 23% (95% CI: 0.6-0.92); in the patient subgroup with hormone receptor negative tumour the risk reduction of disease recurrence was 10% (95% CI: 0.7-1.11).

In the first-line treatment of metastatic breast cancer, the efficacy and safety of paclitaxel were evaluated in two pivotal, phase III, randomised, controlled open-label trials.

In the first study (BMS CA139-278), the combination of bolus doxorubicin (50 mg/m<sup>2</sup>) followed after 24 hours by paclitaxel (220 mg/m<sup>2</sup> by 3-hour infusion) (AT), was compared versus standard FAC regimen (5-FU 500 mg/m<sup>2</sup>, doxorubicin 50 mg/m<sup>2</sup>, cyclophosphamide 500 mg/m<sup>2</sup>), both administered every three weeks for eight courses. In this randomised study, 267 patients with metastatic breast cancer, who had either received no prior chemotherapy or only non-anthracycline chemotherapy in the adjuvant setting, were enrolled. Results showed a significant difference in time to progression for patients receiving AT compared to those receiving FAC (8.2 vs. 6.2 months; p= 0.029). The median survival was in favour of Paclitaxel/doxorubicin vs. FAC (23.0 vs. 18.3 months; p= 0.004). In the AT and FAC treatment arm 44% and 48% respectively received follow-up chemotherapy which included taxanes in 7% and 50% respectively. The overall response rate was also significantly higher in the AT arm compared to the FAC arm (68% vs. 55%). Complete responses were seen in 19% of the Paclitaxel/doxorubicin arm patients vs. 8% of the FAC arm patients. All efficacy results have been subsequently confirmed by a blinded independent review.

In the second pivotal study, the efficacy and safety of the paclitaxel and trastuzumab combination was evaluated in a planned subgroup analysis (metastatic breast cancer patients who formerly received adjuvant anthracyclines) of the study HO648g. The efficacy of trastuzumab in combination with paclitaxel in patients who did not receive prior adjuvant anthracyclines has not been proven. The combination of trastuzumab (4 mg/kg loading dose then 2 mg/kg weekly) and paclitaxel (175 mg/m<sup>2</sup>) 3-hour infusion, every three weeks was compared to single-agent paclitaxel (175 mg/m<sup>2</sup>) three-hour infusion, every three weeks in 188 patients with metastatic breast cancer over expressing HER2 (2+ or 3+ as measured by immunohistochemistry), who had previously been treated with anthracyclines. Paclitaxel was administered every three weeks for at least six courses while trastuzumab was given weekly until disease progression.

The study showed a significant benefit for the paclitaxel /trastuzumab combination in terms of time to progression (6.9 vs. 3.0 months), response rate (41% vs. 17%), and duration of response (10.5 vs. 4.5 months) when compared to paclitaxel alone. The most significant toxicity observed with the paclitaxel /trastuzumab combination was cardiac dysfunction (see 4.8).

In the treatment of advanced NSCLC, paclitaxel 175 mg/m<sup>2</sup> followed by cisplatin 80 mg/m<sup>2</sup> has been evaluated in two phase III trials (367 patients on paclitaxel containing regimens). Both were randomised trials, one compared to treatment with cisplatin 100 mg/m<sup>2</sup>, the other used teniposide 100 mg/m<sup>2</sup> followed by cisplatin 80 mg/m<sup>2</sup> as comparator (367 patients on comparator). Results in each trial were similar. For the primary outcome of mortality, there was no significant difference between the paclitaxel containing regimen and the comparator (median survival times 8.1 and 9.5 months on paclitaxel containing regimens, 8.6 and 9.9 months on comparators). Similarly, for progression-free survival there was no significant difference between treatments. There was a significant benefit in terms of clinical response rate. Quality of life results are suggestive of a benefit on paclitaxel containing regimens in terms of appetite loss and provide clear evidence of the inferiority of paclitaxel containing regimens in terms of peripheral neuropathy (p < 0.008).

In the treatment of AIDS-related KS, the efficacy and safety of paclitaxel were investigated in a non-comparative study in patients with advanced KS, previously treated with systemic chemotherapy. The primary end-point was best tumour response. Of the 107 patients, 63 were considered resistant to liposomal anthracyclines. This subgroup is considered to constitute the core efficacy population. The overall success rate (complete/partial response) after 15 cycles of treatment was 57% (CI 44 - 70%) in liposomal anthracycline-resistant patients. Over 50% of the responses were apparent after the first three cycles. In liposomal anthracycline-resistant patients, the response rates were comparable for patients who had never received a protease inhibitor (55.6%) and those who received one at least two months prior to treatment with paclitaxel (60.9%). The median time to progression in the core population was 468 days (95% CI 257-NE). Median survival could not be computed, but the lower 95% bound was 617 days in core patients.

## 5.2 Pharmacokinetic properties

Following intravenous administration, paclitaxel exhibits a biphasic decline in plasma concentrations.

The pharmacokinetics of paclitaxel were determined following three and 24 hour infusions at doses of 135 and 175 mg/m<sup>2</sup>.

Mean terminal half-life estimates ranged from 3.0 to 52.7 hours, and mean, non-compartmentally derived, values for total body clearance ranged from 11.6 to 24.0 l/hr/m<sup>2</sup>; total body clearance appeared to decrease with higher plasma concentrations of paclitaxel. Mean steady-state volume of distribution ranged from 198 to 688 l/m<sup>2</sup>, indicating extensive extravascular distribution and/or tissue binding. With the 3-hour infusion, increasing doses result in non-linear pharmacokinetics. For the 30% increase in dose from 135 mg/m<sup>2</sup> to 175 mg/m<sup>2</sup>, the C<sub>max</sub> and AUC-∞ values increased 75% and 81%, respectively.

Following an intravenous dose of 100 mg/m<sup>2</sup> given as a three-hour infusion to 19 KS patients, the mean C<sub>max</sub> was 1,530 ng/ml (range 761 - 2,860 ng/ml) and the mean AUC 5,619 ng.hr/ml (range 2,609 - 9,428 ng.hr/ml). Clearance was 20.6 l/h/m<sup>2</sup> (range 11-38) and the volume of distribution was 291 l/m<sup>2</sup> (range 121-638). The terminal elimination half-life averaged 23.7 hours (range 12 - 33).

Inpatient variability in systemic paclitaxel exposure was minimal. There was no evidence for accumulation of paclitaxel with multiple treatment courses.

*In vitro* studies of binding to human serum proteins indicate that 89-98% of drug is bound. The presence of cimetidine, ranitidine, dexamethasone or diphenhydramine did not affect protein binding of paclitaxel.

The disposition of paclitaxel has not been fully elucidated in humans. Mean values for cumulative urinary recovery of unchanged drug have ranged from 1.3 to 12.6% of the dose, indicating extensive non-renal clearance. Hepatic metabolism and biliary clearance may be the principal mechanism for disposition of paclitaxel. Paclitaxel appears to be metabolised primarily by cytochrome P450 enzymes. Following administration of a radiolabelled paclitaxel, an average of 26, 2 and 6% of the radioactivity was excreted in the faeces as 6α-hydroxypaclitaxel, 3'-p-hydroxypaclitaxel, and 6α-3'-p-dihydroxy-paclitaxel, respectively. The formation of these hydroxylated metabolites is catalysed by CYP2C8, -3A4, and both -2C8 and -3A4 respectively. The effect of renal or hepatic dysfunction on the disposition of paclitaxel following a 3-hour infusion has not been investigated formally. Pharmacokinetic parameters obtained from one patient undergoing haemodialysis who received a three-hour infusion of paclitaxel 135 mg/m<sup>2</sup> were within the range of those defined in non-dialysis patients.

In clinical trials where paclitaxel and doxorubicin were administered concomitantly, the distribution and elimination of doxorubicin and its metabolites were prolonged. Total plasma exposure to doxorubicin was 30% higher when paclitaxel immediately followed doxorubicin than when there was a 24-hour interval between drugs.

For use of paclitaxel in combination with other therapies, please consult the Summary of Product Characteristics of cisplatin, doxorubicin or trastuzumab for information on the use of these medicinal products.

### 5.3 Preclinical safety data

The carcinogenic potential of paclitaxel has not been studied. However, paclitaxel is a potential carcinogenic and genotoxic agent, based upon its pharmacodynamic mechanism of action. Paclitaxel has been shown to be mutagenic in both *in vitro* and *in vivo* mammalian test systems.

## 6 PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Citric acid, anhydrous  
Macrogolglycerol ricinoleate  
Ethanol, 96%  
Nitrogen

## 6.2 Incompatibilities

Macroglycerol ricinoleate can result in DEHP [di-(2-ethylhexyl) phthalate] leaching from plasticised polyvinyl chloride (PVC) containers, at levels which increase with time and concentration. Consequently, the preparation, storage and administration of diluted paclitaxel concentrate should be carried out using non-PVC-containing equipment.

## 6.3 Shelf life

Under recommended storage conditions, the unopened product is stable for up to 24 months

For diluted solution, chemical and physical in-use stability has been demonstrated for 24 hours at 25°C. From a microbiological point of view, the diluted product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C, unless reconstitution/dilution has taken place in controlled and validated aseptic conditions.

Diluted solution should be for single use only.

## 6.4 Special precautions for storage

Do not store above 25°C.

Keep vial in the outer carton, in order to protect from light.

After dilution:

Chemical and physical in use stability has been demonstrated for 24 hours at 25°C.

From a microbiological point of view, once opened, the product may be stored for a maximum of 28 days at 25°C. Other in use storage times and conditions are the responsibility of the user.

The diluted solution should not be refrigerated or frozen

## 6.5 Nature and contents of container

Colourless Type I glass vial with fluopolymer-coated chlorobutyl rubber stoppers and aluminium overseal.

Packs of 1 vial containing 5ml, 16.7ml and 50ml of Paclitaxel 6mg/ml Concentrate for Solution for Infusion.

Not all pack sizes may be marketed.

## 6.6 Special precautions for disposal

Handling: as with all antineoplastic agents, caution should be exercised when handling Paclitaxel 6mg/ml Concentrate for Solution for Infusion. Dilution should be carried out under aseptic conditions by trained personnel in a designated area. -

### *Protection instructions for preparation of Paclitaxel solution for infusion*

1. Protective chamber should be used and protective gloves as well as protective gown should be worn. If there is no protective chamber available mouth cover and goggles should be used.
2. Pregnant women or women who may become pregnant, should not handle this product.
3. Opened containers, like injection vials and infusion bottles and used canules, syringes, catheters, tubes, and residuals of cytostatics should be considered as hazardous waste and undergo disposal according to local guidelines for the handling of HAZARDOUS WASTE.

4. Follow the instructions below in case of spillage:- protective clothing should be worn - broken glass should be collected and placed in the container for HAZARDOUS WASTE - contaminated surfaces should be flushed properly with copious amounts of cold water - the flushed surfaces should then be wiped thoroughly and the materials used for wiping should be disposed as HAZARDOUS WASTE
5. In the event of Paclitaxel being in contact with the skin, the area should be rinsed with plenty of running water and then washed with soap and water. In case of contact with mucous membranes, wash the contacted area thoroughly with water. If you have any discomfort, contact a doctor.
5. In case of contact of Paclitaxel with eyes, wash them thoroughly with plenty of cold water. Contact an ophthalmologist immediately.

If unopened vials are refrigerated, a precipitate may form that redissolves with little or no agitation upon reaching room temperature. Product quality is not affected. If the solution remains cloudy or if an insoluble precipitate is noted, the vial should be discarded.

The Chemo-Dispensing Pin device or similar devices with spikes should not be used since they can cause the vial stopper to collapse, resulting in loss of sterile integrity.

Preparation for IV administration: Prior to infusion, Paclitaxel 6mg/ml Concentrate for Solution for Infusion must be diluted, using aseptic techniques, in 0.9% sodium chloride injection, or 5% glucose injection, or 5% glucose and 0.9% sodium chloride injection, or 5% glucose in Ringer's Injection, to a final concentration of 0.3 to 1.2 mg/mL.

Solutions prepared for infusion are stable for 24 hours at 25°C. Following multiple needle entries and product withdrawals, paclitaxel infusion multidose vials maintain microbial, chemical and physical stability for up to 28 days at 25°C. Other in-use storage times and conditions are the responsibility of the user. Diluted solutions should not be refrigerated.

Upon preparation, solutions may show haziness, which is attributed to the formulation vehicle, and is not removed by filtration. Paclitaxel infusion should be administered through an in-line filter with a microporous membrane  $\leq 0.22\mu\text{m}$ . No significant losses in potency have been noted following simulated delivery of the solution through IV tubing containing an in-line filter.

There have been rare reports of precipitation during paclitaxel infusions, usually towards the end of a 24 hour infusion period. Although the cause of this precipitation has not been elucidated, it is probably linked to the supersaturation of the diluted solution. To reduce the precipitation risk, paclitaxel should be used as soon as possible after dilution and excessive agitation, vibration or shaking should be avoided. The infusion sets should be flushed thoroughly before use. During infusion the appearance of the solution should be inspected regularly and the infusion should be stopped if precipitation is present.

To minimise patient exposure to DEHP, which may be leached from plasticised PVC infusion bags, sets, or other medical instruments, diluted paclitaxel solutions should be stored in non-PVC bottles (glass, polypropylene) or plastic bags (polypropylene, polyolefin) and administered through polyethylene-lined administration sets. Use of filter devices (eg. IVEX-2®) which incorporate short inlet and/or outlet plasticised PVC tubing has not resulted in significant leaching of DEHP.

Diluted solution should be for single use only.

Disposal: All items used for preparation, administration or otherwise coming into contact with paclitaxel injection should undergo disposal according to local guidelines for the handling of cytotoxic compounds.

## 7 MARKETING AUTHORISATION HOLDER

Peckforton Pharmaceuticals Ltd,  
Crewe Hall,  
Crewe,  
Cheshire,  
CW1 6UL  
United Kingdom

**8 MARKETING AUTHORISATION NUMBER**

PA946/3/1

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 8 January 2010

**10 DATE OF REVISION OF THE TEXT**

February 2016