

What Andrews Liver Salts do

Andrews Liver Salts quickly and effectively relieve:



Indigestion



Excess Acid



Symptoms of
Over-Indulgence



Upset Stomach



Constipation

How to use Andrews Liver Salts

For oral use only.

Check before you take this medicine if you need a laxative dose every day or you have persistent abdominal pain (tummy ache).

Further information

Keep out of the sight and reach of children.

Do not use after the expiry date (Exp) which refers to the last day of that month. Store below 25°C. Keep the container tightly closed.

Use within 3 months of opening. Check inner seal is intact before first use.

Contents sold by weight and may settle after packaging.

Andrews Liver Salts is an effervescent powder

Active ingredients: Sodium Hydrogen Carbonate 22.6% w/w, Citric Acid 19.5% w/w and Magnesium Sulphate Dihydrate 17.4% w/w.

Other ingredients: Sucrose 40.5% w/w. Carbohydrate content: 2g per 5ml teaspoon.

Andrews is a registered trade mark of the GSK group of companies.

The Product Authorisation holder is GlaxoSmithKline Consumer Healthcare (Ireland) Limited, 12 Riverwalk, CityWest Business Campus, Dublin 24, Ireland and all enquiries should be sent to this address.

The manufacturer is GlaxoSmithKline SA, 28806 Alcala de Henares, Madrid, Spain. Information was last revised in June 2015.

How to take Andrews Liver Salts

How to open:

Lift end of lid. Check inner seal is intact before first use. Break inner seal carefully. Do not take off lid in order to remove inner seal completely.

How to take:

For oral use only.

Adults: For upset stomach, indigestion and over-indulgence, measure one to two level teaspoonfuls into a glass of water and drink.

For constipation, measure two level teaspoonfuls into a glass of water and drink. Take before breakfast or at bedtime.

Drink before the effervescence subsides.

Children aged 6 years or over: Use half the adult dose.

Not suitable for children under 6 years of age except on medical advice.

Close the lid after use.

DO NOT EXCEED THE STATED DOSE

Prolonged, continuous use is not recommended.

If you take too much, you may get diarrhoea.

If this happens, reduce the dose and the effect should go away.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRAs Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2;

Tel: +353 1 6764971; Fax: +353 1 6762517.

Website: www.hpra.ie; E-mail: medsafety@hpra.ie.

By reporting side effects you can help provide more information on the safety of this medicine.

Ask your doctor or pharmacist before you take Andrews Liver Salts:

- if you are taking any prescribed medicines; particularly **tetracycline antibiotics** (e.g. minocycline).
- if you are **pregnant** or are **breastfeeding**, or if you are planning to become pregnant or to breastfeed,
- if you have an intolerance to some sugars, as this product contains sucrose.
- if you are on a **sodium-controlled diet**. Each 5ml spoonful contains **319mg of sodium**.
- if you have **high blood pressure, heart or kidney disease or diabetes** - and avoid frequent use.
- if you suffer from an inflammatory bowel disease or abdominal pain of unknown origin.

Tell your doctor or pharmacist:

- if you need a laxative dose every day or you have persistent abdominal pain (tummy ache).
- if symptoms persist.
- if you experience any unwanted effects.

IMPORTANT:

Do not take if you are allergic to any of the ingredients listed.



Lot:

Exp:

PA 678/43/1

11000010002724

250g



5 000347 069673 >