Rare side-effects:

 bain in your hands or feet (less than 1 in 1,000 people are affected)

· redness of the skin - this usually disappears within a few days

· stomach discomfort and/or pain

Reporting of side-effects

side-effects you can help provide more information on the safety of this side effects directly via HPRA Pharmacovigilance website: By reporting includes any possible side effects not listed in this leaflet. You can also report If you get any side effects, talk to your doctor, pharmacist or nurse. This

D How to store Nicorette Invisi Patch

 Remember that there is still nicotine left in the used patches. Keep this medicine out of the sight and reach of children

· Do not store above 25°C.

Do not use this medicine after the expiry date which is stated on the carton

and sachet after "EXP". The expiry date refers to the last day of that

away from the reach of children and animals. throwing away. Always dispose of used Nicorette Invisi Patches sensibly, inside the empty sachet (or wrapping in a piece of aluminium foil) before • Dispose of Nicorette Invisi Patch as directed by folding it in half and placing

6 Contents of the pack and other information

What Nicorette Invisi Patch contains

The active substance is nicotine.

Each 13.5 cm² transdermal patch contains nicotine 1.75 mg/cm², releasing a

nominal 15 mg nicotine over 16 hours".

side aluminised, both sides siliconised. Aluminium acetylacetonate, Polyethylene terephthalate film (PET) film single copolymer adhesive solution, potassium hydroxide, croscarmellose sodium, triglycerides, medium-chain, basic butyl methacrylate copolymer, acrylic The other ingredients are: polyethylene terephthalate (PET) film,

of a backing layer, a nicotine source layer, a skin contact adhesive layer and Micorette Invisi patch is beige, semi-transparent transdermal patch consisting What Nicorette Invisi Patch looks like and contents of the pack

tace of the patch a siliconised and aluminised release layer, with "nicorette" printed on the top

Each Nicorette Invisi patch is packed in a heat-sealed laminate pouch

Parallel Product Authorisation Holder: supplied in a carton of 7 patches

Road, Dublin 2, Ireland. Lexon Pharmaceuticals (Ireland) Limited, Block 3, Harcourt Centre, Harcourt

Business Park, Clonshaugh, Dublin 17, Ireland. Lexon Pharmaceuticals (Ireland) Limited, Units 22 & 25, Block 4, Port Tunnel **Кераскадед by:**

Andernach, Germany. LTS Lohmann Therapie-Systeme AG, Lohmannstrasse 2, D-56626 Manufactured by:

Nicorette is a registered trademark of GlaxoSmithKline Group of Companies

PPA23176/051/001

Revision date: 31/08/2022

Blind or partially sighted?

Is this leaflet hard to see or read?

Tel: +35318710550 for help.

Phone Lexon Pharmaceuticals (Ireland) Limited,

smokers quit line on 1800 201 203 or visit www.quit.ie. For further information on smoking cessation please ring the national

* if you are pregnant or breast-feeding.

medicine (listed in section 6).

over 16 hours.

Do not use Nicorette Invisi Patch:

det when you try to stop smoking.

6 Contents of the pack and other information

How to store Nicorette Invisi Patch

3 How to use Nicorette Invisi Patch

doctor or pharmacist have told you.

Possible side effects

What is in this leaflet

* if you are a non-smoker or are under 18 years of age.

* if you are allergic to nicotine or any of the other ingredients of this

What you need to know before you use Nicorette Invisi

relieve the unpleasant nicotine withdrawal symptoms and your cravings to

smoke (craving). This indicates that you were dependent on nicotine.

passes into your body through the skin. The nicotine released is sufficient to

When you apply a Nicorette Invisi Patch to the skin, nicotine is released and

absorbing. You may experience unpleasant feelings and a strong desire to

When you stop smoking, your body misses the nicotine that you have been

Nicorette® Invisi 10mg Patch: Each patch releases 10 milligrams of nicotine,

Nicorette® Invisi 15mg Patch: Each patch releases 15 milligrams of nicotine,

combination with either Nicorette 2mg Gum or Nicorette 15mg Inhaler for the

relieve withdrawal symptoms and reduce the cravings for nicotine that you

Nicorette Invisi Patch is a nicotine replacement therapy (NRT). It is used to

What Nicorette Invisi Patch is and what it is used for

What you need to know before you use Nicorette Invisi Patch

This includes any possible side effects not listed in this leaflet. See section

Always take this medicine exactly as described in this leaflet or as your

Read all of this leaflet carefully before you start using this medicine

What Nicorette Invisi Patch is and what it is used for

- If you get any side effects, talk to your doctor or pharmacist.

- Ask your pharmacist if you need more information or advice.

- Keep this leaflet. You may need to read it again.

because it contains important information for you.

treatment of tobacco dependence as part of a stop smoking programme.

Nicorette Invisi 10mg and 15mg Patch can sometimes be used in

aking this medicine. planning to have a baby, ask your doctor or pharmacist for advice before

If you are pregnant or breast-feeding, think you might be pregnant or are

flecainide or adenosine to treat irregular or fast heart beats

imipramine, clomipramine, fluvoxamine to treat depression

* If any of these applies, talk to your doctor or pharmacist.

* if you have persistent indigestion or pains in the chest.

smoking may require the doses of these medicines to be adjusted:

take any other medicines including the following medicines as stopping

Tell your doctor or pharmacist if you are taking, have recently taken or might

Nicorette Invisi Patch should be removed before you have an MRI scan.

* if you have had a stomach ulcer or oesophagitis (inflammation of the

* if you have a skin disorder such as psoriasis, eczema or hives (urticaria)

starting to use Nicorette Invisi Patches as you may find your insulin or

(a fumour of the adrenal gland that can affect blood pressure) – your

Talk to your doctor or pharmacist before using Nicorette Invisi Patch

if you have diabetes - monitor your blood sugar levels more often when

* if you have an overactive thyroid gland or have a phaeochromocytoma

if you have heart disease (including heart attack, disorders of heart rate

occur with this medicine, but is unusual and is both less harmful and easier dependence syndrome after continued use. Transferred dependence can * Transferred dependence: Nicotine in any dose form can result in a

Pregnancy and breastfeeding

ropinirole for Parkinson's disease

tacrine for Alzheimer's disease,

theophylline to treat asthma

* if you have bad circulation.

* if you have high blood pressure.

covering a large area of your skin.

medication requirements after.

* if you have liver or kidney disease.

doctor will have told you this.

or rnytinm, or stroke)

Warnings and Precautions

Micorette Invisi 15mg/16 hours Transdermal Patch

Package leaflet: Information for the User

clozapine or olanzapine for schizophrenia

Other medicines and Nicorette Invisi Patch

to preak than smoking dependence.

are far less than smoking, however you should talk to your doctor or NRT. If you can't manage this, you can use NRT as the risks to your baby If you are pregnant: ideally, you should try to give up smoking without

Ref:23176/051/001/310822/1/F

43 How to use Nicorette Invisi Patch

Always use this medicine exactly as described in this leaflet or as your


pnarmacist it you are not sure. doctor, nurse or pharmacist has told you. Check with your doctor or

Choosing where to apply the patch Follow the instructions below. How to use Nicorette Invisi Patch

2. Avoid placing the patch onto any area of skin that is red, cut or irritated. dry area of hairless skin on the front or side of the chest, upper arm or hip. 1. Before applying your Micorette Invisi Patch, choose a completely clean,

to help avoid irritating that site. 4. It is important that you do not use the same area on two consecutive days as this may prevent it from sticking properly. 3. Do not apply oil or talcum powder to the skin before putting on the patch

How to apply the patch



1. Wash your hands before applying the patch.



with your iingers. aluminium backing away. Avoid touching the sticky surface of the patch 3. Remove the patch from its sachet and then peel one part of the silvery



then peel off the remaining half of the silvery aluminium backing foil. 4. Carefully apply the sticky part of the patch to the chosen area of skin and



5. Press the patch firmly onto the skin with your palm or finger tips.

Invisi Patch to the skin One 2mg gum or the Adults and Apply one new 15mg Removing and disposing of the patch əɓ∀ Dosage of Inhaler Dosage of Invisi Patch 6. Run your fingers around the edge to ensure it sticks firmly.

and over

children

aged 18 years

Patch is not designed to be worn when you go to sleep at night or for 1. The patch should be removed before you go to bed as Nicorette Invisi

placed inside the opened sachet or a piece of aluminium foil. 2. After removal, the patch should be folded in half, sticky side inwards and more than 16 hours each day.

out of reach of children and animals. 3. The patch should then be disposed of carefully in the household rubbish,

Do not give this product to children under 18 years. Children under 18 years

Adults and Children aged 18 years and over

stop using the patches. You should aim to do this within 12 weeks (3 nicotine that you are getting by switching to lower strength patches. You then cravings to smoke. After achieving this you then slowly reduce the amount of The idea is to stop smoking immediately and use the patch to relieve the Stopping Smoking

· Remove 16 hours later which is usually at (usually in the morning). 18 years and over Apply one new patch (of appropriate strenath) to the skin strength and children aged strength) to the skin when you wake up

replace it with another patch. If you lose a patch whilst swimming, bathing or showering you can Do not use more than one patch at a time.

they are out of reach of children and pets. Dispose of the patches carefully after you have removed them, ensuring

or pharmacist. than 6 months in total without asking for help and advice from a doctor Adults aged 18 years and over should not use the patches for longer

patch for the last 4 weeks of treatment. patch and use it for 8 weeks and then step down to the lower strength 10mg If you are a lighter smoker, you should start on the medium strength 15mg If you are a lighter smoker (less than 20 cigarettes a day)

patch and use it for 8 weeks and then step down to the lower strength 15mg If you are a heavier smoker, you should start on the extra strength 25mg If you are a heavier smoker (more than 20 cigarettes a day)

who experience breakthrough cravings, or those who have failed with single a single therapy. However, heavy smokers (more than 20 cigarettes a day), Nicotine replacement therapy often provides sufficient nicotine when used as patch for 2 weeks and then the 10mg patch for the last 2 weeks of treatment.

Combination Therapy Guide

· flushing, high blood pressure

 aching muscles, muscle tenderness or weakness not caused by exercise excessive sweating shortness of breath

(may affect up to 1 in 100 people)

(may affect up to 1 in 10 people)

Very common side-effects:

Other side-effects include:

Pins and needles

(may affect more than 1 in 10 people)

itching - this usually disappears within a few days

• Chest pain, palpitations, an abnormal or fast heart beat

swelling of the nasal passages and back of the throat

dizziness, lightheadedness, blurry vision, nausea

· night time awakening or sleep disturbance

seek medical help immediately:
Swelling of the face, lips, tongue, throat or other parts of the body,

If you experience any of the following, stop using the patch and

may persist or it there is a more severe skin reaction, you should stop using has been. This will usually disappear after a few days. Rarely the reaction

skin reaction. This is usually redness or itching of the skin where the patch

When you use the Micorette Invisi Patch for the first time it may cause a mild

wheezing or difficulty breathing or swallowing

the patches and consult a doctor or pharmacist.

Side-effects of Nicorette Invisi Patch

the mouth in the mouth

lowering of heart rate

book concentration

· irritability or aggression

These effects include:

restlessness

wol gnileef •

anxiety

ntges to smoke (craving)

· increased appetite or weight gain

· month ulcers · constibation

condus sug colds

Uncommon side-effects:

· hives (urticaria) or rash

sickness (vomiting)

feeling sick (nausea)

Common side-effects:


рездасре

- Lash where the patch is applied
- · chest discomfort and pain

silergic reaction

- feeling unwell or tired

Ref:23176/051/001/310822/1/B



Do not exceed 15 pieces of chewing gum or 6 inhaler cartridges per

relieve cravings

Inhaler as required to

Remove the patch 16 hours after application, usually at bedtime.

for the next 4 weeks

when you wake. Do this

Invisi Patch to the skin

when you wake. Do this

Apply one new 10mg

for the first 8 weeks

Use only one Invisi Patch and chew only one piece of gum at a time.

If you use too many Nicorette Invisi Patches

have reduced the amount of nicotine you are taking.

also occur when nicotine is obtained by smoking.

Effects related to stopping smoking (nicotine withdrawal)

disturbances or weakness. in your abdomen, diarrhoea, sweating, headache, dizziness, hearing Patch, you may experience nausea (feeling sick), increased salivation, pain If you have used more than the recommended dosage of Nicorette Invisi

rinse the application site with water. Take this leaflet and the pack with you. Accident and Emergency department immediately. Remove the patch and • If you do get any of these effects contact a doctor or your nearest hospital

If a child has used or swallowed Nicorette Invisi Patches

• Contact a doctor or your nearest hospital Accident and Emergency

this medicine. Take this leaflet and the pack with you. department immediately if a child under 18 years uses, chews or swallows

You may experience unwanted effects because by stopping smoking you

everyone gets them. As many of the effects are due to nicotine, they can

Like all medicines, this medicine can cause side-effects, although not

Possible side effects

Combination Therapy

therapy may benefit from using combination therapy with patch and 2mg gum

Directions for combination use

