

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING -
Label**

**1. NAME OF THE MEDICINAL PRODUCT AND ROUTE(S) OF
ADMINISTRATION**

Lemsip Dry Cough 0.25ml/5ml Oral Solution

2. METHOD OF ADMINISTRATION

For oral use

3. EXPIRY DATE

EXP

4. BATCH NUMBER <DONATION AND PRODUCT CODES>

BN

5. CONTENTS BY WEIGHT, BY VOLUME OR BY UNIT

0.25ml/5ml

6. OTHER

For the symptomatic relief of dry tickly coughs and sore throats

DIRECTIONS

How much to take: Adults & Children 12 and over: 2 x 5ml spoonfuls (10ml) per dose, 3 or 4 times daily. Children 1-12 years: 5ml spoonful per dose, 3 or 4 times daily. Do not give to children aged less than 1 year. If you take too much medicine contact your doctor or pharmacist for advice.

Symptoms of overdose may include headache, nausea, vomiting, diarrhoea, thirst, dizziness, confusion and palpitations. Erosion of the teeth and mouth irritation is possible with citric acid overdose

WARNINGS

For safe use: KEEP OUT OF THE SIGHT AND REACH OF CHILDREN.

If symptoms persist consult your doctor. Do not take this medicine if you are intolerant to sugar. Contains 3.3 g of sucrose and 500 mg invert sugars per 5ml. This should be taken into account in patients with diabetes mellitus. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Warnings: Each 5ml dose contains 0.04mg of ethanol (alcohol).

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

Contains parahydroxybenzoates which may cause allergic reactions (possibly delayed).

Contains benzyl alcohol which may cause allergic reactions. Do not use for more than a

week in young children (less than 3 years old), unless advised by your doctor or pharmacist. Ask your doctor or pharmacist for advice if you are pregnant or breast feeding or have liver or kidney disease. This is because large amounts of benzyl alcohol can build-up in your body and may cause side effects (called "metabolic acidosis").

Side effects: Tell your doctor or pharmacist if you have any side-effects after taking this product. You can also report side effects directly via HPRA Pharmacovigilance, Website: www.hpra.ie

DO NOT EXCEED THE STATED DOSE

INGREDIENTS (Each 5ml dose contains)

Glycerol 0.25ml, honey, sucrose, citric acid, lemon oil terpeneless (benzyl alcohol), ethanol (96%), sodium parabens mixture (sodium ethyl p-hydroxybenzoate (E215), sodium propyl p-hydroxybenzoate and sodium methyl p-hydroxybenzoate (E219) and purified water.

Manufacturer: Reckitt Benckiser Healthcare (UK) Limited, Hull, HU8 7DS, UK and RB NL Brands B.V. Schiphol Blvd 207, 1118 BH, Schiphol, NL.

PA Holder: Reckitt Benckiser Ireland Limited, Dublin 24.

PA979/25/1

Do not use after expiry date shown.

Do not store above 25°C

Date of last revision: January 2022