

Package leaflet: Information for the user
Singulair® Paediatric 5 mg chewable tablets
montelukast

Read all of this leaflet carefully before you or your child start taking this medicine because it contains important information for you or your child.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you or your child only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as your or your child's.
- If you or your child get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Singulair Paediatric is and what it is used for
2. What you need to know before you take Singulair Paediatric
3. How to take Singulair Paediatric
4. Possible side effects
5. How to store Singulair Paediatric
6. Contents of the pack and other information

1. What Singulair Paediatric is and what it is used for

What Singulair Paediatric is

Singulair Paediatric is a leukotriene receptor antagonist that blocks substances called leukotrienes.

How Singulair Paediatric works

Leukotrienes cause narrowing and swelling of airways in the lungs. By blocking leukotrienes, Singulair Paediatric improves asthma symptoms and helps control asthma.

When Singulair Paediatric should be used

Your doctor has prescribed Singulair Paediatric to treat asthma, preventing your asthma symptoms during the day and night.

- Singulair Paediatric is used for the treatment of paediatric patients 6 to 14 years of age who are not adequately controlled on their medication and need additional therapy.
- Singulair Paediatric may also be used as an alternative treatment to inhaled corticosteroids for 6 to 14 year old patients who have not recently taken oral corticosteroids for their asthma and have shown that they are unable to use inhaled corticosteroids.
- Singulair Paediatric also helps prevent the narrowing of airways triggered by exercise.

Your doctor will determine how Singulair Paediatric should be used depending on the symptoms and severity of your or your child's asthma.

What is asthma?

Asthma is a long-term disease.

Asthma includes:

- difficulty breathing because of narrowed airways. This narrowing of airways worsens and improves in response to various conditions.
 - sensitive airways that react to many things, such as cigarette smoke, pollen, cold air, or exercise.
 - swelling (inflammation) in the lining of the airways.
- Symptoms of asthma include: Coughing, wheezing, and chest tightness.

2. What you need to know before you take Singulair Paediatric

Tell your doctor about any medical problems or allergies you or your child has now or has had.

Do not take Singulair Paediatric

- if you or your child is allergic to montelukast or any of the other ingredients of this medicine (listed in section 6)

Warnings and precautions

Talk to your doctor or pharmacist before you or your child take Singulair Paediatric

- If you or your child's asthma or breathing gets worse, tell your doctor immediately.
- Oral Singulair Paediatric is not meant to treat acute asthma attacks. If an attack occurs, follow the instructions your doctor has given you or your child. Always have your inhaled rescue medicine for asthma attacks with you.
- It is important that you or your child take all asthma medications prescribed by your doctor. Singulair Paediatric should not be used instead of other asthma medications your doctor has prescribed for you or your child.
- Any patient on anti-asthma medicines should be aware that if you develop a combination of symptoms such as flu-like illness, pins and needles or numbness of arms or legs, worsening of pulmonary symptoms, and/or

rash, you should consult your doctor.

- You or your child should not take acetyl-salicylic acid (aspirin) or anti-inflammatory medicines (also known as non-steroidal anti-inflammatory drugs or NSAIDs) if they make your asthma worse.

Various neuropsychiatric events (for example behaviour and mood-related changes, depression and suicidality) have been reported in patients of all ages treated with montelukast (see section 4). If you develop such symptoms while taking montelukast, you should contact your doctor.

Children and adolescents

Do not give this medicine to children less than 6 years of age.

There are different form(s) of this medicine available for paediatric patients under 18 years of age based on age range.

Other medicines and Singulair Paediatric

Tell your doctor or pharmacist if you or your child are taking or have recently taken or might take any other medicines including those obtained without a prescription.

Some medicines may affect how Singulair Paediatric works, or Singulair Paediatric may affect how other medicines work.

Tell your doctor if you or your child is taking the following medicines before starting Singulair Paediatric:

- phenobarbital (used for treatment of epilepsy)
- phenytoin (used for treatment of epilepsy)
- rifampicin (used to treat tuberculosis and some other infections)

Singulair Paediatric with food and drink

Singulair Paediatric 5 mg chewable tablets should not be taken immediately with food; it should be taken at least 1 hour before or 2 hours after food.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking Singulair Paediatric.

Pregnancy

Your doctor will assess whether you can take Singulair Paediatric during this time.

Breast-feeding

It is not known if Singulair Paediatric appears in breast milk. You should consult your doctor before taking Singulair Paediatric if you are breast-feeding or intend to breast-feed.

Driving and using machines

Singulair Paediatric is not expected to affect your ability to drive a car or operate machinery. However, individual responses to medication may vary. Certain side effects (such as dizziness and drowsiness) that have been reported with Singulair Paediatric may affect some patients' ability to drive or operate machinery.

Singulair Paediatric 5 mg chewable tablets contain aspartame, sodium and benzyl alcohol

This medicine contains 1.5 mg aspartame in each 5 mg chewable tablet which is equivalent to 0.842 mg phenylalanine. Aspartame is a source of phenylalanine. It may be harmful if you or your child have phenylketonuria (PKU), a rare genetic disorder in which phenylalanine builds up because the body cannot remove it properly.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

This medicine contains benzyl alcohol. Benzyl alcohol may cause allergic reactions.

Ask your doctor or pharmacist for advice if you are pregnant or breast-feeding. This is because large amounts of benzyl alcohol can build-up in your body and may cause side effects (called "metabolic acidosis"). Ask your doctor or pharmacist for advice if you or your child have a liver or kidney disease. This is because large amounts of benzyl alcohol can build-up in the body and may cause side effects (called "metabolic acidosis").

3. How to take Singulair Paediatric

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

- You or your child should take only one chewable tablet of Singulair Paediatric once a day as prescribed by your doctor.
- It should be taken even when you or your child has no symptoms or has an acute asthma attack.

For children 6 to 14 years of age:

The recommended dose is one 5 mg chewable tablet daily to be taken in the evening.

If you or your child are taking Singulair Paediatric, be sure that you or your child do not take any other products that contain the same active ingredient, montelukast.

This medicine is for oral use.

The tablets are to be chewed before swallowing. Singulair Paediatric 5 mg chewable tablets should not be taken immediately with food; it should be taken at least 1 hour before or 2 hours after food.

If you or your child takes more Singulair Paediatric than you should

Contact your doctor immediately for advice.

There were no side effects reported in the majority of overdose reports. The most frequently occurring symptoms reported with overdose in adults and children included abdominal pain, sleepiness, thirst, headache, vomiting, and hyperactivity.

If you forget to take Singulair Paediatric or give Singulair Paediatric to your child

Try to take Singulair Paediatric as prescribed. However, if you or your child misses a dose, just resume the usual schedule of one chewable tablet once daily.

Do not take a double dose to make up for a forgotten dose.

If you or your child stops taking Singulair Paediatric

Singulair Paediatric can treat you or your child's asthma only if you or your child continues to take it. It is important to continue taking Singulair Paediatric for as long as your doctor prescribes. It will help control you or your child's asthma.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

In clinical studies with Singulair Paediatric 5 mg chewable tablets, the most commonly reported side effects (may affect up to 1 in 10 people) thought to be related to Singulair Paediatric were:

- headache

Additionally, the following side effect was reported in clinical studies with Singulair 10 mg film-coated tablets:

- abdominal pain
- These were usually mild and occurred at a greater frequency in patients treated with Singulair than placebo (a pill containing no medication).

Serious side effects

Talk with your doctor immediately if you notice any of the following side effects, which may be serious, and for which you or your child may need urgent medical treatment.

Uncommon: the following may affect up to 1 in 100 people

- allergic reactions including swelling of the face, lips, tongue, and/or throat which may cause difficulty in breathing or swallowing
- behaviour and mood related changes: agitation including aggressive behaviour or hostility, depression
- seizure

Rare: the following may affect up to 1 in 1,000 people

- increased bleeding tendency
- tremor
- palpitations

Very rare: the following may affect up to 1 in 10,000 people

- combination of symptoms such as flu-like illness, pins and needles or numbness of arms and legs, worsening of pulmonary symptoms and/or rash (Churg-Strauss syndrome) (see section 2)
- low blood platelet count
- behaviour and mood related changes: hallucinations, disorientation, suicidal thoughts and actions
- swelling (inflammation) of the lungs
- severe skin reactions (erythema multiforme) that may occur without warning
- inflammation of the liver (hepatitis)

Other side effects while the medicine has been on the market

Very common: the following may affect more than 1 in 10 people

- upper respiratory infection

Common: the following may affect up to 1 in 10 people

- diarrhoea, nausea, vomiting
- rash
- fever
- elevated liver enzymes

Uncommon: the following may affect up to 1 in 100 people

- behaviour and mood related changes: dream abnormalities, including nightmares, trouble sleeping, sleepwalking, irritability, feeling anxious, restlessness
- dizziness, drowsiness, pins and needles/numbness
- nosebleed
- dry mouth, indigestion
- bruising, itching, hives
- joint or muscle pain, muscle cramps
- bedwetting in children
- weakness/tiredness, feeling unwell, swelling

Rare: the following may affect up to 1 in 1,000 people

- behaviour and mood related changes: disturbance in attention, memory impairment, uncontrolled muscle movements

Very rare: the following may affect up to 1 in 10,000 people

- tender red lumps under the skin, most commonly on your shins (erythema nodosum)
- behaviour and mood related changes: obsessive-compulsive symptoms, stuttering

Reporting of side effects

If you or your child get any side effects, talk to your doctor, pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRC Pharmacovigilance. Website www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Singulair Paediatric

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last date of that month.
- Store in the original package in order to protect from light and moisture.
- Store below 30 °C.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Singulair Paediatric contains

- The active substance is montelukast. Each chewable tablet contains montelukast sodium which corresponds to 5 mg of montelukast.
- The other ingredients are: Mannitol (E 421), microcrystalline cellulose, hypolose (E 463), red ferric oxide (E 172), croscarmellose sodium, cherry flavour (containing benzyl alcohol (E 1519)), aspartame (E 951), and magnesium stearate.

What Singulair Paediatric looks like and contents of the pack

5 mg Singulair Paediatric chewable tablets are pink, round, biconvex with SINGULAIR engraved on one side and MSD 275 on the other.

Blisters in packages of 28 tablets.

Parallel Product Authorisation Holder: IMED Healthcare Ltd, Unit 625 Kilshane Avenue, Northwest Business Park, Ballycoolin, Dublin 15, Ireland.

Manufacturer: Merck Sharp & Dohme B.V., Waarderweg 39, 2031 BN Haarlem, The Netherlands.

Repackaged by: Cast Healthcare Ltd, Unit E, The Business Centre, 5-7 Tobermore Road, Draperstown, Magherafelt, BT45 7AG, UK(NI) or IMED Healthcare Ltd, Unit 625 Kilshane Avenue, Northwest Business Park, Ballycoolin, Dublin 15, Ireland.

Information is given by:

The Asthma Society of Ireland, 42 - 43 Amiens Street, Dublin 1, tel: 01 817 8886. Alternatively call the Asthma Adviceline on 1800 44 5464.

(The Asthma Society of Ireland is an independent charity working to conquer asthma and is not associated with IMED Healthcare Ltd.)

This medicine is authorised in the Member States of the European Economic Area under the following names:

Austria, Belgium, Denmark, Finland, France, Greece, Spain, Sweden, Germany, Ireland, Italy, Luxembourg, Netherlands, Portugal
Singulair

This package leaflet was last revised in July 2023.