

Package Leaflet: Information for the user  
Zolepant 20 mg gastro-resistant tablets  
pantoprazole

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet:**

1. What Zolepant is and what it is used for
2. What you need to know before you take Zolepant
3. How to take Zolepant
4. Possible side effects
5. How to store Zolepant
6. Contents of the pack and other information

**1. What Zolepant is and what it is used for**

Zolepant is a selective "proton pump inhibitor", a medicine which reduces the amount of acid produced in your stomach. It is used for treating acid-related diseases of the stomach and intestine.

**Zolepant is used for:**

*Adults and adolescents 12 years of age and above:*

- Treating symptoms (e.g. heartburn, acid regurgitation, pain on swallowing) associated to gastro-oesophageal reflux disease caused by reflux of acid from the stomach.
- Long-term management of reflux oesophagitis (inflammation of the oesophagus accompanied by the regurgitation of stomach acid) and preventing its return.

*Adults:*

- Preventing duodenal and stomach ulcers caused by non-steroidal anti-inflammatory drugs (NSAIDs, for example, ibuprofen) in patients at risk who need to take NSAIDs continuously.

**2. What you need to know before you take Zolepant**

**Do not take Zolepant:**

- If you are allergic to pantoprazole, sorbitol or any of the other ingredients of this medicine (listed in section 6)
- If you are allergic to medicines containing other proton pump inhibitors.

**Warnings and Precautions**

**Talk to your doctor or pharmacist before taking Zolepant**

- If you have severe liver problems. Please tell your doctor if you have ever had problems with your liver. He will check your liver enzymes more frequently, especially when you are taking Zolepant as a long-term treatment. In the case of a rise of liver enzymes the treatment should be stopped.
- If you need to take medicines called NSAIDs continuously and receive Zolepant because you have an increased risk of developing stomach and intestinal complications. Any increased risk will be assessed according to your own personal risk factors such as your age (65 years old or more), a history of stomach or duodenal ulcers or of stomach or intestinal bleeding.
- If you have reduced body stores or risk factors for reduced vitamin B12 and receive pantoprazole long-term treatment. As with all acid reducing agents, pantoprazole may lead to a reduced absorption of vitamin B12.
- If you are taking HIV protease inhibitors such as atazanavir (for the treatment of HIV-infection) at the same time as pantoprazole, ask your doctor for specific advice.
- Taking a proton pump inhibitor like pantoprazole, especially over a period of more than one year, may slightly increase your risk of fracture in the hip, wrist or spine.
- Tell your doctor if you have osteoporosis or if you are taking corticosteroids (which can increase the risk of osteoporosis).
- If you are on Zolepant for more than three months it is possible that the levels of magnesium in your blood may fall. Low levels of magnesium can be seen as fatigue, involuntary muscle contractions, disorientation, convulsions, dizziness or increased heart rate. If you get any of these symptoms, please tell your doctor promptly. Low levels of magnesium can also lead to a reduction in potassium or calcium levels in the blood. Your doctor may decide to perform regular blood tests to monitor your levels of magnesium.
- If you have ever had a skin reaction after treatment with a medicine similar to Zolepant that reduces stomach acid.
- If you get a rash on your skin, especially in areas exposed to the sun tell your doctor as soon as you can, as you may need to stop your treatment with Zolepant. Remember to also mention any other ill-effects like pain in your joints.
- If you are due to have a specific blood test (Chromogranin A).

**Tell your doctor immediately** if you notice any of the following symptoms:

- an unintentional loss of weight
- vomiting, particularly if repeated
- vomiting blood; this may appear as dark coffee grounds in your vomit
- you notice blood in your stools; which may be black or tarry in appearance
- difficulty in swallowing or pain when swallowing
- you look pale and feel weak (anaemia)
- chest pain
- stomach pain
- severe and/or persistent diarrhoea, as Zolepant has been associated with a small increase in infectious diarrhoea.

Your doctor may decide that you need some tests to rule out malignant disease because pantoprazole also alleviates the symptoms of cancer and could cause delay in diagnosing it. If your symptoms continue in spite of your treatment, further investigations will be considered.

If you take Zolepant on a long-term basis (longer than 1 year) your doctor will probably keep you under regular surveillance. You should report any new and exceptional symptoms and circumstances whenever you see your doctor.

**Children and adolescents**

These tablets are not recommended for use in children below 12 years.

**Other medicines and Zolepant**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. Zolepant may influence the effectiveness of other medicines, so tell your doctor if you are taking

- Medicines such as ketoconazole, itraconazole and posaconazole (used to treat fungal infections) or erlotinib (used for certain types of cancer) because Zolepant may stop these and other medicines from working properly.
- Warfarin and phenprocoumon, which affect the thickening, or thinning of the blood. You may need further checks.
- Medicines used to treat HIV-infection, such as atazanavir.
- Methotrexate (used to treat rheumatoid arthritis, psoriasis, and cancer) - if you are taking methotrexate your doctor may temporarily stop your Zolepant treatment because pantoprazole can increase levels of methotrexate in the blood.
- Fluvoxamine (used to treat depression and other psychiatric diseases) - if you are taking fluvoxamine your doctor may reduce the dose.
- Rifampicin (used to treat infections)
- St John's wort (*Hypericum perforatum*)(used to treat mild depression).

**Zolepant with food and drink**

Take the tablets 1 hour before a meal without chewing or breaking them and swallow them whole with some water.

**Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. There are no adequate data from the use of pantoprazole in pregnant women. Excretion into human milk has been reported. You should use this medicine, only if your doctor considers the benefit for you greater than the potential risk for your unborn child or baby.

**Driving and using machines**

Zolepant has no or negligible influence on the ability to drive and use machines. If you experience side effects like dizziness or disturbed vision, you should not drive or operate machines.

**Zolepant contains sorbitol and sodium.**

This medicine contains 18 mg sorbitol in each tablet. This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

**3. How to take Zolepant**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

**When and how should you take Zolepant**

Take the tablets 1 hour before a meal without chewing or breaking them and swallow them whole with some water.

Unless told otherwise by your doctor, the recommended dose is:

*Adults and adolescents 12 years of age and above:*

**To treat symptoms (e.g. heartburn, acid regurgitation, pain on swallowing) associated to gastro-oesophageal reflux disease**

The recommended dose is one tablet a day. This dose usually brings relief within 2-4 weeks - at most after another 4 weeks. Your doctor will tell you how long to continue taking the medicine. After this any recurring symptoms can be controlled by **taking one tablet daily**, when required.

**For long-term management and for preventing the return of reflux oesophagitis**

The recommended dose is one tablet a day. If the illness returns, your doctor can double the dose, in which case you can use Zolepant 40 mg tablets instead, one a day. After healing, you can reduce the dose back again to one tablet 20 mg a day.

*Adults:*

**To prevent duodenal and stomach ulcers in patients who need to take NSAIDs continuously**

The recommended dose is one tablet a day.

**Special patient groups:**

- If you suffer from severe liver problems, you should not take more than one 20 mg tablet a day.

**Use in children and adolescents**

Children below 12 years.  
These tablets are not recommended for use in children below 12 years.

**If you take more Zolepant than you should**

Tell your doctor or pharmacist. There are no known symptoms of overdose.

**If you forget to take Zolepant**

Do not take a double dose to make up for a forgotten dose. Take your next, normal dose at the usual time.

**If you stop taking Zolepant**

Do not stop taking these tablets without first talking to your doctor or pharmacist. If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

24LF01695PD

Direction of fibre

Direction of reading PhC

70 mm ± 0,5 mm

70 mm ± 0,5 mm

Direction of reading PhC

Direction of reading PhC

130 mm ± 0,5 mm

70 mm ± 0,5 mm

Direction of reading PhC

70 mm ± 0,5 mm

Direction of reading PhC

70 mm ± 0,5 mm

Direction of reading PhC



Pharma code

Pharma code

#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**If you get any of the following side effects, stop taking these tablets and tell your doctor immediately, or contact the casualty department at your nearest hospital:**

##### **Serious allergic reactions (frequency rare (may affect up to 1 in 1,000 people)):**

- swelling of the tongue and/or throat,
- difficulty in swallowing,
- hives (nettle rash),
- difficulties in breathing,
- allergic facial swelling (Quincke's oedema / angioedema),
- severe dizziness with very fast heartbeat and heavy sweating.

##### **Serious skin conditions (frequency not known (frequency cannot be estimated from the available data)):**

- blistering of the skin and rapid deterioration of your general condition,
- erosion (including slight bleeding) of eyes, nose, mouth/lips or genitals (Stevens-Johnson-Syndrome, Lyell-Syndrome, Erythema multiforme)
- sensitivity to light.

##### **Other serious conditions (frequency not known (frequency cannot be estimated from the available data)):**

- yellowing of the skin or whites of the eyes (severe damage to liver cells, jaundice)
- fever
- rash
- enlarged kidneys sometimes with painful urination and lower back pain (serious inflammation of the kidneys, possibly progressing to kidney failure).

If you are on pantoprazole for more than three months it is possible that the levels of magnesium in your blood may fall. Low levels of magnesium can be seen as fatigue, involuntary muscle contractions, disorientation, convulsions, dizziness, increased heart rate. If you get any of these symptoms, please tell your doctor promptly. Low levels of magnesium can also lead to a reduction in potassium or calcium levels in the blood. Your doctor may decide to perform regular blood tests to monitor your levels of magnesium.

##### **Frequency Rare: (may affect up to 1 in 1,000 people)**

- severe reduction in number of white blood cells which makes infections more likely. Report any signs of sudden fever, sore throat, extreme tiredness, which may be signs of infection.

##### **Very rare (may affect up to 1 in 10,000 people)**

- A reduction in the number of blood platelets, which may cause you to bleed or bruise more than normal. Please report any unexplained bruising or bleeding to your doctor.

##### **Other side effects are:**

##### **Common (may affect up to 1 in 10 people)**

- Benign polyps in the stomach.

##### **Uncommon (may affect up to 1 in 100 people)**

- Headache;
- diarrhoea;
- vomiting;
- constipation;
- abdominal pain and discomfort;
- itching;
- feeling weak, exhausted or generally unwell;
- dizziness;
- feeling sick;
- bloating and flatulence (wind);
- dry mouth;
- skin rash, exanthema, eruption;
- fracture of the hip, wrist or spine;
- sleep disorders.

##### **Rare (may affect up to 1 in 1,000 people)**

- Disturbances in vision such as blurred vision;
- pain in the joints;
- weight changes;
- swelling of the extremities (peripheral oedema);
- allergic reactions;
- breast enlargement in males;
- hives;
- muscle pains;
- raised body temperature;
- depression;
- taste disorders.

##### **Very rare (may affect up to 1 in 10,000 people)**

- Disorientation.

##### **Not known (frequency cannot be estimated from the available data)**

- Hallucination, confusion (especially in patients with a history of these symptoms);
- decreased sodium level in blood;
- decreased potassium level in blood;
- muscle spasm;
- inflammation in the large bowel, that causes persistent watery diarrhoea.
- decreased calcium level in blood;
- tingling or numbness in the hands or feet;
- rash, possibly with pain in the joints;

##### **Side effects identified through blood tests:**

##### **Uncommon (may affect up to 1 in 100 people)**

- An increase in liver enzymes.

##### **Rare (may affect up to 1 in 1,000 people)**

- An increase in bilirubin;
- increased fats in the blood;
- severe reduction in number of white blood cells which makes infections more likely.

##### **Very rare (may affect up to 1 in 10,000 people)**

- A reduction in the number of blood platelets, which may cause you to bleed or bruise more than normal;
- a reduction in the number of white blood cells, which may lead to more frequent infections;
- severe reduction in blood cells which can cause weakness, bruising or make infections more likely.

##### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRÁ Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: [www.hpra.ie](http://www.hpra.ie); e-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

#### 5. How to store Zolepant

Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiry date which is stated on the packaging after EXP. The expiry date refers to the last day of that month. This medicine does not require any special temperature storage conditions.

Blister pack: Store in the original package in order to protect from moisture.

Container: Keep the container tightly closed in order to protect from moisture.

After first opening of the container, the product should be used within 3 months.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

#### 6. Contents of the pack and other information

##### **What Zolepant contains**

- The active substance is pantoprazole. Each gastro-resistant tablet contains 20 mg pantoprazole (as pantoprazole sodium sesquihydrate).
- The other ingredients are mannitol, crospovidone (type A, type B), sodium carbonate, sorbitol (E420), calcium stearate in the tablet core and hypromellose, povidone (K25), titanium dioxide (E171), yellow iron oxide (E172), propylene glycol, methacrylic acid - ethyl acrylate copolymer, sodium laurilsulfate, polysorbate 80, macrogol 6000 and talc in the film-coating.

##### **What Zolepant looks like and contents of the pack**

The 20 mg gastro-resistant tablets are light brownish yellow, oval, slightly biconvex tablets.

##### **Pack sizes:**

Cartons of 7, 14, 15, 20, 28, 30, 50, 50 x 1, 56, 60, 84, 90, 98, 100, 100 x 1, 112 and 140 gastro-resistant tablets in blister packs. A plastic container of 100 and 250 gastro-resistant tablets. Not all pack sizes may be marketed.

##### **Marketing Authorisation Holder**

Pinewood Laboratories Ltd., Ballymacarbry, Clonmel, Co. Tipperary, Ireland.

##### **Manufacturer**

KRKA d.d., Novo Mesto, Smarjeska Cesta 6, 8501 Novo Mesto, Slovenia.  
TAD Pharma GmbH, Heinz-Lohmann-Str.5, D-27472 Cuxhaven, Germany.



##### **Product Authorisation Number**

PA 281/140/1

##### **This medicinal product is authorised in the Member States of the EEA under the following names:**

AT	PANTOPRAZOL TEVA	BE	PANTOPRAZOLE
DK	PANTOPRAZOL KRKA	FI	PANTOPRAZOLE KRKA
FR	PANTOPRAZOLE TEVA	DE	PANTOPRAZOL TAD
IE	ZOLEPANT	IT	PANTOPRAZOLE KRKA
UK	PANTOPRAZOLE	NL	PANTOPRAZOL
NO	PANTOPRAZOL KRKA	SE	PANTOPRAZOL KRKA
ES	PANTOPRAZOL TEVA	PT	PANTOPRAZOLE KRKA
PL	NOLPAZA	SK	NOLPAZA
LV	NOLPAZA	EE	NOLPAZA
LT	NOLPAZA	CZ	NOLPAZA

This leaflet was last revised in September 2019



Pharma code

KRKA article number