

**Canesten 200mg Pessary
Clotrimazole 200mg**

Read all of this leaflet carefully before you start using this medicine because it contains important information for you. Always use this medicine exactly as described in this leaflet or as your doctor, pharmacist or nurse have told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See Section 4.
- You must contact a doctor if your symptoms worsen or do not improve in 7 days.

WHAT IS IN THIS LEAFLET

1. What is Canesten 200mg Pessary and what is it used for?
2. What you need to know before you use Canesten 200mg Pessary
3. How to use Canesten 200mg Pessary
4. Possible side effects
5. How to store Canesten 200mg Pessary
6. Contents of the pack and other information

1. WHAT IS CANESTEN® 200mg PESSARY AND WHAT IS IT USED FOR?

Canesten 200mg Pessary is indicated for the treatment of vaginal thrush. It is inserted into the vagina for treatment at the site of infection. Canesten 200mg pessaries are used to treat adults and children aged 12 years and over.

Only use this product if you have been previously diagnosed by your doctor as having vaginal thrush.

The active substance in Canesten 200mg Pessary is clotrimazole. Clotrimazole belongs to a group of medicines called azoles and is an antifungal agent which fights the cause of infections such as vaginal thrush.

2. WHAT YOU NEED TO KNOW BEFORE YOU USE CANESTEN® 200mg PESSARY

Do not use Canesten® 200mg Pessary:

- If you are allergic (hypersensitive) to clotrimazole or any of the other ingredients of Canesten 200mg Pessary (listed in section 6).
- During your period as it may be less effective.

Warnings and Precautions:

Talk to your doctor or pharmacist before using **Canesten® 200mg Pessary**

- You are unsure whether you have thrush or this is the first time you have had these symptoms.
- You have had more than two infections of thrush in the last six months.

- You or your partner have ever had a sexually transmitted disease.
- You are aged under 12 or over 60.
- You have ever had an allergic reaction to Canesten or any of the other ingredients (listed in section 6)
- You have any of the following symptoms:
 - Irregular vaginal bleeding.
 - Abnormal vaginal bleeding or a blood-stained discharge.
 - Ulcers, blisters or sores of the vagina or vulva.
 - Redness, irritation or swelling associated with recent use of Canesten 200mg Pessary.
 - Lower abdominal pain.
 - Back pain.
 - Shoulder pain.
 - Pain or difficulty in passing urine.
 - Fever or chills.
 - Feeling sick or vomiting.
 - Diarrhoea.
 - A foul smelling discharge from the vagina.

This is because Canesten 200mg Pessary may not be the right treatment for you.

Taking other medicines:

If you are taking tacrolimus or sirolimus (immunosuppressant medicine), then you should tell your doctor, because he or she will monitor the levels of this medicine in your blood.

Special precautions:

Avoid sexual intercourse while using this product, as the infection could be transferred to your partner. As with other pessaries, this product may reduce the effectiveness of rubber contraceptives, such as condoms or diaphragms.

Consequently, you should use alternative precautions for at least five days after using this product. Do not use tampons, spermicides or other vaginal products while using this product.

Pregnancy and breastfeeding:

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine. Canesten 200mg Pessary can be used in pregnancy and breastfeeding, but only under the supervision of your doctor or midwife. You must follow his/her instructions for use.

Use of the applicator is not advised during pregnancy.

3. HOW TO USE CANESTEN® 200mg PESSARY

Always use this medicine exactly as described in this leaflet or as your pharmacist or doctor has told you. Check with your pharmacist if you are not sure.

If Canesten 200mg Pessary has been prescribed for you by your doctor, follow any instructions he/she may have given you. If you purchased this product without a prescription, follow these directions closely:


The applicator should be used to insert the pessary as high as possible into the vagina, preferably before going to sleep on **three consecutive nights** for convenient and comfortable treatment.

Wash your hands before handling the applicator and the foil blister pack and again afterwards when you have used the applicator.


1. Remove the applicator from the packaging and pull out the plunger and red lock from the applicator body.



2. Remove the pessary from the foil blister pack and place into the open end of the applicator body with the curved edge of the pessary facing down. Push the plunger and lock into the applicator body until you feel a click.



3. Once you have felt the click, remove the lock from the plunger and then carefully insert the applicator as deep as is comfortable into the vagina (this is easiest when lying on your back with your knees bent up) up to the patterned grip zone, without pressing on the plunger.



4. Hold the applicator body at the patterned grip zone and slowly push the plunger all the way until it stops, so that the pessary is deposited in the vagina.



5. Remove the applicator. After use, disassemble all the components for cleaning and wash thoroughly in warm (not boiling) soapy water, rinse for 30 seconds and wipe the parts carefully. Store it in a clean and safe place.



For re-use, ensure the red lock is replaced on the plunger and repeat the process from step 2. After completing the treatment, dispose of the applicator and red lock in a safe place, out of the reach of children. The applicator cannot be flushed down the toilet.

Since the pessary dissolves in the vagina, it may be helpful to wear a panty liner because it is quite common to notice a white chalky residue after using the pessary. This does not mean that the treatment has not worked. However, if you notice pieces of undissolved pessary, speak to your doctor or a pharmacist, as the treatment may not have worked properly. If your vagina and vulva are itchy and sore, Canesten Cream should also be used. The symptoms of thrush should disappear within three days of treatment. If no improvement is seen after seven days you must tell your doctor. If the infection returns after seven days you may use one further treatment, but if you have more than two infections within two months you should see the doctor.

Canesten® 200mg Pessary is for use in the vagina only:

Do not put the pessary in your mouth or swallow it.

Avoid contact with the eyes. If the pessary is swallowed accidentally, tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital.

If you forget to use Canesten® 200mg Pessary:

Canesten 200mg Pessary should be used for three consecutive nights. However, if you forget to insert a pessary, follow the advice given below:

When you remember, continue with your course of treatment. If you have forgotten for more than one day, your infection may not be fully treated, so if symptoms persist upon completion of this course, see your doctor.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Canesten 200mg Pessary can cause side effects, although not everybody gets them.

As with all medicines, some people may be allergic to the pessary. If you are allergic, a reaction will occur soon after you have used the medicine. If you experience an allergic reaction or the redness, burning, pain, itching or swelling get worse, stop using this product and tell your doctor straight away or contact the Accident and Emergency Department of your nearest hospital.

Signs of an allergic reaction may include:

- Rash.
- Swallowing or breathing problems.
- Swelling of your lips, face, throat or tongue.
- Weakness, feeling dizzy or faint.
- Nausea.
- Low blood pressure.

Rarely, immediately after you insert the pessary you might experience:

- Mild burning or irritation.

Other potential side effects that have been reported:

- Discomfort of the vagina or vulva, redness, swelling, burning, bleeding, peeling skin in the genital area, irritation, itching, rash, vaginal discharge, pain (e.g. abdominal or pelvic).

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE CANESTEN® 200mg PESSARY

Keep this medicine out of the sight and reach of children.

Do not store above 25°C.

Do not use Canesten 200mg Pessary after the expiry date which is stated at one end of the carton and on the foil blister strip of the pessaries. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to dispose of medicines you no longer use. These measures will help to protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Canesten® 200mg Pessary contains:

- The active substance is clotrimazole at a strength of 200mg.
- The other ingredients are calcium lactate pentahydrate, maize starch, crospovidone, colloidal anhydrous silica, lactic acid, lactose monohydrate, magnesium stearate, hypromellose and microcrystalline cellulose.

What Canesten® 200mg Pessary looks like and contents of the pack:

Canesten 200mg Pessary contains three white pessaries held inside a foil blister pack and one applicator for insertion of the pessary into the vagina.

Marketing Authorisation Holder:

Bayer Ltd,
The Atrium,
Blackthorn Road,
Dublin 18.

Manufacturer:

GP Grenzach Produktions GmbH, Germany.

Remember: If you have any doubts about using Canesten® 200mg Pessary correctly, seek the advice of your doctor or pharmacist.

Further information about vaginal thrush:

Vaginal thrush (candidiasis) is a common infection that most women suffer from at some time in their lives and is not caused by lack of personal hygiene. Thrush is caused by a yeast (fungus) called Candida which lives harmlessly in the vagina and other parts of the body, without you even noticing it. However, the natural balance that keeps Candida under control can be upset by many factors such as hormonal changes (menstruation, contraceptive pill, pregnancy, menopause), poor health, antibiotics, perfumed soaps, bath additives and tight clothing.

If the natural pH balance is altered, the level of yeast increases and can develop into a thrush infection causing any of the following symptoms: persistent burning and/or itching around the vagina and vulva, redness, swelling and soreness of the tissues of the vagina and vulva and a whitish, odourless discharge from the vagina. Not everybody who has thrush has all these symptoms; you may have only one of them.

How to avoid future recurrences:

Wear cotton knickers and loose clothing.

Wash daily.

After going to the toilet, wipe yourself from the front to back as a thrush infection may be transferred from the bowel.

Change your sanitary protection regularly.

Try to avoid wearing tights, nylon knickers and close fitting jeans.

Try to avoid washing with perfumed soaps or using vaginal deodorants.

Do not wash or rub yourself hard with sponges or flannels and avoid hot baths with strong perfumed oils.

If you are still worried or have any questions about the symptoms or the treatment of thrush, do not hesitate to ask your doctor or pharmacist for advice.

This leaflet was last approved XXXX 2022.

Canesten is a registered trademark of Bayer AG, Germany.