

(GSK logo)

**Package leaflet: Information for the user**

**Zovirax® 30 mg/g Eye Ointment**  
**aciclovir**

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**Whats in this leaflet:**

- 1 What Zovirax is and what it is used for
- 2 What you need to know before you use Zovirax
- 3 How to use Zovirax
- 4 Possible side effects
- 5 How to store Zovirax
- 6 Contents of the pack and other information

## **1 What Zovirax is and what it is used for**

Zovirax 30 mg/g Eye Ointment (called 'Zovirax' in this leaflet) contains a medicine called aciclovir. This belongs to a group of medicines called antivirals. It is used to treat eye infections caused by the Herpes simplex virus. It works by stopping the growth of viruses on the front of your eyeball (cornea).

## **2 What you need to know before you use Zovirax**

**Do not use Zovirax:**

- if you are allergic to aciclovir or valaciclovir or any of the other ingredients of this medicine (listed in Section 6).

If you are not sure, talk to your doctor or pharmacist before using Zovirax.

**Warnings and precautions**

Talk to your doctor or pharmacist before using Zovirax if:

- you wear contact lenses. Stop wearing them while using Zovirax.

If you are not sure, talk to your doctor or pharmacist before using Zovirax.

**Other medicines and Zovirax**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes medicines obtained without prescription, including herbal medicines.

**Zovirax with food and drink**

Food and drink should not affect the absorption of your medicine.

**Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine

**Driving and using machines**

Your vision may be blurred for 5 to 10 minutes after applying Zovirax. Wait until you can see clearly again before driving or using any tools or machines.

## **3 How to use Zovirax**

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

### Adults and children

- The usual dose is 1cm (about ½ an inch) of Zovirax applied to the infected eye five times a day. You should space each application of Zovirax by four hours. Suggested times are: **7am, 11am, 3pm, 7pm and 11pm.**
- You should carry on using Zovirax for at least 3 days after your eye has got better.
- Your vision may be blurred for 5 to 10 minutes after applying Zovirax. Wait until you can see clearly again before driving or using any tools or machines.

### How to apply Zovirax to your eye



- 1 Wash your hands.
- 2 Use your finger to gently pull down the lower eyelid of your infected eye.
- 3 Tilt your head slightly backwards and look up.
- 4 Apply 1cm (about ½ an inch) of Zovirax to the inside of your lower eyelid. **Try to avoid the tip of the tube touching any part of your eye.**
- 5 Close your eye for 30 seconds.
- 6 Wash your hands after applying Zovirax.

### If you use more Zovirax than you should

Talk to your doctor or pharmacist if you use too much Zovirax. Zovirax eye ointment is for use in the eye, if you swallow any talk to your doctor or pharmacist.

### If you forget to use Zovirax

- If you forget to use Zovirax, use it as soon as you remember. However, if it is nearly time for your next application, skip the missed application.
- Do not use a double application to make up for a forgotten application.

## 4 Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. The following side effects may happen with this medicine:

### Allergic reactions (may affect up to 1 in 10,000 people)

If you have an allergic reaction, **stop using Zovirax and see a doctor straight away.** The signs may include:

- rash, itching or hives on your skin
- swelling of your face, lips, tongue or other parts of your body
- shortness of breath, wheezing or trouble breathing
- unexplained fever (high temperature) and feeling faint, especially when standing up

Other side effects include:

### Very common (may affect more than 1 in 10 people)

- burning or irritation, photophobia/light sensitivity or feeling as if there is something in your eye (superficial punctate keratopathy)

### Common (may affect up to 1 in 10 people)

- mild stinging in your eye straight after using Zovirax
- swollen, runny eyes (conjunctivitis)

### Rare (may affect up to 1 in 1,000 people)

- swollen, red, irritated, itchy eyelids (blepharitis)

### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL – Dublin 2; Tel: +353 1 6764971; Fax +353 1 6762517. Website: [www.hpra.ie](http://www.hpra.ie); E-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie).

By reporting side effects you can help provide more information on the safety of this medicine.

## **5 How to store Zovirax**

- Keep this medicine out of the sight and reach of children.
- Do not store above 25°C.
- Keep the tube tightly closed when you are not using Zovirax.
- Do not use Zovirax if more than one month has passed since you first opened the tube.
- Do not use this medicine after the expiry date which is stated on the carton and tube after EXP. The expiry date refers to the last day of that month.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment

## **6 Contents of the pack and other information**

### **What Zovirax 30 mg/g Eye Ointment contains**

- The active substance is aciclovir.
- Each gram of ointment contains 30 mg aciclovir equivalent to 3% w/w aciclovir.
- The other ingredient is white petrolatum.

### **What Zovirax looks like and contents of the pack**

Each carton contains a 4.5 g tube of Zovirax. The tube has a screw cap and contains an off-white ointment.

### **Marketing authorisation holder and manufacturer**

#### **Marketing authorisation holder**

GlaxoSmithKline (Ireland) Ltd, 12 Riverwalk, Citywest Business Campus, Dublin 24

#### **Manufacturer**

Glaxo Operations UK Ltd, Barnard Castle, Durham, DL12 8DT, UK

**This leaflet was last revised in July 2015.**

Zovirax is a registered trade mark of the GSK group of companies.

©2015 GSK group of companies. All rights reserved.