

Package leaflet: Information for the user

HBVAXPRO® 10 micrograms, suspension for injection in pre-filled syringe Hepatitis B vaccine (rDNA)

Read all of this leaflet carefully before you are vaccinated because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What HBVAXPRO 10 micrograms is and what it is used for
2. What do you need to know before you receive HBVAXPRO 10 micrograms
3. How HBVAXPRO 10 micrograms is given
4. Possible side effects
5. How to store HBVAXPRO 10 micrograms
6. Contents of the pack and other information

1. What HBVAXPRO 10 micrograms is and what it is used for

This vaccine is indicated for active immunisation against hepatitis B virus infection caused by all known subtypes in individuals 16 years of age or more considered at risk of exposure to hepatitis B virus.

It can be expected that hepatitis D will also be prevented by immunisation with HBVAXPRO as hepatitis D does not occur in the absence of hepatitis B infection.

The vaccine will not prevent infection caused by other agents such as hepatitis A, hepatitis C and hepatitis E and other pathogens known to infect the liver.

2. What you need to know before you receive HBVAXPRO 10 micrograms

Do not use HBVAXPRO 10 micrograms

- if you are allergic to hepatitis B surface antigen or to any of the other ingredients of HBVAXPRO (see section 6.)
- if you have a severe illness with fever

Warnings and precautions

The container of this vaccine contains latex rubber. Latex rubber may cause severe allergic reactions.

Talk to your doctor, pharmacist or nurse before you receive HBVAXPRO 10 micrograms.

Others vaccines and HBVAXPRO 10 micrograms

HBVAXPRO can be administered at the same time as with hepatitis B immunoglobulin, at a separate injection site.

HBVAXPRO can be used to complete a primary immunisation course or as a booster dose in subjects who have previously received another hepatitis B vaccine.

HBVAXPRO can be administered at the same time as with other vaccines, using separate sites and syringes.

Tell your doctor, pharmacist or nurse if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Pregnancy and breast-feeding

Caution should be exercised when prescribing the vaccine to pregnant or breast-feeding women. Ask your doctor, pharmacist or nurse for advice before taking any medicine.

Driving and using machines

HBVAXPRO is expected to have no or negligible influence on the ability to drive and use machines.

HBVAXPRO 10 micrograms contains sodium: This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially "sodium- free".

3. How HBVAXPRO 10 micrograms is given

Dosage

The recommended dose for each injection (1 ml) is 10 micrograms for individuals 16 years of age or more.

A course of vaccination should include at least three injections.

Two immunisation schedules can be recommended:

- two injections with an interval of one month followed by a third injection 6 months after the first administration (0,1,6 months)
- if immunity is needed quickly: three injections with an interval of one month and a fourth dose 1 year later (0,1,2,12 months).

In case of a recent exposure to the hepatitis B virus, a first dose of HBVAXPRO together with the appropriate dose of immunoglobulin can be given.

Some local vaccination schedules currently include recommendations for a booster dose. Your doctor, pharmacist or nurse will inform you if a booster dose should be given.

For individuals less than 16 years of age, HBVAXPRO 10 micrograms is not recommended. The appropriate strength for administration to individuals from birth to 15 years of age is HBVAXPRO 5 micrograms.

Method of administration

The doctor or nurse will give the vaccine as an injection into muscle. The upper arm muscle is the preferred site for injection in adults and adolescents.

This vaccine should never be given into a blood vessel.

Exceptionally, the vaccine may be administered subcutaneously in patients with thrombocytopenia (diminution of blood platelets) or to persons at risk of haemorrhage.

If you forget one dose of HBVAXPRO 10 micrograms

If you miss a scheduled injection, talk to your doctor, pharmacist or nurse. Your doctor or nurse will decide when to give the missed dose.

If you have any further questions on the use of this product, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this vaccine can cause side effects, although not everybody gets them.

As with other hepatitis B vaccines, in many instances, the causal relationship of side effects to the vaccine has not been established.

The most common side effects seen are injection-site reactions: soreness, redness and hardening.

Other side effects are reported very rarely:

- Low platelet count, Lymph node disease
- Allergic reactions
- Nervous system disorders such as pins and needles, Facial paralysis, Nerve inflammations including Guillain-Barre Syndrome, Inflammation of the nerve of the eye that leads to impaired vision, Brain inflammation, Exacerbation of multiple sclerosis, Multiple sclerosis, Convulsions, Headache, Dizziness and Fainting
- Low blood pressure, Blood vessel inflammation
- Asthma-like symptoms
- Vomiting, Nausea, Diarrhoea, Abdominal pain
- Skin reactions such as eczema, Rash, Itching, Hives and Skin blistering, Hair loss
- Joint pain, Arthritis, Muscle pain, Pain in extremity
- Fatigue, Fever, Vague illness, Flu-like symptoms
- Elevation of liver enzymes
- Inflammation of the eye which causes pain and redness

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

United Kingdom

Yellow Card Scheme at: www.mhra.gov.uk/yellowcard

Ireland

HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; e-mail: medsafety@hpra.ie

5. How to store HBVAXPRO 10 micrograms

Keep this vaccine out of the sight and reach of children.

Do not use this vaccine after the expiry date which is stated on the label.

Store in a refrigerator between 2°C and 8°C.

Do not freeze.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What HBVAXPRO 10 micrograms contains

The active substance is:

Hepatitis B virus surface antigen, recombinant (HBsAg) *10 micrograms

Adsorbed on amorphous aluminium hydroxyphosphate sulfate (0.50 milligram Al⁺)

* produced in *Saccharomyces cerevisiae* (strain 2150-2-3) yeast by recombinant DNA technology.

The other ingredients are sodium chloride, borax and water for injections.

What HBVAXPRO 10 micrograms looks like and contents of the pack

HBVAXPRO 10 micrograms is a suspension for injection in a syringe.

Pack sizes of 1, 10 and 20 pre-filled syringes with 2 separate needles.

Pack sizes of 1 and 10 pre-filled syringes without needle, or with 1 separate needle.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder:

MSD VACCINS

162 avenue Jean Jaurès

69007 Lyon

France

Manufacturer Responsible for Batch Release:

Merck Sharp and Dohme, B.V.

Waarderweg, 39

2031 BN Haarlem

The Netherlands

For any information about this vaccine, please contact the local representative of the Marketing Authorisation Holder.

Ireland

Merck Sharp & Dohme Ireland (Human Health) Limited

Tel: +353 (0)1 2998700

medinfo_ireland@merck.com

United Kingdom

Merck Sharp & Dohme Limited
Tel: +44 (0) 1992 467272
medicalinformationuk@merck.com

España

Merck Sharp & Dohme de España, S.A.
Tel: +34 91 321 06 00
msd_info@merck.com

Italia

MSD Italia S.r.l.
Tel: +39 06 361911
medicalinformation.it@merck.com

This leaflet was last approved in January 2017.

Detailed information on this medicine is available on the European Medicines Agency website:
<http://www.ema.europa.eu>.

© Merck Sharp & Dohme Limited, 2017. All rights reserved.

PIL.HBV.10MCG.17.UK.4997-IA (MSD Vaccins)

The following information is intended for medical or health care professionals only:

Instructions

The vaccine should be inspected visually prior to administration for any foreign particulate matter and/or abnormal physical appearance. The syringe should be well shaken until a slightly opaque white suspension is obtained.

The needle is attached by twisting in clockwise direction, until the needle fits securely on the syringe.

PIL.HBV.10MCG.17.UK.4997-IA (MSD Vaccins)