

## PATIENT INFORMATION LEAFLET

# Scheriproct® 1.5 mg/g + 5 mg/g Rectal Ointment

Prednisolone, Cinchocaine hydrochloride

**Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If you experience any side effect and this becomes serious, tell your doctor or pharmacist.

### In this leaflet:

1. What Scheriproct is and what it is used for
2. Before you use Scheriproct
3. How to use Scheriproct
4. Possible side effects
5. How to store Scheriproct
6. Further information

## 1. What Scheriproct is and what it is used for

Scheriproct contains a substance which reduces inflammation (prednisolone), and a local anaesthetic (cinchocaine) which relieves pain.

This medicine is used for the relief of the inflammation, swelling, itching and soreness of piles (haemorrhoids) and to relieve itching of the anus (back passage). It is used short-term usually for 5 to 7 days.

## 2. Before you use Scheriproct

### Do not use Scheriproct if you:

- are allergic (hypersensitive) to prednisolone caproate, cinchocaine hydrochloride, other local anaesthetics or any of the other ingredients of Scheriproct.
- have a viral infection (e.g. herpes, shingles, chicken-pox)
- have any bacterial or fungal infections of the skin for which you are not receiving treatment

### Take special care with Scheriproct

- Long-term continuous treatment should be avoided because it can cause the skin to thin and deteriorate in the affected area (see section 4. "Possible Side Effects") and some of the medicine may be absorbed into the blood stream.
- If the area treated with Scheriproct is also infected your doctor should prescribe another medicine to use with Scheriproct, to treat the infection.
- Do not allow Scheriproct to come into contact with the eyes.

If symptoms do not improve, consult the doctor again.

### Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

### Pregnancy and breast feeding

Contact your doctor for advice before using this medicine if you are pregnant or intending to become pregnant or are breastfeeding.

There may be a very small risk to the development of a baby in pregnant women treated with Scheriproct. As with most medicines, this risk is likely to be greatest during the first 3 months of pregnancy.

### Driving and using machines:

Scheriproct has no influence on the ability to drive and use machines.

## 3. How to use Scheriproct

Always use Scheriproct exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. Do not use Scheriproct for more than 7 days.

Always wash your hands before and after applying Scheriproct.

### How to use Scheriproct Ointment:

1. Before you use the ointment, gently but thoroughly wash and dry the anus (back passage) and the skin around it.
2. If the ointment is to be used for the area **around** the anus:
  - squeeze a small quantity (about the size of a pea) on to the top of your finger. The amount of ointment in the nozzle is one dose.
  - spread it gently over the skin and just inside the anus.
  - do not rub it in.
3. If the ointment is to be used **inside** the anus:
  - put the separate nozzle (applicator) onto the tube.
  - squeeze the tube until the nozzle is full of ointment.
  - insert the nozzle very carefully into the anus until the whole length of the nozzle is inside.
  - then, while squeezing the tube gently, withdraw the nozzle.
  - wash the nozzle carefully in hot soapy water and rinse thoroughly.

Generally, the ointment should be applied thinly twice a day, but it may be applied three or four times on the first day, to obtain quick relief.

### If you use more Scheriproct than you should

If you accidentally swallow Scheriproct it is unlikely to be dangerous but contact your doctor or pharmacist if you are worried.

### If you forget to use Scheriproct

Do not use or take a double dose to make up for a forgotten dose. When you remember, use/take the next prescribed dose and continue with the treatment. See your doctor or pharmacist if you are worried.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

## 4. Possible side effects

Like all medicines, Scheriproct can cause side effects, though not everybody gets them.

Some thinning of the skin may occur if too much Scheriproct is applied for long periods of time (much longer than 5 - 7 days).

Allergic skin reactions may occur in rare cases.

If you experience any of these side effects and this becomes serious, or if you notice any side effects not listed in this leaflet please tell your doctor or pharmacist.

## 5. How to store Scheriproct

Store Scheriproct out of the reach and sight of children.

Do not store above 25°C.

Once opened, use within 3 months.

Replace the cap tightly after use.

The expiry date is printed on one flap of the carton and on the back of the tube after "EXP".

Do not use Scheriproct after the expiry date. The expiry date refers to the last day of that month.

## 6. Further information

### What Scheriproct Ointment contains

- The active substances are prednisolone caproate and cinchocaine hydrochloride. 1 g of Scheriproct Ointment contains prednisolone caproate equivalent to 1.5 mg of prednisolone and 5 mg cinchocaine hydrochloride.
- The other ingredients are:  
octyldodecanol, castor oil refined, castor oil hydrogenated, polyethylene glycol-400-monoricinoleate, chypre perfume oil.

### What Scheriproct looks like and contents of the pack

Scheriproct Ointment is a colourless to slightly yellowish ointment supplied in tubes of 30 g.

### Procured from within the EU by the Parallel Product

**Authorisation Holder:** Clear Pharmacy, 157-173 Roden Street, Belfast BT12 5QA, United Kingdom.

### Manufacturer:

Intendis Manufacturing SpA , Milan, Italy.

### Repackager:

Pharmaram Ltd Unit F4 Tom Bill Way, Ashby de la Zouch, Leicestershire LE65 2UY, UK.

*This medicinal product is authorized in the UK as Scheriproct.*

**Leaflet Date: July 2013**

20130814

*SCHERIPROCT is a registered trademark of Schering AG.*