

Gaviscon Liquid Aniseed Flavour Oral suspension.
Sodium Alginate 500mg/10ml, Sodium Bicarbonate 267mg/10ml, Calcium carbonate 160mg/10ml
Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to take this medicine carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve after 7 days.**
- If any of the side-effects gets serious, or if you notice any side-effect not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

- What is Gaviscon Liquid and what is it used for?
- Before taking this medicine.
- How to take this medicine.
- Possible side effects.
- How to store this medicine.
- Further information.

1. What is this medicine and what is it used for?
Gaviscon Liquid belongs to a group of medicines called "reflux suppressants".
This product forms a protective layer that floats on top of the stomach contents. This layer prevents reflux and keeps the stomach

RB46586

Pregnancy and breast-feeding:
You can take this product if you are pregnant or breast-feeding. As with all medicines, the treatment duration should be limited as much as possible.

Important information about some of the ingredients of Gaviscon Liquid:
This product contains methyl (E218) and propyl (E216) para-hydroxybenzoates which may cause allergic reactions (possibly delayed).

Driving and using machines:
This medicine has no influence on the ability to drive and use machines.

3. How to take this medicine
Check that the cap seal is unbroken before first using this product. Shake well before use.
Adults and children 12 years and over: 10-20ml (two to four 5 ml spoonfuls) after meals and at bedtime.
Children under 12 years: Should only be taken on medical advice. Not for use in children of 6 years or under.
If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.
Leave a time interval of 2 hours between taking Gaviscon and other medicinal products. Refer to Section 2 of this leaflet.
If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist if this doesn't go away.

4. Possible side effects
Very rarely (less than 1 in 10,000 patients) there is a chance of an allergic reaction to the ingredients. Symptoms of this may include skin rash, itching, difficulty breathing, dizziness or swelling of the face, lips, tongue or throat. If you experience these or any other side effects stop taking the product and consult your doctor immediately.

RB46586

contents away from the lining of the food pipe to relieve the symptoms of heartburn and acid indigestion.
This medicine is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion (related to reflux), for example, following meals, or during pregnancy, or in patients with symptoms related to reflux oesophagitis.

2. Before taking this medicine
Do not take Gaviscon Liquid:
- If you know you are allergic (hypersensitive) to any of the ingredients in this product (see 'Further Information' for a full list of ingredients).
Take special care with Gaviscon Liquid:
This medicinal product contains 6.2 mmol (or 141 mg) sodium per 10 ml.
To be taken into consideration by patients on a controlled sodium diet.
Consult your doctor if you are over 40 years and have never suffered with heartburn and acid indigestion before.
If symptoms persist after 7 days consult your doctor.
Taking other medicines:
Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without prescription.
A time interval of 2 hours should be considered between Gaviscon intake and the administration of other medicinal products, especially tetracyclines, fluoroquinolones, iron salts, thyroid hormones, chloroquine, bisphosphonates, and estramustine.

RB46586

Reporting of side effects
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via IMB Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517.
Website: www.imb.ie; e-mail imbpharmacovigilance@imb.ie
By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store this medicine
Do not use this product after the expiry date (EXP: month/year) shown. Do not store above 30°C. Keep out of the reach and sight of children. **Do not refrigerate.** Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Further Information
Each 10 ml of oral suspension contains 500 mg sodium alginate, 267mg sodium bicarbonate and 160mg calcium carbonate as the active ingredients. The other ingredients are carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, sodium hydroxide, aniseed flavour derived from fennel, erythrosine and purified water. This product does not contain sugar or gluten.
Gaviscon Liquid Oral Suspension is available in bottles of 150 ml, 300 ml or 600 ml.
Manufacturer: Reckitt Benckiser Healthcare (UK) Limited, Hull, HU8 7DS, UK.
PA Holder: Reckitt Benckiser Ireland Ltd., 7 Riverwalk, Citywest Business Campus, Dublin 24.
Date of preparation: September 2011.
Gaviscon and the sword and circle are trademarks.

RB46586

CUSTOMER INFO:
Minimum Point Size
Front Label = 7 pt

Minimum Point Size
Internal = 9 pt

TBRXXXX-D8055387

Gaviscon
Liquid
Aniseed

RB Artwork and Print Specification

Trident Reference No: RB146586
ZEN Ref: TR783820
Action: **B**
Brand: Gaviscon
Category: Adult
Segment Group: Standard Liquid
Segment: Aniseed
Pack Size: 150ml
Market/Country: EIRE
Date: 22/04/14

RBH Contact: Hazel Roche

Artwork Type: Submission
Component Code (2D if applicable): RB46586
Parent Technical Packaging Specification: D8055387
Finished Goods Code: Na
Supply Point: RB Hull
Pharmacode No/NE: N/A
Edgemark Position: Na

CAD Cam Ref: G-lbl-D8055387-60x155mm_3_page
Printer: Chesapeake (Bourne, UK)
Substrate: Self Adhesive Paper White

Technical & Non Printing Items

Cutter	Cutter 2 (if applicable)
Guides	Guides 2 (if applicable)

Colours (Label _Peel Read)

PANTONE Reflex Blue C	PANTONE 2375 C	PANTONE 320 C	PANTONE 185 C
1 FLX	2 FLX	3 FLX	4 FLX
Process Yellow	Braille	Vamh - Fluorescent WPR	Adhesive - Deadener
5 FLX	6 FLX	7 FLX	8 FLX

BARCODE INFO

Barcode Type: EAN 13
Barcode Number: 5000158065284
Magnification: 85%
Truncated By: 0.1 mm
Full Height: 22.2 mm
Bar Height (Smallest Bar): 19.4 mm
BWR: 30 microns
Encoded Data: N/A

Reckitt Benckiser

Reckitt Benckiser Healthcare (UK) Ltd,
Dansom Lane, Hull, HU8 7DS, U.K.

TRIDENT

Connaught House, Connaught Road,
Kingswood Business Park, Hull, HU7 3AP, England.
T: +44 (0) 1482 828100

Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.

STUDIO USE ONLY	Georgina Merry	v2.0
Sm•Art check results:	G=0; O=0; R=0; - GM - 22/04/14 08:07:18	

ARTWORK APPROVAL





Approved by Sonoco Trident on behalf of Reckitt Benckiser

Signed: _____

Date: _____

NOT APPROVED FOR COLOUR

UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM TRIDENT.



RB Artwork and Print Specification

Trident Reference No: RB146586
ZEN Ref: TR783820
Action: **B**
Brand: Gaviscon
Category: Adult
Segment Group: Standard Liquid
Segment: Aniseed
Pack Size: 150ml
Market/Country: EIRE
Date: 22/04/14

RBH Contact: Hazel Roche

Artwork Type: Submission

Component Code (2D if applicable):	RB46586
Parent Technical Packaging	
Specification:	D8055387
Finished Goods Code:	Na
Supply Point:	RB Hull
Pharmacode No/NE:	N/A
Edgemark Position:	Na

CAD Cam Ref: G-|b|-D8055387-60x155mm_3_pa
Printer: Chesapeake (Bourne, UK)
Substrate: Self Adhesive Paper White

Technical & Non Printing Items

 Cutter
 Guides
 Cutter 2 (if applicable)
 Guides 2 (if applicable)

Colours (Label _Peel Read)			
PANTONE Reflex Blue C		PANTONE 2375 C	
1	2	3	4
FLX	FLX	FLX	FLX
Process Yellow		Varnish - Fluorescent UVF	
5	6	7	8
FLX	FLX	FLX	FLX
Braille		Adhesive _Dead	

BARCODE INFO

Barcode Type:	EAN 13
Barcode Number:	5000158065284
Magnification:	85%
Truncated By:	0.1 mm
Full Height:	22.2 mm
Bar Height (Smallest Bar):	19.4 mm
BWR:	30 microns
Encoded Data:	N/A



Reckitt Benckiser Healthcare (UK) Ltd,
Dansom Lane, Hull, HU8 7DS, U.K.



Connaught House, Connaught Road,
Swood Business Park, Hull, HU7 3AP, England.
T: +44 (0) 1482 828100

Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.

STUDIO USE ONLY Georgina Merry

Sm•Art check results: G=0; O=0; R=0; - GM - 22/04/14 08:07:1

ARTWORK APPROVAL







Approved by Sonoco Trident on behalf of Reckitt Benckiser

Signed: _____

Date: _____

NOT APPROVED FOR COLOUR

UNDER NO CIRCUMSTANCES SHOULD THIS
ARTWORK BE ALTERED WITHOUT PRIOR
PERMISSION FROM TRIDENT.



RB Artwork
and Print Specification

Trident Reference No: RB146567

ZEN Ref: TR783737

Action: B

Brand: Gaviscon

Category: Adult

Segment Group: Standard Liquid

Segment: Aniseed

Pack Size: 300ml

Market/Country: EIRE

Date: 16/04/14

RBH Contact: Hazel Roche

Artwork Type: **Submission**

Component Code (2D if applicable): RB46567

Parent Technical Packaging Specification: D8081417

Finished Goods Code: Na

Supply Point: RB Hull

Pharmacode No/NE: N/A

Edgemark Position: Na

CAD Cam Ref: G-HL-D8081417 197x75mm_6_page

Printer: Chesapeake (Bourne, UK)

Substrate: PP-White SA

Technical & Non Printing Items

Cutter

Guides

Cutter 2 (if applicable)

Guides 2 (if applicable)

Colours (Label - Peel Read)

PANTONE Reflex Blue C

PANTONE 2375 C

PANTONE 520 C

PANTONE 185 C

Process Yellow

Brill

Vanish - Huescent 88F

Adhesive - Dye/Printer

BARCODE INFO

Barcode Type: EAN 13

Barcode Number: 5000158065925

Magnification: 100%

Truncated By: 0.0 mm

Full Height: 26.1 mm

Bar Height (Smallest Bar): 22.9 mm

BWR: 30 microns

Encoded Data: N/A

Reckitt
Benckiser

Reckitt Benckiser Healthcare (UK) Ltd,
Dansom Lane, Hull, HU8 7DS, U.K.

TRIDENT

Connaught House, Connaught Road,
Kingswood Business Park, Hull, HU7 3AP, England.
T: +44 (0) 1482 828100

Please note that any low resolution paper Canon colour
copies associated with this job should be referred to
for content, layout and colour separation only.

STUDIO USE ONLY

Sm•Art check results: G=0; O=0; R=0; - G/M - 16/04/14 09:23:44

ARTWORK APPROVAL

Approved by Sonoco Trident on behalf of Reckitt Benckiser

Signed: _____

Date: _____












NOT APPROVED FOR COLOUR

UNDER NO CIRCUMSTANCES SHOULD THIS
ARTWORK BE ALTERED WITHOUT PRIOR
PERMISSION FROM TRIDENT.

TBR1401-D8000089 - English

GAVISCON
LIQUID
ANISEED

CUSTOMER INFO:
Minimum Point Size = 6.50pt



What is this product? Each 10ml of oral suspension contains 500mg sodium alginate, 267mg sodium bicarbonate and 160mg calcium carbonate.

Dosage: Check that the cap seal is unbroken before first using this product. Shake well before use. Read the package leaflet before use. For oral use. Adults and children 12 years and over: Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime. Children 6 to 12 years: 5-10 ml (one to two 5ml spoonfuls) after meals and at bedtime. Not for use in children of 6 years and under. **If symptoms persist after 7 days consult your doctor. Consult your doctor if you are over 40 years and have never suffered with heartburn and indigestion before.** Contains sodium, methyl (E218) and propyl (E216) para-hydroxybenzoates. See leaflet for further information. Do not store above 30°C. **Do not refrigerate.** **Keep out of the reach and sight of children. Do not use this product after the expiry date (EXP: month/year shown).** Peel here. Do not remove.

 **e 600ml**



GAVISCON
LIQUID ANISEED FLAVOUR
ORAL SUSPENSION

Each 10ml contains:
sodium alginate 500mg
sodium bicarbonate 267 mg
calcium carbonate 160 mg

FAST SOOTHING RELIEF
Heartburn and indigestion • Long lasting •

ANISEED FLAVOUR

Manufacturer: Reckitt Benckiser Healthcare (UK) Ltd., Hull HU8 7DS, UK.
PA Holder: Reckitt Benckiser Ireland Ltd., Citywest Business Campus, Dublin 24.
PA 979/15/3
For product queries please call (01) 6305429



BN EXP

Gaviscon Liquid Aniseed Flavour Oral Suspension
Sodium alginate 500mg/10ml, Sodium bicarbonate 267mg/10ml, Calcium carbonate 160mg/10ml
Read all of this leaflet carefully because it contains important information for you.
This medicine is available without prescription. However, you still need to take this medicine carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve after 7 days.**
- If any of the side-effects gets serious, or if you notice any side-effect not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

- What is Gaviscon Liquid and what is it used for?
- Before taking this medicine.
- How to take this medicine.
- Possible side effects.
- How to store this medicine.
- Further information.

1. What is this medicine and what is it used for?
Gaviscon Liquid belongs to a group of medicines called 'reflux suppressants'.
This product forms a protective layer that floats on top of the stomach contents. This layer prevents reflux and keeps the stomach contents away from the lining of the food pipe to relieve the symptoms of heartburn and acid indigestion.
This medicine is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion (related to reflux), for example, following meals, or during pregnancy, or in patients with symptoms related to reflux oesophagitis.

2. Before taking this medicine
Do not take Gaviscon Liquid:
- If you know you are allergic (hypersensitive) to any of the ingredients in this product (see 'Further Information' for a full list of ingredients).
Take special care with Gaviscon Liquid:
This medicinal product contains 6.2 mmol (or 141 mg) sodium per 10 ml. To be taken into consideration by patients on a controlled sodium diet.

If symptoms persist after 7 days consult your doctor. Consult your doctor if you are over 40 years and have never suffered with heartburn and acid indigestion before.
Taking other medicines:
Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without prescription
A time interval of 2 hours should be considered between Gaviscon intake and the administration of other medicinal products, especially tetracyclines, fluoroquinolones, iron salts, thyroid hormones, chloroquine, bisphosphonates, and estramustine.
Pregnancy and breast-feeding:
You can take this product if you are pregnant or breast-feeding. As with all medicines, the treatment duration should be limited as much as possible.
Important information about some of the ingredients of Gaviscon Liquid:
This product contains methyl (E218) and propyl (E216) para-hydroxybenzoates which may cause allergic reactions (possibly delayed).

BB46148

Driving and using machines:
This medicine has no influence on the ability to drive and use machines.

3. How to take this medicine
Check that the cap seal is unbroken before first using this product. Shake well before use.

Adults and children 12 years and over: 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime.
Children 6-12 years: Take 5-10ml (one to two 5ml spoonfuls) after meals and at bedtime.
Children under 6 years: not recommended.
If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.
Leave a time interval of 2 hours between taking Gaviscon and other medicinal products. Refer to Section 2 of this leaflet.
If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist if this doesn't go away.

4. Possible side effects
Very rarely (less than 1 in 10,000 patients) there is a chance of an allergic reaction to the ingredients.
Symptoms of this may include skin rash, itching, difficulty


breathing, dizziness or swelling of the face, lips, tongue or throat. If you experience these or any other side effects stop taking the product and consult your doctor immediately.

Reporting of side effects
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via IMB Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.imb.ie; e-mail imbpharmacovigilance@imb.ie
By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store this medicine
Do not use this product after the expiry date (EXP: month/year) shown. Do not store above 30°C. Keep out of the reach and sight of children. **Do not refrigerate.** Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.


6. Further Information
Each 10 ml of oral suspension contains 500 mg sodium alginate and 267mg sodium bicarbonate and 160mg calcium carbonate as the active ingredients. The other ingredients are, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, sodium hydroxide, aniseed flavour derived from fennel, erythrosine and purified water. This product does not contain sugar or gluten.
Gaviscon Liquid Oral Suspension is available in bottles of 150 ml, 300 ml or 600ml.
Manufacturer: Reckitt Benckiser Healthcare (UK) Limited, Hull, HU8 7DS, UK.
PA Holder: Reckitt Benckiser Ireland Ltd., 7 Riverwalk, Citywest Business Campus, Dublin 24.
Date of preparation: September 2011.
Gaviscon and the sword and circle are trademarks.

BB46148



CUSTOMER INFO:
Minimum Point Size = 7.00pt
(Refers to the FOP label, the copy on the inside leaflet is a minimum of 9pt).

TBR1279-D0014922



Gaviscon
Liquid
Aniseed

RB Artwork and Print Specification





Trident Reference No: RB146148
ZEN Ref: TR781450
Action: A
Brand: Gaviscon
Category: Adult
Segment Group: Standard Liquid
Segment: Aniseed
Pack Size: 600ml
Market/Country: EIRE
Date: 02/04/14

RBH Contact: Hazel Roche








Artwork Type: Submission
Component Code (2D if applicable): RB46148
Parent Technical Packaging
Specification: D0014922
Finished Goods Code: Na
Supply Point: RB Hull
Pharmacode No/NE: N/A
Edgemark Position: Na

CAD Cam Ref: G-lbl-D0014922-197x75mm
Printer: Chesapeake (Bourne, UK)
Substrate: Self Adhesive Paper White

Technical & Non Printing Items


 Cutter	 Cutter 2 (if applicable)
 Guides	 Guides 2 (if applicable)


Colours (Label _Peel Read)

BARCODE INFO

Barcode Type: EAN 13
Barcode Number: 5000158065932
Magnification: 100%
Truncated By: 0.0 mm
Full Height: 26.2 mm
Bar Height (Smallest Bar): 22.9 mm
BWR: 30 microns
Encoded Data: N/A


Reckitt Benckiser Healthcare (UK) Ltd,
Dansom Lane, Hull, HU8 7DS, U.K.


TRIDENT
Connaught House, Connaught Road,
Kingswood Business Park, Hull, HU7 3AP, England.
T: +44 (0) 1482 828100

Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.

STUDIO USE ONLY Rebecca Milner v2.0
Sm•Art check results: G=0; O=0; R=0; - RM - 02/04/14 16:16:48

ARTWORK APPROVAL

Approved by Sonoco Trident on behalf of Reckitt Benckiser

Signed: _____

Date: _____

NOT APPROVED FOR COLOUR

UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM TRIDENT.

What is this product? Each 10ml of oral suspension contains 500mg sodium alginate, 267mg sodium bicarbonate and 160mg calcium carbonate.

Dosage: Check that the cap seal is unbroken before first using this product. Shake well before use. Read the package leaflet before use. For oral use. Adults and children 12 years and over: Take 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime. Children 6 to 12 years: 5-10 ml (one to two 5ml spoonfuls) after meals and at bedtime. Not for use in children of 6 years and under. If symptoms persist after 7 days consult your doctor. Consult your doctor if you are over 40 years and have never suffered with heartburn and indigestion before. Contains sodium, methyl (E218) and propyl (E216) para-hydroxybenzoates. See leaflet for further information. Do not store above 30°C. Do not refrigerate. Keep out of the reach and sight of children. Do not use this product after the expiry date (EXP: month/year shown). Peel here. Do not remove.

GAVISCON
LIQUID ANISEED FLAVOUR
ORAL SUSPENSION

Each 10ml contains:
sodium alginate 500mg
sodium bicarbonate 267 mg
calcium carbonate 160 mg

FAST SOOTHING RELIEF
Heartburn and indigestion • Long lasting •

ANISEED FLAVOUR

Manufacturer: Reckitt Benckiser Healthcare (UK) Ltd., Hull HU8 7DS, UK.
PA Holder: Reckitt Benckiser Ireland Ltd., Citywest Business Campus, Dublin 24.
PA 979/15/3
For product queries please call (01) 6305429

Gaviscon Liquid Aniseed Flavour Oral Suspension
Sodium alginate 500mg/10ml, Sodium bicarbonate 267mg/10ml, Calcium carbonate 160mg/10ml
Read all of this leaflet carefully because it contains important information for you.
This medicine is available without prescription. However, you still need to take this medicine carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve after 7 days.**
- If any of the side-effects gets serious, or if you notice any side-effect not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

- What is Gaviscon Liquid and what is it used for?
- Before taking this medicine.
- How to take this medicine.
- Possible side effects.
- How to store this medicine.
- Further information.

1. What is this medicine and what is it used for?
Gaviscon Liquid belongs to a group of medicines called 'reflux suppressants'.
This product forms a protective layer that floats on top of the stomach contents. This layer prevents reflux and keeps the stomach contents away from the lining of the food pipe to relieve the symptoms of heartburn and acid indigestion.
This medicine is used for the treatment of symptoms of gastro-oesophageal reflux such as acid regurgitation, heartburn and indigestion (related to reflux), for example, following meals, or during pregnancy, or in patients with symptoms related to reflux oesophagitis.

2. Before taking this medicine
Do not take Gaviscon Liquid:
- If you know you are allergic (hypersensitive) to any of the ingredients in this product (see 'Further Information' for a full list of ingredients).
Take special care with Gaviscon Liquid:
This medicinal product contains 6.2 mmol (or 141 mg) sodium per 10 ml. To be taken into consideration by patients on a controlled sodium diet.

If symptoms persist after 7 days consult your doctor. Consult your doctor if you are over 40 years and have never suffered with heartburn and acid indigestion before.
Taking other medicines:
Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without prescription.
A time interval of 2 hours should be considered between Gaviscon intake and the administration of other medicinal products, especially tetracyclines, fluoroquinolones, iron salts, thyroid hormones, chloroquine, bisphosphonates, and estramustine.
Pregnancy and breast-feeding:
You can take this product if you are pregnant or breast-feeding. As with all medicines, the treatment duration should be limited as much as possible.
Important information about some of the ingredients of Gaviscon Liquid:
This product contains methyl (E218) and propyl (E216) para-hydroxybenzoates which may cause allergic reactions (possibly delayed).

Driving and using machines:
This medicine has no influence on the ability to drive and use machines.

3. How to take this medicine
Check that the cap seal is unbroken before first using this product. Shake well before use.

Adults and children 12 years and over: 10-20ml (two to four 5ml spoonfuls) after meals and at bedtime.
Children 6-12 years: Take 5-10ml (one to two 5ml spoonfuls) after meals and at bedtime.
Children under 6 years: not recommended.
If you forget a dose it is not necessary to double the dose next time, just carry on taking as before.
Leave a time interval of 2 hours between taking Gaviscon and other medicinal products. Refer to Section 2 of this leaflet.
If you take too much of this product you may feel bloated. It is unlikely to cause you any harm, but please consult your doctor or pharmacist if this doesn't go away.


4. Possible side effects
Very rarely (less than 1 in 10,000 patients) there is a chance of an allergic reaction to the ingredients.
Symptoms of this may include skin rash, itching, difficulty

breathing, dizziness or swelling of the face, lips, tongue or throat. If you experience these or any other side effects stop taking the product and consult your doctor immediately.

Reporting of side effects
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via IMB Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.imb.ie; e-mail imbpharmacovigilance@imb.ie
By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store this medicine
Do not use this product after the expiry date (EXP: month/year) shown. Do not store above 30°C. Keep out of the reach and sight of children. **Do not refrigerate.** Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Further Information
Each 10 ml of oral suspension contains 500 mg sodium alginate and 267mg sodium bicarbonate and 160mg calcium carbonate as the active ingredients. The other ingredients are, carbomer, methyl (E218) and propyl (E216) parahydroxybenzoates, sodium saccharin, sodium hydroxide, aniseed flavour derived from fennel, erythrosine and purified water. This product does not contain sugar or gluten.
Gaviscon Liquid Oral Suspension is available in bottles of 150 ml, 300 ml or 600ml.
Manufacturer: Reckitt Benckiser Healthcare (UK) Limited, Hull, HU8 7DS, UK.
PA Holder: Reckitt Benckiser Ireland Ltd., 7 Riverwalk, Citywest Business Campus, Dublin 24.
Date of preparation: September 2011.
Gaviscon and the sword and circle are trademarks.



CUSTOMER INFO:
Minimum Point Size = 7.00pt
(Refers to the FOP label, the copy on the inside leaflet is a minimum of 9pt).

TBR1279-D0014922

Gaviscon Liquid Aniseed

RB Artwork and Print Specification

Trident Reference No: RB146148
ZEN Ref: TR781450
Action: A
Brand: Gaviscon
Category: Adult
Segment Group: Standard Liquid
Segment: Aniseed
Pack Size: 600ml
Market/Country: EIRE
Date: 02/04/14

RBH Contact: Hazel Roche

Artwork Type: Submission
Component Code (2D if applicable): RB46148
Parent Technical Packaging
Specification: D0014922
Finished Goods Code: Na
Supply Point: RB Hull
Pharmacode No/NE: N/A
Edgemark Position: Na

CAD Cam Ref: G-lbl-D0014922-197x75mm
Printer: Chesapeake (Bourne, UK)
Substrate: Self Adhesive Paper White

Technical & Non Printing Items

Cutter	Cutter 2 (if applicable)
Guides	Guides 2 (if applicable)

Colours (Label _Peel Read)

PANTONE Reflex Blue C	PANTONE 320 C	PANTONE 2375 C	PANTONE 185 C
1 FLX	2 FLX	3 FLX	4 FLX
Process Yellow	Black	Variable - fluorescent UVF	
5 FLX	6 FLX	7 FLX	

BARCODE INFO

Barcode Type: EAN 13
Barcode Number: 5000158065932
Magnification: 100%
Truncated By: 0.0 mm
Full Height: 26.2 mm
Bar Height (Smallest Bar): 22.9 mm
BWR: 30 microns
Encoded Data: N/A

Reckitt Benckiser Healthcare (UK) Ltd,
Dansom Lane, Hull, HU8 7DS, U.K.

TRIDENT
Connaught House, Connaught Road,
Kingswood Business Park, Hull, HU7 3AP, England.
T: +44 (0) 1482 828100

Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.

STUDIO USE ONLY Rebecca Milner v2.0
Sm•Art check results: G=0; O=0; R=0; - RM - 02/04/14 16:16:48

ARTWORK APPROVAL

Approved by Sonoco Trident on behalf of Reckitt Benckiser

Signed: _____

Date: _____

NOT APPROVED FOR COLOUR

UNDER NO CIRCUMSTANCES SHOULD THIS ARTWORK BE ALTERED WITHOUT PRIOR PERMISSION FROM TRIDENT.