

Ivabradine Rowex 5 mg Film-coated tablets
Ivabradine Rowex 7.5 mg Film-coated tablets

ivabradine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

- 1. What Ivabradine Rowex is and what it is used for
- 2. What you need to know before you take Ivabradine Rowex
- 3. How to take Ivabradine Rowex
- 4. Possible side effects
- 5. How to store Ivabradine Rowex
- 6. Contents of the pack and other information



1. What Ivabradine Rowex is and what it is used for

Ivabradine Rowex (ivabradine) is a heart medicine used to treat:

- Symptomatic stable angina pectoris (which causes chest pain) in adult patients whose heart rate is over or equal to 70 beats per minute. It is used in adult patients who do not tolerate or cannot take heart medicines called beta-blockers. It is also used in combination with beta-blockers in adult patients whose condition is not fully controlled with a beta-blocker.
- Chronic heart failure in adult patients whose heart rate is over or equal to 75 beats per minute. It is used in combination with standard therapy, including beta-blocker therapy or when beta-blockers are contraindicated or not tolerated.

About stable angina pectoris (usually referred to as “angina”):
Stable angina is a heart disease which happens when the heart does not receive enough oxygen. It usually appears between 40 and 50 years of age. The most common symptom of angina is chest pain or discomfort. Angina is more likely to happen when the heart beats faster in situations such as exercise, emotion, exposure to the cold or after eating. This increase in heart rate can cause the chest pain in people who suffer from angina.

About chronic heart failure:
Chronic heart failure is a heart disease which happens when your heart cannot pump enough blood to the rest of your body. The most common symptoms of heart failure are breathlessness, fatigue, tiredness and ankle swelling.

How does Ivabradine Rowex work?
Ivabradine Rowex mainly works by reducing the heart rate by a few beats per minute. This lowers the heart’s need for oxygen especially in the situations when an angina attack is more likely to happen. In this way Ivabradine Rowex helps to control and reduce the number of angina attacks.
Furthermore as elevated heart rate adversely affects the heart functioning and vital prognosis in patients with chronic heart failure, the specific heart rate lowering action of ivabradine helps to improve the heart functioning and vital prognosis in these patients.

2. What you need to know before you take Ivabradine Rowex

Do not take Ivabradine Rowex

- if you are allergic to ivabradine or any of the other ingredients of this medicine (listed in section 6)
- if your resting heart rate before treatment is too slow (below 70 beats per minute)
- if you are suffering from cardiogenic shock (a heart condition treated in hospital)
- if you suffer from a heart rhythm disorder
- if you are having a heart attack
- if you suffer from very low blood pressure
- if you suffer from unstable angina (a severe form in which chest pain occurs very frequently and with or without exertion)
- if you have heart failure which has recently become worse
- if your heartbeat is exclusively imposed by your pacemaker
- if you suffer from severe liver problems
- if you are already taking medicines for the treatment of fungal infections (such as ketoconazole, itraconazole), macrolide antibiotics (such as josamycin, clarithromycin, telithromycin or erythromycin given orally), medicines to treat HIV infections (such as nelfinavir, ritonavir) or nefazodone (medicine to treat depression) or diltiazem, verapamil (used for high blood pressure or angina pectoris)
- if you are a woman able to have children and not using reliable contraception
- if you are pregnant or trying to become pregnant
- if you are breast-feeding.

Warnings and precautions

Talk to your doctor or pharmacist before taking Ivabradine Rowex

- if you suffer from heart rhythm disorders (such as irregular heartbeat, palpitation, increase in chest pain) or sustained atrial fibrillation (a type of irregular heartbeat), or an abnormality of electrocardiogram (ECG) called ‘long QT syndrome’
- if you have symptoms such as tiredness, dizziness or shortness of breath (this could mean that your heart is slowing down too much)
- if you suffer from symptoms of atrial fibrillation (pulse rate at rest unusually high (over 110 beats per minute) or irregular, without any apparent reason, making it difficult to measure)
- if you have had a recent stroke (cerebral attack)
- if you suffer from mild to moderate low blood pressure
- if you suffer from uncontrolled blood pressure, especially after a change in your antihypertensive treatment

- if you suffer from severe heart failure or heart failure with abnormality of ECG called ‘bundle branch block’
- if you suffer from chronic eye retinal disease
- if you suffer from moderate liver problems
- if you suffer from severe renal problems.

If any of the above applies to you, talk straight away to your doctor before or while taking Ivabradine Rowex.

Children and adolescents

Ivabradine Rowex is not intended for use in children and adolescents younger than 18 years.

Other medicines and Ivabradine Rowex

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Make sure to tell your doctor if you are taking any of the following medicines, as a dose adjustment of Ivabradine Rowex or monitoring should be required:

- fluconazole (an antifungal medicine)
- rifampicin (an antibiotic)
- barbiturates (for difficulty sleeping or epilepsy)
- phenytoin (for epilepsy)
- *Hypericum perforatum* or St John’s Wort (herbal treatment for depression)
- QT prolonging medicines to treat either heart rhythm disorders or other conditions:
 - quinidine, disopyramide, ibutilide, sotalol, amiodarone (to treat heart rhythm disorders)
 - bepridil (to treat angina pectoris)
 - certain types of medicines to treat anxiety, schizophrenia or other psychoses (such as pimozone, ziprasidone, sertindole)
 - anti-malarial medicines (such as mefloquine or halofantrine)
 - intravenous erythromycin (an antibiotic)
 - pentamidine (an antiparasitic medicine)
 - cisapride (against the gastro-oesophageal reflux).

Some types of diuretics which may cause decrease in blood potassium level, such as furosemide, hydrochlorothiazide, indapamide (used to treat oedema, high blood pressure).

Ivabradine Rowex with food and drink

Avoid grapefruit juice during treatment with Ivabradine Rowex.

Pregnancy and breast-feeding

Do not take Ivabradine Rowex if you are pregnant or are planning to have a baby (see “Do not take Ivabradine Rowex”).
If you are pregnant and have taken Ivabradine Rowex, talk to your doctor.
Do not take Ivabradine Rowex if you are able to become pregnant unless you use reliable contraceptive measures (see “Do not take Ivabradine Rowex”).
Do not take Ivabradine Rowex if you are breast-feeding (see “Do not take Ivabradine Rowex”). Talk to your doctor if you are breast-feeding or intending to breast-feed as breast-feeding should be discontinued if you take Ivabradine Rowex.
If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

Ivabradine Rowex may cause temporary luminous visual phenomena (a temporary brightness in the field of vision, see “Possible side effects”). If this happens to you, be careful when driving or using machines at times when there could be sudden changes in light intensity, especially when driving at night.

Ivabradine Rowex contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Ivabradine Rowex

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.
Ivabradine Rowex should be taken during meals.
The tablet of 5 mg strength can be divided into equal doses.

If you are being treated for stable angina pectoris
The starting dose should not exceed one tablet of Ivabradine Rowex 5 mg twice daily. If you still have angina symptoms and if you have tolerated the 5 mg twice daily dose well, the dose may be increased. The maintenance dose should not exceed 7.5 mg twice daily. Your doctor will prescribe the right dose for you. The recommended dose is one tablet in the morning and one tablet in the evening. In some cases (e.g. if you are elderly), your doctor may prescribe half the dose i.e., one half 5 mg tablet of Ivabradine Rowex 5 mg (corresponding to 2.5 mg ivabradine) in the morning and one half 5 mg tablet in the evening.

If you are being treated for chronic heart failure
The recommended starting dose is one tablet of Ivabradine Rowex 5 mg twice daily increasing if necessary to one tablet of Ivabradine Rowex 7.5 mg twice daily. Your doctor will decide the right dose for you. The usual dose is one tablet in the morning and one tablet in the evening. In some cases (e.g. if you are elderly), your doctor may prescribe half the dose i.e., one half 5 mg tablet of Ivabradine Rowex 5 mg (corresponding to 2.5 mg ivabradine) in the morning and one half 5 mg tablet in the evening.

If you take more Ivabradine Rowex than you should:
A large dose of Ivabradine Rowex could make you feel breathless or tired because your heart slows down too much. If this happens, contact your doctor immediately.

If you forget to take Ivabradine Rowex:
If you forget to take a dose of Ivabradine Rowex, take the next dose at the usual time. Do not take a double dose to make up for a forgotten dose.

[For calendar blister packs]The calendar printed on the blister containing the tablets should help you remember when you last took a tablet of Ivabradine Rowex.

If you stop taking Ivabradine Rowex:
As the treatment for angina or chronic heart failure is usually life-long, you should discuss with your doctor before stopping this medicinal product.
If you think that the effect of Ivabradine Rowex is too strong or too weak, talk to your doctor or pharmacist.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The most common adverse reactions with this medicine are dose dependent and related to its mode of action:

Very common (may affect more than 1 in 10 people):
Luminous visual phenomena (brief moments of increased brightness, most often caused by sudden changes in light intensity). They can also be described as a halo, coloured flashes, image decomposition or multiple images. They generally occur within the first two months of treatment after which they may occur repeatedly and resolve during or after treatment.

Common (may affect up to 1 in 10 people):
Modification in the heart functioning (the symptoms are a slowing down of the heart rate). They particularly occur within the first 2 to 3 months of treatment initiation.

Other side effects have also been reported:

Common (may affect up to 1 in 10 people):
Irregular rapid contraction of the heart, abnormal perception of heartbeat, uncontrolled blood pressure, headache, dizziness and blurred vision (cloudy vision).

Uncommon (may affect up to 1 in 100 people):
Palpitations and cardiac extra beats, feeling sick (nausea), constipation, diarrhoea, abdominal pain, spinning sensation (vertigo), difficulty breathing (dyspnoea), muscle cramps, changes in laboratory parameters: high blood levels of uric acid, an excess of eosinophils (a type of white blood cell) and elevated creatinine in blood (a breakdown product of muscle), skin rash, angioedema (such as swollen face, tongue or throat, difficulty in breathing or swallowing), low blood pressure, fainting, feeling of tiredness, feeling of weakness, abnormal ECG heart tracing, double vision, impaired vision.

Rare (may affect up to 1 in 1,000 people):
Urticaria, itching, skin reddening, feeling unwell.

Very rare (may affect up to 1 in 10, 000 people):
Irregular heartbeats.

Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL – Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Ivabradine Rowex

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, the tablet container and blister after ‘EXP’. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not use the medicine packed in bottles longer than 6 months after first opening.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Ivabradine Rowex contains
– The active substance is ivabradine (as oxalate).
Ivabradine Rowex 5 mg: one film-coated tablet contains 5 mg ivabradine (equivalent to 5.961 mg ivabradine as oxalate).
Ivabradine Rowex 7.5 mg: one film-coated tablet contains 7.5 mg ivabradine (equivalent to 8.941 mg ivabradine as oxalate).
– The other ingredients in the tablet core are: lactose anhydrous; silica, colloidal anhydrous; croscarmellose sodium (E468); butylhydroxytoluene (E321),

magnesium stearate (E470b) and in the tablet coating: Hypromellose (E464), titanium dioxide (E171), macrogol 6000, glycerol (E422), magnesium stearate (E470b), yellow iron oxide (E172), red iron oxide (E172).

What Ivabradine Rowex looks like and contents of the pack

Ivabradine Rowex 5 mg tablets are yellow coloured, round, film-coated tablets debossed with 5 on one side and scored on other side. The tablet can be divided into equal doses.
Ivabradine Rowex 7.5 mg tablets are orange-yellow coloured, round, biconvex, film-coated tablets debossed with 7.5 on one side.

The tablets are available in blister packs (Aluminium-Aluminium; PVC/PE/PVdC-Aluminium) of 10, 14, 28, 30, 56, 60, 84, 90, 98, 100, 112 film-coated tablets. The tablets are available in calendar blister packs (Aluminium-Aluminium; PVC/PE/PVdC-Aluminium) of 14, 28, 30, 56, 60, 84, 100, 112 film-coated tablets. The tablets are available in tablet containers (HDPE) of 100, 250 film-coated tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturers
Marketing Authorisation Holder
Rowex Ltd., Bantry, Co. Cork, Ireland.

Manufacturers
Combino Pharm Ltd., HF60 Hal Far Industrial Estate, BBG3000 Malta.
HBM Pharma s.r.o., Sklabinská 30, Martin, 036 80 Slovak Republic.
Salutas Pharma GmbH, Otto-von-Guericke-Allee 1, Sachsen-Anhalt, Barleben 39179, Germany.

This medicinal product is authorised in the Member States of the EEA under the following names:

The Netherlands	Ivabradine Sandoz 5 mg, filmomhulde tabletten Ivabradine Sandoz 7,5 mg, filmomhulde tabletten
Austria	Ivabradin Sandoz 5 mg – Filmtabletten Ivabradin Sandoz 7,5 mg – Filmtabletten
Belgium	Ivabradine Sandoz 5 mg filmomhulde tabletten Ivabradine Sandoz 7,5 mg filmomhulde tabletten
Cyprus	Ivabradine Sandoz 5 mg Ivabradine Sandoz 7.5 mg
Czech Republic	Ivabradin Sandoz 5 mg Ivabradin Sandoz 7,5 mg
Germany	Ivabradin - 1 A Pharma 5 mg Filmtabletten Ivabradin - 1 A Pharma 7,5 mg Filmtabletten
Greece	Ivabradine/Sandoz
Spain	Ivabradina Sandoz Farmacéutica 5 mg comprimidos recubiertos con película EFG Ivabradina Sandoz Farmacéutica 7.5 mg comprimidos recubiertos con película EFG
France	IVABRADINE SANDOZ 5 mg, comprimé pelliculé sécable IVABRADINE SANDOZ 7,5 mg, comprimé pelliculé
Croatia	Ivabradin Sandoz 5 mg filmom obložene tablete Ivabradin Sandoz 7,5 mg filmom obložene tablete
Hungary	Ivabradin Sandoz 5 mg filmtabletta Ivabradin Sandoz 7,5 mg filmtabletta
Ireland	Ivabradine Rowex 5 mg Film-coated tablets Ivabradine Rowex 7.5 mg Film-coated tablets
Luxemburg	Ivabradin Sandoz 5 mg comprimés pelliculés Ivabradin Sandoz 7,5 mg comprimés pelliculés
Portugal	Ivabradina Sandoz
Romania	Ivabradină Sandoz 5 mg comprimate filmate Ivabradină Sandoz 7,5 mg comprimate filmate
Slovakia	Ivabradín Sandoz 5 mg Ivabradín Sandoz 7,5 mg

This leaflet was last revised in 10/2016.

I.M. L/963a+b 11-16
article no.

Internal codes, article number, pharma codes and measurements will be printed during the packaging process.

Artwork Checklist		Job Description	Version Number	Size	Inks	ACCF 19098										
		Ivabradine Rowex 5 mg & 7.5 mg Film-coated tablets	0.5	165 x 500												
					Colour	Black										
		Market	Proof Date	Cutter Ref.			Front									
		IM	21/12/2016	N/A			Back									
							Prints On									