

Ibandronic Acid Mylan 150 mg Film-coated Tablets

ibandronic acid

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

- | | |
|--|---|
| 1. What Ibandronic Acid Mylan is and what it is used for | 3. How to take Ibandronic Acid Mylan |
| 2. What you need to know before you take Ibandronic Acid Mylan | 4. Possible side effects |
| | 5. How to store Ibandronic Acid Mylan |
| | 6. Contents of the pack and other information |

1. What Ibandronic Acid Mylan is and what it is used for

Ibandronic Acid Mylan belongs to a group of medicines called **bisphosphonates**. It contains the active substance ibandronic acid. Ibandronic Acid Mylan may reverse bone loss by stopping more loss of bone and increasing bone mass in most women who take it, even though they won't be able to see or feel a difference. Ibandronic Acid Mylan may help lower the chances of breaking bones (fractures). This reduction in fractures was shown for the spine but not for the hip.

Ibandronic Acid Mylan is prescribed to you to treat postmenopausal osteoporosis because you have an increased risk of fractures. Osteoporosis is a thinning and weakening of the bones, which is common in women after the menopause. At the menopause, a woman's ovaries stop producing the female hormone, oestrogen, which helps to keep her skeleton healthy.

The earlier a woman reaches the menopause, the greater her risk of fractures in osteoporosis. Other things that can increase the risk of fractures include:

- not enough calcium and vitamin D in the diet
- smoking, or drinking too much alcohol
- not enough walking or other weight-bearing exercise
- a family history of osteoporosis

A **healthy lifestyle** will also help you to get the most benefit from your treatment. This includes:

- eating a balanced diet rich in calcium and vitamin D
- walking or any other weight-bearing exercise
- not smoking; and not drinking too much alcohol.

2. What you need to know before you take Ibandronic Acid Mylan

Do not take Ibandronic Acid Mylan:

- If you are allergic to ibandronic acid or any of the other ingredients of this medicine listed in section 6.
- If you have certain problems with your gullet/food pipe (oesophagus) such as narrowing or difficulty swallowing.
- If you can't stand or sit upright for at least one hour (60 minutes) at a time.
- **If you have, or had in the past low blood calcium.** Please consult your doctor.

Warnings and precautions

A side effect called osteonecrosis of the jaw (ONJ) (bone damage in the jaw) has been reported very rarely in the post marketing setting in patients receiving ibandronic acid for osteoporosis. ONJ can also occur after stopping treatment.

It is important to try and prevent ONJ developing as it is a painful condition that can be difficult to treat. In order to reduce the risk of developing osteonecrosis of the jaw, there are some precautions you should take.

Before receiving treatment, tell your doctor/nurse (health care professional) if:

- you have any problems with your mouth or teeth such as poor dental health, gum disease, or a planned tooth extraction
- you don't receive routine dental care or have not had a dental check up for a long time
- you are a smoker (as this may increase the risk of dental problems)
- you have previously been treated with a bisphosphonate (used to treat or prevent bone disorders)
- you are taking medicines called corticosteroids (such as prednisolone or dexamethasone)
- you have cancer

Your doctor may ask you to undergo a dental examination before starting treatment with Ibandronic Acid Mylan.

While being treated, you should maintain good oral hygiene (including regular teeth brushing) and receive routine dental check-ups. If you wear dentures you should make sure these fit properly. If you are under dental treatment or will undergo dental surgery (e.g. tooth extractions), inform your doctor about your dental treatment and tell your dentist that you are being treated with ibandronic acid.

Contact your doctor and dentist immediately if you experience any problems with your mouth or teeth such as loose teeth, pain or swelling, or non-healing of sores or discharge, as these could be signs of osteonecrosis of the jaw.

Some people need to be especially careful while they are taking ibandronic acid. Talk to your doctor or pharmacist before taking Ibandronic Acid Mylan:

- If you have any disturbances of mineral metabolism (such as vitamin D deficiency).
- If your kidneys are not functioning normally.
- If you have any swallowing or digestive problems.

Irritation, inflammation or ulceration of the gullet/food pipe (oesophagus) often with symptoms of severe pain in the chest, severe pain after swallowing food and/or drink, severe nausea, or vomiting may occur, especially if you do not drink a full glass of water and/or if you lie down within an hour of taking ibandronic acid. If you develop these symptoms, stop taking Ibandronic Acid Mylan and tell your doctor straight away (see section 3).

Children and adolescents

Do not give Ibandronic Acid Mylan to children or adolescents below 18 years.

Other medicines and Ibandronic Acid Mylan

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. Especially:

- **Supplements containing calcium, magnesium, iron, or aluminium** as they could possibly influence the effects of ibandronic acid.
- Acetylsalicylic acid and other non-steroidal anti-inflammatory medicines (NSAIDs) (including ibuprofen, diclofenac sodium and naproxen) may irritate the stomach and intestine. Bisphosphonates (like ibandronic acid) may also do so. So be especially careful if you take painkillers or anti-inflammatories while you're taking ibandronic acid.

After swallowing your monthly Ibandronic Acid Mylan tablet, **wait for 1 hour before taking any other medication**, including indigestion tablets, calcium supplements, or vitamins.

Ibandronic Acid Mylan with food and drink

Do not take Ibandronic Acid Mylan with food. Ibandronic Acid Mylan is less effective if it is taken with food. **You can drink water but no other drinks.**

After you have taken Ibandronic Acid Mylan, please wait for 1 hour before you have your first food and further drinks (see 3. How to take Ibandronic Acid Mylan).

Pregnancy and breast-feeding

Ibandronic Acid Mylan is for use only by postmenopausal women and must not be taken by women who could still have a baby.

Do not take Ibandronic Acid Mylan if you are pregnant or breast-feeding.

Ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

You can drive and use machines, as it is expected that Ibandronic Acid Mylan has no or negligible effect on your ability to drive and use machines.

Ibandronic Acid Mylan contains Lactose

This medicine contains lactose monohydrate. If your doctor has told you that you have an intolerance to some sugars (e.g. if you have a galactose intolerance, the Lapp lactase deficiency or have problems with glucose-galactose absorption), talk to your doctor before taking this medicine.

3. How to take Ibandronic Acid Mylan

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The usual dose of Ibandronic Acid Mylan is one tablet once a month.

Taking your monthly tablet

It's important to follow these instructions carefully. They are designed to help your Ibandronic Acid Mylan tablet reach your stomach quickly, so it's less likely to cause irritation.

- **Take Ibandronic Acid Mylan 150 mg once a month.**
- **Choose one day of the month** that will be easy to remember. You can choose either the same date (such as the 1st of each month) or the same day (such as the first Sunday of each month) to take your Ibandronic Acid Mylan tablet. Choose the date that best fits your routine.
- Take your Ibandronic Acid Mylan tablet **at least 6 hours after you last had anything** to eat or drink except water.

50064555



Description		Ibandronate Sodium 150 mg 1		Date: 02 AUG 2017		Time: 14:53	
Component Type	Leaflet	Pharma Code	5298	No. of colours	1	Page Count	1/2
Affiliate Item Code	1048484	SAP No.	N/A	Colours	Black		
Superseded Affiliate Item Code	785510	Vendor Job No.	317982	Non-Print Colours			
TrackWise PR No.	1048484	Trackwise Proof No.	2	Equate CMYK with			
MA No.	PA0577/111/001	Client Market	Ireland	Main Font	Times New Roman	Body Text Size	9 pt
Packing Site/Printer	Komarom	Keyline/Drawing No.	N/A	Dimensions	170 x 360 mm	Min Text Size used	9 pt
Supplier Code	50064555	Barcode Info	N/A				
Sign-offs							
v2/Oct 2016							

- Take your Ibandronic Acid Mylan tablet
 - **after you first get up for the day**, and
 - **before you have anything to eat or drink** (on an empty stomach)
 - **Swallow your tablet with a full glass of water** (at least 180ml).
- Do not** take your tablet with water with a high concentration of calcium, fruit juice or any other drinks. If there is a concern regarding potentially high levels of calcium in the tap water (hard water) it is advised to use bottled water with a low mineral content.
- **Swallow your tablet whole** – do not chew it, crush it or let it dissolve in your mouth.
 - **For the next hour (60 minutes)** after you have taken your tablet
 - **do not lie down**; if you do not stay upright (standing or sitting), some of the medicine could leak back into your oesophagus
 - **do not eat anything**
 - **do not drink anything** (except water if you need it)
 - **do not take any other medicines**
 - After you have waited for an hour you can have your first food and drink of the day. Once you have eaten it is OK to lie down if you wish, and to take any other medication you need.

Do not take your medicine at bedtime or before you get up for the day.

Continuing to take Ibandronic Acid Mylan

It's important to keep taking Ibandronic Acid Mylan every month, as long as your doctor prescribes it for you. After 5 years of using Ibandronic Acid Mylan, please consult with your doctor whether you should continue to take Ibandronic Acid Mylan.

If you take more Ibandronic Acid Mylan than you should If you've taken more than one tablet by mistake, **drink a full glass of milk and talk to your doctor straight away.**

Do not make yourself vomit, and do not lie down – this could cause Ibandronic Acid Mylan to irritate your oesophagus.

If you forget to take Ibandronic Acid Mylan

- If you forget to take your tablet on the morning of your chosen day, **do not take a tablet later in the day.** Instead, consult your calendar and find out when your next scheduled dose is.
- **If you forget to take your tablet on your chosen day and your next scheduled dose is only 1 to 7 days away: Never take two Ibandronic Acid Mylan tablets within the same week.** You should wait until the next scheduled dose is due and take it as normal then, continue taking one tablet once a month on the scheduled days you've marked on your calendar.
- **If you forget to take your tablet on your chosen day and your next scheduled dose is more than 7 days away:** You should take one tablet the next morning after the day you remember; then, continue taking one tablet once a month on the scheduled days you've marked on your calendar.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Talk to a nurse or a doctor straight away if you notice any of the following serious side effects - you may need urgent medical treatment:

Uncommon (may affect up to 1 in 100 people):

- severe pain in the chest, severe pain after swallowing food or drink, severe nausea, or vomiting, difficulty in swallowing. You may have a severe inflammation of your gullet/food pipe, possibly with sores or constriction of the gullet/food pipe

Rare (may affect up to 1 in 1,000 people):

- itching, swelling of your face, lips, tongue and throat, with difficulty breathing.
- persistent eye pain and inflammation
- new pain, weakness or discomfort in your thigh, hip or groin. You may have early signs of a possible unusual fracture of the thigh bone

Very rare (may affect up to 1 in 10,000 people):

- pain or sore in your mouth or jaw. You may have early signs of severe jaw problems (necrosis (dead bone tissue) in the jaw bone)
- talk to your doctor if you have ear pain, discharge from the ear, and/or an ear infection. These could be signs of bone damage in the ear
- severe skin reactions including itchy red skin rash with raised patches and severe blistering or peeling of the skin
- serious, potentially life-threatening allergic reaction

Other possible side effects

Common (may affect up to 1 in 10 people):

- headache
- heartburn, discomfort in swallowing, stomach or tummy pain (may be due to an inflammation of the stomach), indigestion, nausea, having diarrhoea (loose bowels)
- muscle cramps, stiffness of your joints and limbs
- flu-like symptoms, including fever, shaking and shivering, feeling of discomfort, bone pain and aching muscles and joints. Talk to a nurse or doctor if any effects become troublesome or last more than a couple of days

- rash

Uncommon (may affect up to 1 in 100 people):

- dizziness
- flatulence (farting, feeling bloated)
- back pain
- feeling tired and exhausted
- asthma attacks

Rare (may affect up to 1 in 1,000 people):

- Inflammation of the duodenum (first section of the bowel) causing stomach pain
- hives

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Ibandronic Acid Mylan

Keep this medicine out of the sight and reach of children.

This medicinal product does not require any special storage conditions.

Do not use this medicine after the expiry date which is stated on the carton after "EXP". The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Ibandronic Acid Mylan contains

The active substance is ibandronic acid 150 mg (as sodium monohydrate).

The other ingredients are:

- tablet core: lactose monohydrate, povidone, microcrystalline cellulose, crospovidone, colloidal anhydrous silica, and magnesium stearate
- tablet coat: hydroxypropylcellulose, macrogol and titanium dioxide (E171)
- The **imprinting ink contains** shellac glaze, iron oxide black and propylene glycol

What Ibandronic Acid Mylan looks like and contents of the pack

Your medicine comes as a white film-coated, capsule-shaped, biconvex tablet, with "G" over "I-150" imprinted in black ink on one side and plain on other.

Ibandronic Acid Mylan is available in blister packs of 1, 3, 6 and 12 tablets. Not all pack sizes may be marketed.

Marketing Authorisation Holder

McDermott Laboratories Ltd. T/A Gerard Laboratories
35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13, Ireland.

Manufacturers

McDermott Laboratories Ltd. T/A Gerard Laboratories
35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13, Ireland.

Generics [UK] Ltd., Station Close, Potters Bar, Hertfordshire EN6 1TL, United Kingdom.

Mylan Hungary Kft, H-2900 Komárom, Mylan utca 1, Hungary.

This medicinal product is authorised in the Member States of the EEA under the following names:

Belgium	Ibandronate Acid Mylan 150 mg filmomhulde tabletten
Bulgaria	Ibandronic Acid Mylan 150 mg
Cyprus	Ibandronic Acid Mylan 150 mg
Czech Republic	Ibandronic Acid Mylan 150 mg Film-coated Tablets
Denmark	Ibandronic Acid Mylan
Greece	Ibandronic Acid Mylan 150 mg
Spain	Ácido Ibandrónico 150 mg
Finland	Ibandronic Acid Mylan 150 mg
Ireland	Ibandronic Acid Mylan 150 mg Film-coated Tablets
Italy	Acido IbandronicoMylan 150 mg
Malta	Ibandronic Acid Mylan 150 mg
Poland	Ibandronic Acid Mylan
Portugal	Ácido ibandrónico Mylan 150 mg
Romania	Acid ibandronic Mylan 150 mg
Slovenia	Ibandronska kislina Mylan 150 mg filmsko obložene tablete
Slovak Republic	Ibandronic Acid Mylan 150 mg
The Netherlands	Ibandroninezuur Mylan 150 mg
United Kingdom	Ibandronic Acid Mylan 150 mg

This leaflet was last revised in 08/2017

Mylan

1048484
50064555

Mylan			Date: 02 AUG 2017		Time: 14:53	
Description	Ibandronate Sodium 150 mg 1		No. of colours	1	Page Count	2/2
Component Type	Leaflet	Pharma Code 5298	Colours	Black		
Affiliate Item Code	1048484	SAP No. N/A	Non-Print Colours			
Superseded Affiliate Item Code	785510	Vendor Job No. 317982	Equate CMYK with			
TrackWise PR No.	1048484	Trackwise Proof No. 2	Main Font	Times New Roman	Body Text Size	9 pt
MA No.	PA0577/111/001	Client Market Ireland	Dimensions	170 x 360 mm	Min Text Size used	9 pt
Packing Site/Printer	Komarom	Keyline/Drawing No. N/A				
Supplier Code	50064555	Barcode Info N/A				
Sign-offs						
v2/Oct 2016						