

Package leaflet: Information for the user

HUMULIN® M3 (Mixture 3) KwikPen®

100 IU/ml suspension for injection

(Insulin human)

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Humulin M3 KwikPen is and what it is used for
2. What you need to know before you use Humulin M3 KwikPen
3. How to use Humulin M3 KwikPen
4. Possible side effects
5. How to store Humulin M3 KwikPen
6. Contents of the pack and other information

1. What Humulin M3 KwikPen is and what it is used for

Humulin M3 KwikPen is a pre-filled pen containing the active substance human insulin, which is used to treat diabetes. You get diabetes if your pancreas does not make enough insulin to control the level of glucose in your blood (blood sugar). Humulin M3 is used to control glucose in the long term. It is a premixed suspension of rapid and long acting insulin. Its action is prolonged by the inclusion of protamine sulfate in the suspension.

Your doctor may tell you to use Humulin M3 KwikPen as well as a longer-acting insulin. Each kind of insulin comes with its own patient information leaflet to tell you about it. Do not change your insulin unless your doctor tells you to. Be very careful if you do change insulin. Each type of insulin has a different colour and symbol on the pack and the pen so that you can easily tell the difference.

2. What you need to know before you use Humulin M3 KwikPen

Humulin M3 in pre-filled pen is only suitable for injecting just under the skin. Speak to your doctor if you need to inject your insulin by another method.

Do not use Humulin M3 KwikPen:

- **If you think hypoglycaemia (low blood sugar) is starting.** Further in this leaflet it tells you how to deal with mild hypoglycaemia (see A in Section 4).
- If you are allergic to human insulin or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor, pharmacist, or nurse before using Humulin M3 KwikPen.

- If your blood sugar levels are well controlled by your current insulin therapy, you may not feel the warning symptoms when your blood sugar is falling too low. Warning signs are listed later in this leaflet. You must think carefully about when to have your meals, how often to exercise and how much to do. You must also keep a close watch on your blood sugar levels by testing your blood glucose often.
- A few people who have had hypoglycaemia (low blood sugar) after switching from animal insulin to human insulin have reported that the early warning symptoms were less obvious or different. If you often have hypoglycaemia or have difficulty recognising the symptoms, please discuss this with your doctor.
- If you answer YES to any of the following questions, tell your diabetes specialist nurse, doctor or pharmacist.
 - Have you recently become ill?
 - Do you have trouble with your kidneys or liver?
 - Are you exercising more than usual?
- The amount of insulin you need may also change if you drink alcohol.
- You should also tell your diabetes specialist nurse, doctor or pharmacist if you are planning to go abroad. The time difference between countries may mean that you have to have your injections and meals at different times to when you are at home.
- Some patients with long-standing type 2 diabetes mellitus and heart disease or previous stroke who were treated with pioglitazone and insulin experienced the development of heart failure. Inform your doctor as soon as possible, if you experience signs of heart failure such as unusual shortness of breath or rapid increase in weight or localised swelling (oedema).

Other medicines and Humulin M3 KwikPen

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Your insulin needs may change if you are taking any of the following:

- steroids,
- thyroid hormone replacement therapy,
- oral hypoglycaemics (antidiabetic medication),

- acetylsalicylic acid (aspirin),
- growth hormone,
- octreotide, lanreotide,
- beta₂ stimulants (for example ritodrine, salbutamol or terbutaline),
- beta-blockers,
- thiazides or some antidepressants (monoamine oxidase inhibitors),
- danazol,
- some angiotensin converting (ACE) inhibitors (for example captopril, enalapril) or angiotensin II receptor blockers.

Pregnancy, breast-feeding and fertility

The amount of insulin you need usually falls during the first three months of pregnancy and increases for the remaining six months. If you are breast-feeding, you may need to alter your insulin intake or diet.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before using this medicine.

Driving and using machines

Your ability to concentrate and react may be reduced if you have hypoglycaemia (low blood sugar). Please remember this in all situations where you might put yourself and others at risk (e.g. driving a car or operating machinery). You should contact your diabetes specialist nurse or doctor about the advisability of driving if you have:

- frequent episodes of hypoglycaemia
- reduced or absent warning signs of hypoglycaemia.

Humulin M3 contains Sodium

This medicine contains less than 1 mmol sodium (23 mg) per dose unit, that is to say essentially 'sodium-free'.

3. How to use Humulin M3 KwikPen

Always check the pack and the label of the pre-filled pen for the name and type of the insulin when you get it from your pharmacy. Make sure you get the Humulin KwikPen that your doctor has told you to use.

Always use this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. To prevent the possible transmission of disease, each pen must be used by you only, even if the needle is changed.

Dosage

- Your doctor has told you which insulin to use, how much, when and how often to inject. These instructions are only for you. Follow them exactly and visit your diabetes clinic regularly.
- If you change your insulin type (for example from animal to human), you may have to take more or less than before. This might just be for the first injection or it may be a gradual change over several weeks or months.
- Humulin M3 in pre-filled pen is only suitable for injecting just under the skin. Speak to your doctor if you need to inject your insulin by another method.

Preparing Humulin M3 KwikPen

- The KwikPen should be rotated in the palms of the hands 10 times and inverted 10 times immediately before use to resuspend insulin until it appears uniformly cloudy or milky. If not, repeat the above procedure until contents are mixed. Cartridges in the pen contain a small glass bead to assist mixing. **Do not** shake vigorously as this may cause frothing which may interfere with the correct measurement of the dose. The cartridges in the pen should be examined frequently and should not be used if clumps of material are present or if solid white particles stick to the bottom or sides of the cartridge, giving it a frosted appearance. Check each time you inject yourself.

Getting the KwikPen ready to use (Please see user manual)

- First wash your hands.
- Read the instructions on how to use your pre-filled insulin pen. Please follow the instructions carefully. Here are some reminders.
 - Use a clean needle. (Needles are not included).
 - Prime your KwikPen before each use. This checks that insulin comes out and clears the air bubbles from your KwikPen. There may still be some small air bubbles left in the KwikPen - these are harmless, but if the air bubble is too big, it may make the dose of your injection less accurate.

Injecting Humulin M3

- Before you make an injection, clean your skin as you have been instructed. Inject under the skin, as you were taught. **Do not** inject directly into a vein. After your injection, leave the needle in the skin for 5 seconds to make sure you have taken the whole dose. Do not rub the area you have just injected. Make sure you inject at least half an inch (1 cm) from where you last injected and that you 'rotate' the places you inject, as you have been taught.

After injecting

- As soon as you have done the injection, unscrew the needle from the KwikPen using the outer needle cap. This will keep the insulin sterile and prevent leaking. It will also stop air going back into the KwikPen and the needle clogging up. **Do not share your needles or your KwikPen.** Replace the cap on the KwikPen.

Item Code NL068FSAA02			Previous Item Code (to be destroyed) NL068FSAA01		
PPD Information Box				ALRP Information Box	
Start Date 29 JUN 2018	Proof No.: 1	Printing Colours BLACK	Technical Colours Die Cut	Affiliate Barcode: Type: N/A Code: N/A	Translations of Variable Data
Technical Information: Layout name FEGPA014A05	Size (mm): 210x315	No. of Pages: 1/2	Overt N/A	Other Regulated Elements N/A	lot: N/A
	Folded Size (mm) 210x52.5				mfg date: N/A
BLUE Project No. 1800031378	Sick Code N/A				
Feed Direction: (For labels only)	N/A <input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Further injections

- Every time you use a KwikPen you must use a new needle. Before every injection, clear any air bubbles. You can see how much insulin is left by holding the KwikPen with the needle pointing down. The scale on the cartridge shows about how many units you have left.
- Do not mix any other insulin in your disposable pen. Once the KwikPen is empty, do not use it again. Please get rid of it carefully - your diabetes specialist nurse or pharmacist will tell you how to do this.

If you use more Humulin M3 than you should

If you use more Humulin M3 than you should, your blood sugar may become low. Check your blood sugar (see A in Section 4).

If you forget to use Humulin M3 KwikPen

If you use less Humulin M3 than you should, your blood sugar levels may increase. Check your blood sugar. Do not inject a double dose to make up for a forgotten dose.

If you stop using Humulin M3 KwikPen

If you use less Humulin M3 than you should, your blood sugar levels may become too high. Do not change your insulin unless your doctor tells you to.

If you have any further questions on the use of this medicinal product, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Human insulin may cause hypoglycaemia (low blood sugar). See more information on hypoglycaemia below in the subsection "Common problems of diabetes".

Possible side effects

Systemic allergy is very rare (affects less than 1 person in 10,000). The symptoms are as follows:

- fall in blood pressure
- rash over the whole body
- difficulty in breathing
- wheezing
- fast heart beat
- sweating

If you think you are having this sort of insulin allergy with Humulin M3, tell your doctor at once.

Local allergy is common (affects less than 1 person in 10). Some people get redness, swelling or itching around the area of the insulin injection. This usually clears up in anything from a few days to a few weeks. If this happens to you, tell your doctor.

Lipodystrophy (thickening or pitting of the skin) is uncommon (affects less than 1 person in 100). If you notice your skin thickening or pitting at the injection site, change your injection site and tell your doctor.

Oedema (e.g. swelling in arms, ankles; fluid retention) has been reported, particularly at the start of insulin therapy or during a change in therapy to improve control of your blood glucose.

Common problems of diabetes

A. Hypoglycaemia

Hypoglycaemia (low blood sugar) means there is not enough sugar in the blood. This can be caused if you:

- take too much Humulin M3 or other insulin;
- miss or delay meals or change your diet;
- exercise or work too hard just before or after a meal;
- have an infection or illness (especially diarrhoea or vomiting);
- have a change in your need for insulin; or
- have trouble with your kidneys or liver which gets worse.

Alcohol and some medicines can affect your blood sugar levels.

The first symptoms of low blood sugar usually come on quickly and include the following:

- tiredness
- rapid heartbeat
- nervousness or shakiness
- feeling sick
- headache
- cold sweat

Until you become confident in recognising your warning symptoms, avoid situations such as driving a car, in which you or others would be put at risk by hypoglycaemia.

Do not use Humulin M3 if you think hypoglycaemia (low blood sugar) is starting.

If your blood sugar is low, eat glucose tablets, sugar or drink a sugary drink. Then eat fruit, biscuits, or a sandwich, as your doctor has advised you and have some rest. This will often get you over mild hypoglycaemia or a minor insulin overdose. If you get worse and your breathing is shallow and your skin gets pale, tell your doctor at once. A glucagon injection can treat quite severe hypoglycaemia. Eat glucose or sugar after the glucagon injection. If you do not respond to glucagon, you will have to go to hospital. Ask your doctor to tell you about glucagon.

B. Hyperglycaemia and diabetic ketoacidosis

Hyperglycaemia (too much sugar in the blood) means that your body does not have enough insulin. Hyperglycaemia can be brought about by:

- not taking your Humulin or other insulin;
- taking less insulin than your doctor tells you to;
- eating a lot more than your diet allows; or
- fever, infection or emotional stress.

Hyperglycaemia can lead to diabetic ketoacidosis. The first symptoms come on slowly over many hours or days. The symptoms include the following:

- feeling sleepy
- no appetite
- flushed face
- fruity smell on the breath
- thirst
- feeling or being sick

Severe symptoms are heavy breathing and a rapid pulse. **Get medical help immediately.**

If hypoglycaemia (low blood sugar) or hyperglycaemia (high blood sugar) are not treated they can be very serious and cause headaches, nausea, vomiting, dehydration, unconsciousness, coma or even death.

Three simple steps to avoid hypoglycaemia or hyperglycaemia are:

- Always keep spare syringes and a spare vial of Humulin M3.
- Always carry something to show you are diabetic.
- Always carry sugar with you.

C. Illness

If you are ill, especially if you feel sick or are sick, the amount of insulin you need may change. **Even when you are not eating normally, you still need insulin.** Test your urine or blood, follow your 'sick rules', and tell your diabetes specialist nurse or doctor.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via **Ireland:** HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2, Tel: +353 1 6764971, Fax: +353 1 6762517, Website: www.hpra.ie, e-mail: medsafety@hpra.ie, **United Kingdom:** Yellow Card Scheme, Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Humulin M3 KwikPen

Keep this medicine out of the sight and reach children.

Before the first use store your Humulin M3 KwikPen in a refrigerator (2°C – 8°C). **Do not freeze.** Keep your 'in use' Humulin M3 KwikPen at room temperature (below 30°C) for up to 28 days. Do not keep your 'in use' pen in the fridge. Do not put it near heat or in the sun.

Do not use this medicine after the expiry date which is stated on the label and the carton. The expiry date refers to the last day of that month.

Do not use this medicine if you notice that clumps of material are present or solid white particles stick to the bottom or sides of the cartridge, giving it a frosted appearance. Check this each time you inject yourself.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

6. Contents of the pack and other information

What Humulin M3 KwikPen contains

- The **active substance** is human insulin. Human insulin is made in the laboratory by a 'recombinant DNA technology' process. It has the same structure as the natural hormone made by the pancreas. It is therefore different from animal insulins. Humulin M3 is a premixed suspension which contains 30% insulin dissolved in water and 70% insulin with protamine sulfate.

- The **other ingredients** are protamine sulfate, metacresol, phenol, glycerol, dibasic sodium phosphate 7H₂O, zinc oxide and water for injection. Sodium hydroxide or hydrochloric acid may have been used during manufacture to adjust the acidity.

What Humulin M3 KwikPen looks like and contents of the pack

Humulin M3 (Mixture 3) KwikPen 100 IU/ml suspension for injection is a white, sterile suspension and contains 100 units of insulin in each millilitre (100 IU/ml).

Each Humulin M3 KwikPen contains 300 units (3 millilitres).

The Humulin M3 KwikPen comes in a pack of 5.

The Humulin M3 in your KwikPen is the same as the Humulin M3, which comes in separate Humulin M3 cartridges. The KwikPen simply has a built in cartridge. When the KwikPen is empty you cannot use it again.

Marketing Authorisation Holder and Manufacturer

Humulin M3 KwikPen is made by: Lilly France S.A.S., Rue du Colonel Lilly, 67640 Fegersheim, France.

The marketing authorisation is held by:

United Kingdom

Eli Lilly and Company Limited, Lilly House, Priestley Road, Basingstoke, Hampshire RG24 9NL, U.K.

Ireland

Eli Lilly Nederland B.V., Papendorpseweg 83, 3528 BJ Utrecht, The Netherlands

This medicinal product is authorised in the Member States of the EEA under the following names:

Huminsulin "Lilly" Profil III KwikPen (Austria)

Humuline 30/70 KwikPen (Belgium, Luxembourg)

Huminsulin Profil III KwikPen (Germany)

Humulina 30:70 KwikPen (Spain)

Humulin M3 KwikPen (Greece, Ireland, Lithuania, Romania, Slovenia, UK)

Umuline profil 30 KwikPen 100 UI/ml (France)

Humulin M3 (30/70) KwikPen (Czech Republic, Slovakia)

This leaflet was last revised in June 2018.

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

Ireland

Eli Lilly and Company (Ireland) Limited, Tel: + 353-(0) 1 661 4377

United Kingdom

Eli Lilly and Company Limited, Tel: + 44-(0) 1256 315000

NL068FSA02

Item Code NL068FSA02		Previous Item Code (to be destroyed) NL068FSA01				
PPD Information Box			ALRP Information Box			
Start Date 29 JUN 2018	Proof No.: 1	Printing Colours BLACK	Technical Colours Die Cut	Affiliate Barcode: Type: N/A Code: N/A	Translations of Variable Data	
Technical Information: Layout name FEGPA014A05	Size (mm): 210x315			Other Regulated Elements N/A	lot: N/A	
	Folded Size (mm) 210x52.5				mfg date: N/A	
	No. of Pages: 2/2		Overt N/A		exp date: N/A	
BLUE Project No. 1800031378	Sick Code N/A				price: N/A	
Feed Direction: (For labels only)	N/A	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	other: N/A