

## Package leaflet: Information for the user

**Gabapentin Aurobindo 100 mg Capsules, hard**  
**Gabapentin Aurobindo 300 mg Capsules, hard**  
**Gabapentin Aurobindo 400 mg Capsules, hard**  
gabapentin

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### **What is in this leaflet:**

1. What Gabapentin Aurobindo is and what it is used for
2. What you need to know before you take Gabapentin Aurobindo
3. How to take Gabapentin Aurobindo
4. Possible side effects
5. How to store Gabapentin Aurobindo
6. Contents of the pack and other information

## **1. What Gabapentin Aurobindo is and what it is used for**

Gabapentin belongs to a group of medicines used to treat epilepsy and peripheral neuropathic pain (long lasting pain caused by damage to the nerves).

The active substance in Gabapentin Aurobindo is gabapentin.

**Epilepsy:** Gabapentin Aurobindo is used to treat various forms of epilepsy (seizures that are initially limited to certain parts of the brain, whether the seizure spreads to other parts of the brain or not). The doctor treating you or your child 6 years of age and older will prescribe Gabapentin to help treat epilepsy when the current treatment is not fully controlling the condition. You or your child 6 years of age and older should take Gabapentin in addition to the current treatment unless told otherwise. Gabapentin can also be used on its own to treat adults and children over 12 years of age.

**Peripheral neuropathic pain:** Gabapentin Aurobindo is used to treat long lasting pain caused by damage to the nerves. A variety of different diseases can cause peripheral neuropathic pain (primarily occurring in the legs and/or arms), such as diabetes or shingles. Pain sensations may be described as hot, burning, throbbing, shooting, stabbing, sharp, cramping, aching, tingling, numbness, pins and needles etc.

## **2. What you need to know before you take Gabapentin Aurobindo**

**Do not take Gabapentin Aurobindo:**

- if you are allergic (hypersensitive) to gabapentin or any of the other ingredients of this medicine (listed in section 6).

### **Warnings and precautions**

Talk to your doctor or pharmacist before taking Gabapentin Aurobindo if:

- you suffer from kidney problems, your doctor may prescribe a different dosing schedule
- you are on haemodialysis (to remove waste products because of kidney failure), tell your doctor if you develop muscle pain and/or weakness
- you develop signs such as persistent stomach pain, feeling sick and being sick contact your doctor immediately as these may be symptoms of acute pancreatitis (an inflamed pancreas)
- if you have nervous system disorders, respiratory disorders, or you are more than 65 years old, your doctor may prescribe you a different dosing regimen

Cases of abuse and dependence have been reported for gabapentin from the post-marketing experience. Talk to your doctor if you have a history of abuse or dependence.

A small number of people being treated with anti-epileptics such as gabapentin have had thoughts of harming or killing themselves. If at any time you have these thoughts, immediately contact your doctor.

### **Important information about potentially serious reactions**

A small number of people taking Gabapentin get an allergic reaction or potentially serious skin reaction, which may develop into more serious problems if they are not treated. You need to know the symptoms to look out for while you are taking Gabapentin.

**Read the description of these symptoms in section 4 of this leaflet** under ‘Contact your doctor immediately if you experience any of the following symptoms after taking this medicine as they can be serious’.

Muscle weakness, tenderness or pain and particularly, if at the same time, you feel unwell or have a high temperature it may be caused by an abnormal muscle breakdown which can be life-threatening and lead to kidney problems. You may also experience discoloration of your urine, and a change in blood test results (notably blood creatine phosphokinase increased). If you experience any of these signs or symptoms, please contact your doctor immediately.

### **Other medicines and Gabapentin Aurobindo**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. In particular, tell your doctor (or pharmacist) if you are taking or have been recently taking any medicines for convulsions, sleeping disorders, depression, anxiety, or any other neurological or psychiatric problems.

#### Medicines containing opioids such as morphine

If you are taking any medicines containing opioids (such as morphine), please tell your doctor or pharmacist as opioids may increase the effect of Gabapentin. In addition, combination of Gabapentin with opioids may cause sleepiness, sedation, decrease in breathing, or death.

#### Antacids for indigestion

If Gabapentin and antacids containing aluminium and magnesium are taken at the same time, absorption of Gabapentin from the stomach may be reduced. It is therefore recommended that Gabapentin is taken at the earliest two hours after taking an antacid.

Gabapentin is not expected to interact with other antiepileptic drugs or the oral contraceptive pill.

Gabapentin may interfere with some laboratory tests, if you require a urine test tell your doctor or hospital what you are taking.

### **Gabapentin Aurobindo with food**

Gabapentin can be taken with or without food.

### **Pregnancy and breast-feeding and fertility**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

#### **Pregnancy**

Gabapentin Aurobindo should not be taken during pregnancy, unless you are told otherwise by your doctor. Effective contraception must be used by women of child-bearing potential.

There have been no studies specifically looking at the use of gabapentin in pregnant women, but other medications used to treat seizures have reported an increased risk of harm to the developing baby, particularly when more than one seizure medication is taken at the same time. Therefore, whenever possible, you should try to take only one seizure medication during pregnancy and only under the advice of your doctor.

Contact your doctor immediately if you become pregnant, think you might be pregnant or are planning to become pregnant while taking Gabapentin Aurobindo. Do not suddenly discontinue taking this medicine as this may lead to a breakthrough seizure, which could have serious consequences for you and your baby.

#### **Breast-feeding**

Gabapentin Aurobindo is passed on through human milk. Because the effect on the baby is unknown, it is not recommended to breast-feed while using gabapentin.

#### **Fertility**

There is no effect on fertility in animal studies.

#### **Driving and using machines**

Gabapentin Aurobindo may produce dizziness, drowsiness and tiredness. You should not drive, operate complex machinery or take part in other potentially hazardous activities until you know whether this medication affects your ability to perform these activities.

#### **Gabapentin Aurobindo contains lactose and sodium**

Gabapentin Aurobindo capsules contain lactose (a type of sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

### **3. How to take Gabapentin Aurobindo**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor will determine what dose is appropriate for you.

Gabapentin Aurobindo is for oral use. Always swallow the capsules whole with plenty of water.

This medicine may be taken with or without food.

Continue taking gabapentin Aurobindo until your doctor tells you to stop.

**Epilepsy, the recommended dose is:**

***Adults and adolescents:***

Take the number of capsules as instructed. Your doctor will usually build up your dose gradually. The starting dose will generally be between 300 mg and 900 mg each day. Thereafter, the dose may be increased as instructed by your doctor up to a maximum of 3600 mg each day and your doctor will tell you to take this in 3 separate doses, i.e. once in the morning, once in the afternoon and once in the evening.

***Children aged 6 years and above:***

The dose to be given to your child will be decided by your doctor as it is calculated against your child's weight. The treatment is started with a low initial dose which is gradually increased over a period of approximately 3 days. The usual dose to control epilepsy is 25-35 mg per kg per day. It is usually given in 3 separate doses, by taking the capsule(s) each day, usually once in the morning, once in the afternoon and once in the evening.

Gabapentin is not recommended for use in children below 6 years of age.

**Peripheral Neuropathic Pain, the recommended dose is:**

***Adults:***

Take the number of capsules as instructed by your doctor. Your doctor will usually build up your dose gradually. The starting dose will generally be between 300 mg and 900 mg each day. Thereafter, the dose may be increased as instructed by your doctor up to a maximum of 3600 mg each day and your doctor will tell you to take this in 3 separate doses, i.e. once in the morning, once in the afternoon and once in the evening.

**If you have kidney problems or are receiving haemodialysis**

Your doctor may prescribe a different dosing schedule and/or dose if you have problems with your kidneys or are undergoing haemodialysis.

**If you are an elderly patient (over 65 years of age)**

You should take the normal dose of Gabapentin unless you have problems with your kidneys. Your doctor may prescribe a different dosing schedule and/or dose if you have problems with your kidneys.

If you have the impression that the effect of Gabapentin Aurobindo is too strong or too weak, talk to your doctor or pharmacist as soon as possible.

**If you take more Gabapentin Aurobindo than you should**

Higher than recommended doses may result in an increase in side effects including loss of consciousness, dizziness, double vision, slurred speech, drowsiness and diarrhoea. Call your

doctor or go to the nearest hospital emergency unit immediately if you take more Gabapentin Aurobindo than your doctor prescribed. Take along any capsules that you have not taken, together with the container and the label so that the hospital can easily tell what medicine you have taken.

#### **If you forget to take Gabapentin Aurobindo**

If you forget to take a dose, take it as soon as you remember unless it is time for your next dose. Do not take a double dose to make up for a forgotten dose.

#### **If you stop taking Gabapentin Aurobindo**

Do not stop taking Gabapentin Aurobindo unless your doctor tells you to. If your treatment is stopped it should be done gradually over a minimum of 1 week. If you stop taking Gabapentin Aurobindo suddenly or before your doctor tells you, there is an increased risk of seizures.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Contact your doctor immediately if you experience any of the following symptoms after taking this medicine as they can be serious:**

- **severe skin reactions that require immediate attention, swelling of the lips and face, skin rash and redness, and/or hair loss (these may be symptoms of a serious allergic reaction)**
- **persistent stomach pain, feeling sick and being sick as these may be symptoms of acute pancreatitis (an inflamed pancreas)**
- **breathing problems, which if severe you may need emergency and intensive care to continue breathing normally**
- **Gabapentin Aurobindo may cause a serious or life-threatening allergic reaction that may affect your skin or other parts of your body such as your liver or blood cells. You may or may not have rash when you get this type of reaction. It may cause you to be hospitalised or to stop gabapentin. Call your doctor right away if you have any of the following symptoms:**
  - **skin rash**
  - **hives**
  - **fever**
  - **swollen glands that do not go away**
  - **swelling of your lip and tongue**
  - **yellowing of your skin or of the whites of the eyes**
  - **unusual bruising or bleeding**
  - **severe fatigue or weakness**
  - **unexpected muscle pain**
  - **frequent infections**

**These symptoms may be the first signs of a serious reaction. A doctor should examine you to decide if you should continue taking gabapentin aurobindo.**

- If you are on haemodialysis, tell your doctor if you develop muscle pain and/or weakness.

Other side effects include:

**Very common (may affect more than 1 in 10 people):**

- viral infection
- feeling drowsy, dizziness, lack of coordination
- feeling tired, fever

**Common (may affect more than 1 in 100 people):**

- pneumonia, respiratory infections, urinary tract infection, inflammation of the ear or other infections
- low white blood cell counts
- Anorexia, increased appetite
- anger towards others, confusion, mood changes, depression, anxiety, nervousness, difficulty with thinking
- convulsions, jerky movements, difficulty with speaking, loss of memory, tremor, difficulty sleeping, headache, sensitive skin, decreased sensation (numbness), difficulty with coordination, unusual eye movement, increased, decreased or absent reflexes
- blurred vision, double vision
- vertigo
- high blood pressure, flushing or dilation of blood vessels
- difficulty breathing, bronchitis, sore throat, cough, dry nosevomiting (being sick), nausea (feeling sick), problems with teeth, inflamed gums, diarrhoea, stomach pain, indigestion, constipation, dry mouth or throat, flatulence
- facial swelling, bruises, rash, itch, acne
- joint pain, muscle pain, back pain, twitching
- difficulties with erection (impotence)
- swelling in the legs and arms, difficulty with walking, weakness, pain, feeling unwell, flu-like symptoms
- decrease in white blood cells, increase in weight
- accidental injury, fracture, abrasion

Additionally in clinical studies in children aggressive behaviour and jerky movements were reported commonly.

**Uncommon (may affect up to 1 in 100 people):**

- agitation (a state of chronic restlessness and unintentional and purposeless motions)
- allergic reactions such as hives
- decreased movement
- racing heartbeat
- difficulty swallowing
- swelling that may involve the face, trunk and limbs
- abnormal blood test results suggesting problems with the liver
- mental impairment
- fall
- Increase in blood glucose levels (most often observed in patients with diabetes)

**Rare (may affect up to 1 in 1,000 people)**

- loss of consciousness
- decrease in blood glucose levels (most often observed in patients with diabetes)
- trouble breathing, shallow breaths (respiratory depression)

**After marketing Gabapentin Aurobindo the following side effects have been reported:**

- decreased platelets (blood clotting cells)
- suicidal thoughts, hallucinations
- problems with abnormal movements such as writhing, jerking movements and stiffness
- ringing in the ears
- a group of side effects that could include swollen lymph nodes (isolated small raised lumps under the skin), fever, rash, and inflammation of liver occurring together
- yellowing of the skin and eyes (jaundice), inflammation of the liver
- acute kidney failure, incontinence
- increased breast tissue, breast enlargement
- adverse events following the abrupt discontinuation of gabapentin (anxiety, difficulty sleeping, feeling sick, pain, sweating), chest pain
- breakdown of muscle fibers (rhabdomyolysis)
- change in blood test results (creatinine phosphokinase increased)
- Problems with sexual functioning including inability to achieve a sexual climax, delayed ejaculation
- low blood sodium level
- anaphylaxis (serious, potentially life threatening allergic reaction including difficulty breathing, swelling of the lips, throat, and tongue, and hypotension requiring emergency treatment)

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, website: [www.hpra.ie](http://www.hpra.ie). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Gabapentin Aurobindo**

Keep this medicine out of the sight and reach of children.

Do not store above 25°C.

Do not use this medicine after the expiry date which is stated on the blister and the outer carton after “EXP”. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Gabapentin Aurobindo contains**

The active substance is gabapentin.

Each hard capsule contains either 100 mg, 300 mg or 400 mg gabapentin.

The other ingredients are:

Capsule contents: lactose (see section 2, ‘Gabapentin contains lactose’), maize starch and talc.

Capsule shell: gelatin, titanium dioxide (E171) and sodium lauryl sulphate.

The 300 mg capsules also contain yellow iron oxide (E172).

The 400 mg capsules also contain red and yellow iron oxides (E172)

**What Gabapentin Aurobindo looks like and contents of the pack**

Capsule, hard. The 100 mg capsules are white, the 300 mg are yellow and the 400 mg capsules are orange.

Each capsule contains a white powder.

Gabapentin Aurobindo is available in blister packs of 10, 20, 30, 50, 90, 100, 200 (2x100), 250, 500 (5x100) and 1,000 (10x100) capsules and unit dose blister packs containing 20x1, 60x1 and 100x1 capsules.

Gabapentin Aurobindo is also available in bottles containing 50, 100 and 250 capsules.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder**

Aurobindo Pharma (Malta) Limited, Vault 14, Level 2, Valletta Waterfront, Floriana FRN 1913, Malta

**Manufacturer**

Mylan Hungary Kft., Mylan utca 1, Komarom 2900, Hungary

McDermott Laboratories Ltd. t/a Gerard Laboratories, 35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13.

**This leaflet was last revised in: April 2022**