

**Reminyl® XL 8 mg prolonged-release capsules, hard**  
**Reminyl® XL 16 mg prolonged-release capsules, hard**  
**Reminyl® XL 24 mg prolonged-release capsules, hard**  
galantamine

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet:**

1. What Reminyl XL is and what it is used for
2. What you need to know before you take Reminyl XL
3. How to take Reminyl XL
4. Possible side effects
5. How to store Reminyl XL
6. Contents of the pack and other information

**1. What Reminyl XL is and what it is used for**

Reminyl XL contains the active substance 'galantamine' an antimentia medicine. It is used in adults to treat the symptoms of mild to moderately severe Alzheimer's disease, a type of dementia that alters brain function.

Alzheimer's disease causes increasing memory loss, confusion and behavioural changes which make it increasingly difficult to carry out normal daily activities.

These effects are thought to be caused by a lack of 'acetylcholine', a substance responsible for sending messages between brain cells. Reminyl increases the amount of acetylcholine in the brain and treats the signs of the disease.

The capsules are made in a 'prolonged-release' form. This means that they release the medicine slowly.

**2. What you need to know before you take Reminyl XL**

**Do not take Reminyl XL**

- if you are allergic to galantamine or to any of the other ingredients of this medicine (listed in section 6)
- if you have severe liver or severe kidney disease

**Warnings and precautions**

Talk to your doctor or pharmacist before taking Reminyl XL. This medicine is only used in Alzheimer's disease, and is not recommended for other types of memory loss or confusion.

**Serious side effects**

Reminyl XL can cause serious skin reactions, heart problems and fits (seizures). You must be aware of these side effects while you are taking Reminyl. See 'Look out for serious side effects' in section 4.

**Before you take Reminyl XL**, your doctor needs to know if you have, or have had, any of the following:

- liver or kidney problems
- a heart condition (such as chest discomfort that is often brought on by physical activity, a heart attack, heart failure, slow or uneven heart beat, prolonged QTc interval)
- changes in 'electrolyte' levels (naturally occurring chemicals in the blood, such as potassium)
- a peptic (stomach) ulcer
- blockage of the stomach or intestines
- a disorder of the nervous system (such as epilepsy or problems controlling movements of the body or limbs (extrapyramidal disorder))
- a respiratory disease or infection that affects breathing (such as asthma, obstructive pulmonary disease, or pneumonia)
- problems passing urine.

Your doctor will decide if Reminyl XL is suitable for you, or if the dose needs to be changed.

**Also tell your doctor if you recently had an operation** on the stomach, intestines or bladder. Your doctor may decide that Reminyl XL is not suitable for you.

**Reminyl XL can cause weight loss.** Your doctor will check your weight regularly while you are taking Reminyl XL.

**Children and adolescents**

Reminyl XL is not recommended for children and adolescents.

**Other medicines and Reminyl XL**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Reminyl XL should not be used with medicines that work in a similar way. These include:

- donepezil or rivastigmine (for Alzheimer's disease)
- ambenonium, neostigmine or pyridostigmine (for severe muscle weakness)
- pilocarpine (when taken by mouth for dry mouth or dry eyes).

Some medicines can make side effects more likely in people taking Reminyl XL. These include:

- medicines affecting the QTc interval
- paroxetine or fluoxetine (antidepressants)
- quinidine (for uneven heart beat)
- ketoconazole (an antifungal)
- erythromycin (an antibiotic)
- ritonavir (for human immunodeficiency virus or 'HIV').

- non-steroidal anti-inflammatory painkillers (such as ibuprofen) which can increase the risk of ulcers
- medicines taken for heart conditions or high blood pressure (such as digoxin, amiodarone, atropine, beta-blockers, or calcium channel blocking agents). If you take medicines for an uneven heart-beat, your doctor may check your heart using an electrocardiogram (ECG).

Your doctor may give you a lower dose of Reminyl XL if you are taking some of these medicines.

Reminyl XL may affect some anaesthetics. If you are going to have an operation under a general anaesthetic, tell the doctor that you are taking Reminyl, well in advance.

If you have any questions, talk to your doctor or pharmacist for advice.

**Pregnancy and breast-feeding**

If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

**You should not breast-feed while you are taking Reminyl XL.**

**Driving and using machines**

Reminyl XL may make you feel dizzy or sleepy, especially during the first few weeks of treatment. If Reminyl XL affects you, do not drive or use any tools or machinery.

**Reminyl XL contains sucrose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

**3. How to take Reminyl XL capsules**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

If you are currently taking Reminyl tablets or oral solution and have been told by your doctor to switch to Reminyl XL prolonged-release capsules, carefully read the instructions under 'Switching from taking Reminyl tablets or oral solution to Reminyl capsules' in this section.

**How much to take**

You will start treatment with Reminyl XL at a low dose. The usual starting dose is 8 mg, taken once a day. Your doctor may gradually increase your dose, every 4 weeks or more until you reach a dose that is suitable for you. The maximum dose is 24 mg, taken once a day.

Your doctor will explain what dose to start with and when the dose should be increased. If you are not sure what to do, or find the effect of Reminyl XL is too strong or too weak, talk to your doctor or pharmacist.

Your doctor will need to see you regularly, to check that this medicine is working and to discuss how you are feeling.

**If you have liver or kidney problems**, your doctor may give you a reduced dose of Reminyl XL, or may decide this medicine is not suitable for you.

**Switching from taking Reminyl tablets or oral solution to Reminyl XL capsules**

If you are currently taking Reminyl tablets or oral solution, your doctor may decide you should switch to Reminyl XL prolonged-release capsules. If this applies to you:

- Take your last dose of Reminyl tablets or oral solution in the evening
- The next morning, take your first dose of Reminyl XL prolonged-release capsule.

**DO NOT take more than one capsule in a day. While you are taking once daily Reminyl XL capsules, DO NOT take Reminyl tablets or oral solution.**

**How to take**

Reminyl XL capsules must be swallowed whole and NOT chewed or crushed. Take your dose of Reminyl XL once a day in the morning, with water or other liquids. Try to take Reminyl XL with food.

Drink plenty of liquids while you are taking Reminyl XL, to keep yourself hydrated.

**If you take more Reminyl XL than you should**

If you take too much Reminyl XL, contact a doctor or hospital straight away. Take any remaining capsules and the packaging with you. The signs of overdose may include,

- severe nausea and vomiting
- weak muscles, slow heart beat, fits (seizures) and loss of consciousness.

**If you forget to take Reminyl XL**

If you forget to take one dose, miss out the forgotten dose completely and take the next dose at the normal time. **Do not take a double dose to make up for a forgotten dose.**

If you forget to take more than one dose, contact your doctor.

**If you stop taking Reminyl XL**

Check with your doctor before you stop taking Reminyl XL. It is important to continue taking this medicine to treat your condition.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Look out for serious side effects**

**Stop taking Reminyl and see a doctor or go to your nearest emergency department immediately** if you notice any of the following;

**Skin reactions**, including:

- Severe rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (Stevens-Johnson syndrome)
- Red rash covered with small pus-filled bumps that can spread over the body, sometimes with a fever (acute generalised exanthematous pustulosis)
- Rash that may blister, with spots that look like small targets

These skin reactions are rare in people taking Reminyl (may affect up to 1 in 1,000 people).

- **Heart problems**, including changes in heart beat (such as a slow beat, extra beats) or palpitations (heart beat feels fast or uneven). Heart problems may show as an abnormal tracing on an electrocardiogram (ECG), and can be common in people taking Reminyl XL (may affect up to 1 in 10 people)

- **Fits (seizures).** These are uncommon in people taking Reminyl XL (may affect up to 1 in 100 people).

**You must stop taking Reminyl and get help immediately** if you notice any of the side effects above.

#### Other side effects:

**Very common side effects:** (may affect more than 1 in 10 people).

- Nausea and vomiting. These side effects are more likely to happen in the first few weeks of treatment or when the dose is increased. They tend to disappear gradually as the body gets used to the medicine and generally only last for a few days. If you have these effects, your doctor may recommend that you drink more liquids, and may prescribe a medicine to stop you being sick.

**Common side effects:** (may affect up to 1 in 10 people).

- Decreased appetite; weight loss
- Seeing, feeling, or hearing things that are not there (hallucinations)
- Depression
- Feeling dizzy or fainting
- Muscle tremors or spasms
- Headache
- Feeling very tired, weak or generally unwell
- Feeling very sleepy with low energy
- High blood pressure
- Stomach pain or discomfort
- Diarrhoea
- Indigestion
- Falls
- Wounds

**Uncommon side effects** (may affect up to 1 in 100 people)

- Allergic reaction
- Not enough water in the body (dehydration)
- Tingling or numb feeling of the skin (pins and needles)
- Change in sense of taste
- Daytime sleepiness
- Problems controlling movements of the body or limbs (extrapyramidal disorder)
- Blurred vision
- Ringing in the ears that does not go away (tinnitus)
- Low blood pressure
- Flushing
- Feeling the need to vomit (retch)
- Excessive sweating
- Weak muscles
- Increased level of liver enzymes in the blood

**Rare side effects** (may affect up to 1 in 1,000 people)

- Inflamed liver (hepatitis).

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below).

HPRA Pharmacovigilance, website: [www.hpra.ie](http://www.hpra.ie).

By reporting side effects you help provide more information on the safety of this medicine.

#### 5. How to store Reminyl XL

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister foil after EXP. The expiry date refers to the last day of that month.

Do not store above 30 °C

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

#### 6. Contents of the pack and other information

##### What Reminyl XL contains:

The active substance is galantamine.

Each prolonged release capsule contains 8 mg, 16 mg or 24 mg galantamine (as hydrobromide)

The other ingredients are:

Diethyl phthalate, ethylcellulose, hypromellose, macrogol 400, maize starch and sucrose, gelatine, titanium dioxide (E171), iron oxide black (E172), shellac, propylene glycol (E1520).

The 16 mg and 24 mg capsules also contain iron oxide red (E172).

The 24 mg capsule also contains iron oxide yellow (E172).

##### What Reminyl XL looks like and contents of the pack

Reminyl XL 8 mg capsules are white and marked 'G8'.

Reminyl XL 16 mg capsules are pink and marked 'G16'.

Reminyl XL 24 mg capsules are caramel and marked 'G24'.

The 8 mg, 16 mg and 24 mg capsules are available in blister packs of 28 capsules.

##### Manufacturer

Janssen-Cilag SpA, Via C. Janssen, 04100 Borgo San Michele, Latina, Italy or  
McGregor Cory Ltd, Exel, Site 1, Middleton Close, Banbury, Oxfordshire, OX16 4RS, UK

##### Product procured from within the EU, repackaged and distributed by the Parallel Product Authorisation Holder:

PCO Manufacturing Ltd., Unit 10, Ashbourne Business Park, Rath, Ashbourne, Co. Meath, Ireland.

##### Parallel Product Authorisation Number

Reminyl XL 8 mg prolonged-release capsules PPA 465/166/2

Reminyl XL 16 mg prolonged-release capsules PPA 465/166/4

Reminyl XL 24 mg prolonged-release capsules PPA 465/166/3

Reminyl is a registered trademark of Shire Pharmaceutical Development Ltd.

This medicinal product is authorised in the Member States of the EEA under the following names:

Austria	Reminyl 8 mg, 16 mg, 24 mg Retardkapseln
Belgium	Reminyl 8 mg, 16 mg, 24 mg harde capsules met verlengde afgifte
Denmark	Reminyl 8 mg, 16 mg, 24 mg depotkapsler, hårde
Finland	Reminyl 8 mg, 16 mg, 24 mg depotkapseli, kova
France	Reminyl LP 8 mg, 16 mg, 24 mg, gélule à libération prolongée
Germany	REMINYL 1x täglich 8 mg, 16 mg, 24 mg Hartkapseln, retardiert
Greece	Reminyl 8mg, 16 mg, 24 mg καψάκια παρατεταμένης αποδέσμευσης, σκληρά
Ireland	Reminyl XL 8 mg, 16 mg, 24 mg prolonged release capsules
Italy	Reminyl 8 mg, 16 mg, 24 mg capsule rigide a rilascio prolungato
Luxembourg	Reminyl 8 mg, 16 mg, 24 mg, gélules dures à libération prolongée
Netherlands	Reminyl retard capsules 1x daags 8 mg, 16 mg, 24 mg, harde capsules met verlengde afgifte
Norway	Reminyl 8 mg, 16 mg, 24 mg depotkapsler, hard
Portugal	Reminyl 8 mg, 16 mg, 24 mg cápsulas de libertação prolongada
Spain	Reminyl 8 mg, 16 mg, 24 mg cápsulas duras de liberación prolongada
Sweden	Reminyl 8 mg, 16 mg, 24 mg depotkapslar, hårda
UK	Reminyl XL 8 mg, 16 mg, 24 mg prolonged release capsules

#### More Information

If you live in the Republic of Ireland, further information, advice and support is available from:

The Alzheimer Society of Ireland, Alzheimer House, 43 Northumberland Avenue, Dun Laoghaire, Co. Dublin. Telephone: (01) 284 6616, Fax: (01) 284 6030, email [info@alzheimer.ie](mailto:info@alzheimer.ie), National Helpline (open Monday to Thursday 10am -4pm): 1 800 341 341 or Western Alzheimer Foundation, Mount Street, Claremorris, Co. Mayo. Telephone: 094 624 80. Fax: 094 62560.

**This leaflet was last revised in May2022**