

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Olmesartan /Amlodipine Mylan 20 mg/ 5 mg film coated tablets

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 20 mg of olmesartan medoxomil and 5 mg of amlodipine (as amlodipine besilate).

### Excipients with known effects

Each Olmesartan /Amlodipine Mylan 20 mg/5 mg film-coated tablet contains 1.0 mg (0.04 mmol) of sodium.

For the full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Film-coated tablet

white, round, biconvex, film-coated tablet, approximately 7.4 mm diameter and debossed with 'M' on one side and 'OA1' on the other side.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic Indications

Treatment of essential hypertension.

Olmesartan /Amlodipine Mylan is indicated in adult patients whose blood pressure is not adequately controlled on olmesartan medoxomil or amlodipine monotherapy (see sections 4.2 and 5.1).

### 4.2 Posology and method of administration

#### Posology

##### Adults

The recommended dosage of Olmesartan /Amlodipine Mylan is 1 tablet per day.

Olmesartan /Amlodipine Mylan 20 mg/5 mg may be administered in patients whose blood pressure is not adequately controlled by 20 mg olmesartan medoxomil or 5 mg amlodipine alone.

A step-wise titration of the dosage of the individual components is recommended before changing to the fixed combination. When clinically appropriate, direct change from monotherapy to the fixed combination may be considered.

For convenience, patients receiving olmesartan medoxomil and amlodipine from separate tablets may be switched to Olmesartan /Amlodipine Mylan tablets containing the same component doses.

Olmesartan /Amlodipine Mylan can be taken with or without food.

##### Elderly (age 65 years or over)

No adjustment of the recommended dose is generally required for elderly people but increase of the dosage should take place with care (see sections 4.4 and 5.2).

If up-titration to the maximum dose of 40 mg olmesartan medoxomil daily is required, blood pressure should be closely monitored.

##### Renal impairment

The maximum dose of olmesartan medoxomil in patients with mild to moderate renal impairment (creatinine clearance of 20 – 60 mL/min) is 20 mg olmesartan medoxomil once daily, owing to limited experience of higher dosages in this patient group.

The use of Olmesartan /Amlodipine Mylan in patients with severe renal impairment (creatinine clearance < 20 mL/min) is not recommended (see sections 4.4 and 5.2).

Monitoring of potassium levels and creatinine is advised in patients with moderate renal impairment.

#### Hepatic impairment

Olmesartan /Amlodipine Mylan should be used with caution in patients with mild to moderate hepatic impairment (see sections 4.4 and 5.2).

In patients with moderate hepatic impairment, an initial dose of 10 mg olmesartan medoxomil once daily is recommended and the maximum dose should not exceed 20 mg once daily. Close monitoring of blood pressure and renal function is advised in hepatically-impaired patients who are already receiving diuretics and/or other antihypertensive agents. There is no experience of olmesartan medoxomil in patients with severe hepatic impairment.

As with all calcium antagonists, amlodipine's half-life is prolonged in patients with impaired liver function and dosage recommendations have not been established. Olmesartan /Amlodipine Mylan should therefore be administered with caution in these patients. The pharmacokinetics of amlodipine have not been studied in severe hepatic impairment. Amlodipine should be initiated at the lowest dose and titrated slowly in patients with impaired liver function. Use of Olmesartan /Amlodipine Mylan in patients with severe hepatic impairment is contraindicated (see section 4.3).

#### Paediatric population

The safety and efficacy of Olmesartan /Amlodipine Mylan in children and adolescents below 18 years has not been established. No data are available.

#### Method of administration

The tablet should be swallowed with a sufficient amount of fluid (e.g. one glass of water). The tablet should not be chewed and should be taken at the same time each day.

### **4.3 Contraindications**

Hypersensitivity to the active substances, to dihydropyridine derivatives or to any of the excipients listed in section 6.1.

Second and third trimesters of pregnancy (see sections 4.4 and 4.6).

Severe hepatic insufficiency and biliary obstruction (see section 5.2).

The concomitant use of Olmesartan /Amlodipine Mylan with aliskiren-containing products is contraindicated in patients with diabetes mellitus or renal impairment (GFR < 60 mL/min/1.73 m<sup>2</sup>) (see sections 4.5 and 5.1).

Due to the component amlodipine Olmesartan /Amlodipine Mylan is also contraindicated in patients with:

- severe hypotension.
- shock (including cardiogenic shock).
- obstruction of the outflow tract of the left ventricle (e.g. high grade aortic stenosis).
- haemodynamically unstable heart failure after acute myocardial infarction

### **4.4 Special warnings and precautions for use**

#### Patients with hypovolaemia or sodium depletion:

Symptomatic hypotension may occur in patients who are volume and/or sodium depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea or vomiting, especially after the first dose. Correction of this condition prior to administration of Olmesartan /Amlodipine Mylan or close medical supervision at the start of the treatment is recommended.

#### Other conditions with stimulation of the renin-angiotensin-aldosterone system:

In patients whose vascular tone and renal function depend predominantly on the activity of the renin-angiotensin-aldosterone system (e.g. patients with severe congestive heart failure or underlying renal disease, including renal artery stenosis), treatment with other medicinal products that affect this system, such as angiotensin II receptor antagonists, has been associated with acute hypotension, azotaemia, oliguria or, rarely, acute renal failure.

#### Renovascular hypertension:

There is an increased risk of severe hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with medicinal products that affect the renin-angiotensin-aldosterone system.

Renal impairment and kidney transplantation:

When Olmesartan /Amlodipine Mylan is used in patients with impaired renal function, periodic monitoring of serum potassium and creatinine levels is recommended. Use of Olmesartan /Amlodipine Mylan is not recommended in patients with severe renal impairment (creatinine clearance <20 mL/min) (see sections 4.2, and 5.2). There is no experience of the administration of olmesartan medoxomil/amlodipine in patients with a recent kidney transplant or in patients with end-stage renal impairment (i.e. creatinine clearance < 12 mL/min).

Dual blockade of the renin-angiotensin-aldosterone system (RAAS):

There is evidence that the concomitant use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren increases the risk of hypotension, hyperkalaemia and decreased renal function (including acute renal failure). Dual blockade of RAAS through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is therefore not recommended (see sections 4.5 and 5.1).

If dual blockade therapy is considered absolutely necessary, this should only occur under specialist supervision and subject to frequent close monitoring of renal function, electrolytes and blood pressure.

ACE-inhibitors and angiotensin II receptor blockers should not be used concomitantly in patients with diabetic nephropathy.

Hepatic impairment:

Exposure to amlodipine and olmesartan medoxomil is increased in patients with hepatic impairment (see section 5.2). Care should be taken when Olmesartan /Amlodipine Mylan is administered in patients with mild to moderate hepatic impairment. In moderately impaired patients, the dose of olmesartan medoxomil should not exceed 20 mg (see section 4.2). In patients with impaired hepatic function, amlodipine should be initiated at the lower end of the dosing range and caution should be used, both on initial treatment and when increasing the dose. Use of Olmesartan /Amlodipine Mylan in patients with severe hepatic impairment is contraindicated (see section 4.3).

Hyperkalaemia:

As with other angiotensin II antagonists and ACE inhibitors, hyperkalaemia may occur during treatment, especially in the presence of renal impairment and/or heart failure (see section 4.5). Close monitoring of serum potassium levels in at-risk patients is recommended.

Concomitant use with potassium supplements, potassium-sparing diuretics, salt substitutes containing potassium, or other medicinal products that may increase potassium levels (heparin, etc.) should be undertaken with caution and with frequent monitoring of potassium levels.

Lithium:

As with other angiotensin II receptor antagonists, the concomitant use of Olmesartan /Amlodipine Mylan and lithium is not recommended (see section 4.5).

Aortic or mitral valve stenosis; obstructive hypertrophic cardiomyopathy:

Due to the amlodipine component of Olmesartan /Amlodipine Mylan, as with all other vasodilators, special caution is indicated in patients suffering from aortic or mitral valve stenosis, or obstructive hypertrophic cardiomyopathy.

Primary aldosteronism:

Patients with primary aldosteronism generally will not respond to antihypertensive medicinal products acting through inhibition of the renin-angiotensin system. Therefore, the use of Olmesartan /Amlodipine Mylan is not recommended in such patients.

Heart failure:

As a consequence of the inhibition of the renin-angiotensin-aldosterone system, changes in renal function may be anticipated in susceptible individuals. In patients with severe heart failure whose renal function may depend on the activity of the renin-angiotensin-aldosterone system, treatment with angiotensin-converting enzyme (ACE) inhibitors and angiotensin receptor antagonists has been associated with oliguria and/or progressive azotaemia and (rarely) with acute renal failure and/or death.

Patients with heart failure should be treated with caution. In a long-term, placebo controlled study of amlodipine in patients with severe heart failure (NYHA III and IV), the reported incidence of pulmonary oedema was higher in the amlodipine group than in the placebo group (see section 5.1). Calcium channel blockers, including amlodipine, should be used with caution in patients with congestive heart failure, as they may increase the risk of future cardiovascular events and mortality.

Sprue-like enteropathy:

In very rare cases severe, chronic diarrhoea with substantial weight loss has been reported in patients taking olmesartan few months to years after drug initiation, possibly caused by a localized delayed hypersensitivity reaction. Intestinal biopsies of patients often demonstrated villous atrophy. If a patient develops these symptoms during treatment with olmesartan, and in the absence of other apparent etiologies, olmesartan treatment should be immediately discontinued and should not be restarted. If diarrhoea does not improve during the week after the discontinuation, further specialist (e.g. a gastro-enterologist) advice should be considered.

Ethnic differences:

As with all other angiotensin II antagonists, the blood pressure lowering effect of Olmesartan /Amlodipine Mylan can be somewhat less in black patients than in non-black patients, possibly because of a higher prevalence of low-renin status in the black hypertensive population.

Elderly:

In the elderly, increase of the dosage should take place with care (see section 5.2).

Pregnancy:

Angiotensin II antagonists should not be initiated during pregnancy. Unless continued angiotensin II antagonist therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with angiotensin II antagonists should be stopped immediately, and, if appropriate, alternative therapy should be started (see sections 4.3 and 4.6).

Other:

As with any antihypertensive agent, excessive blood pressure decrease in patients with ischaemic heart disease or ischaemic cerebrovascular disease could result in a myocardial infarction or stroke.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

#### **4.5 Interaction with other medicinal products and other forms of interactions**

Potential interactions related to the olmesartan medoxomil/amlodipine combination:

*To be taken into account with concomitant use*

Other antihypertensive agents:

The blood pressure lowering effect of Olmesartan /Amlodipine Mylan can be increased by concomitant use of other antihypertensive medicinal products (e.g. alpha blockers, diuretics).

Potential interactions related to the olmesartan medoxomil component of Olmesartan /Amlodipine Mylan:

Concomitant use not recommended

ACE-inhibitors, angiotensin II receptor blockers or aliskiren:

Clinical trial data has shown that dual blockade of the renin-angiotensin-aldosterone-system (RAAS) through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is associated with a higher frequency of adverse events such as hypotension, hyperkalaemia and decreased renal function (including acute renal failure) compared to the use of a single RAAS-acting agent (see sections 4.3, 4.4 and 5.1).

Medicinal products affecting potassium levels:

Concomitant use of potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium or other medicinal products that may increase serum potassium levels (e.g. heparin, ACE inhibitors) may lead to increases in serum potassium (see section 4.4). If medicinal products which affect potassium levels are to be prescribed in combination with Olmesartan /Amlodipine Mylan, monitoring of serum potassium levels is recommended.

Lithium:

Reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with angiotensin converting enzyme inhibitors and, rarely, with angiotensin II antagonists. Therefore concomitant use of Olmesartan /Amlodipine Mylan and lithium is not recommended (see section 4.4). If concomitant use of Olmesartan /Amlodipine Mylan and lithium proves necessary, careful monitoring of serum lithium levels is recommended.

*Concomitant use requiring caution*Non-steroidal anti-inflammatory medicinal products (NSAIDs) including selective COX-2 inhibitors, acetylsalicylic acid (> 3 g/day) and non-selective NSAIDs:

When angiotensin II antagonists are administered simultaneously with NSAIDs, attenuation of the antihypertensive effect may occur. Furthermore, concomitant use of angiotensin II antagonists and NSAIDs may increase the risk of worsening of renal function and may lead to an increase in serum potassium. Therefore monitoring of renal function at the beginning of such concomitant therapy is recommended, as well as adequate hydration of the patient.

Bile acid sequestering agent colesevelam:

Concurrent administration of the bile acid sequestering agent colesevelam hydrochloride reduces the systemic exposure and peak plasma concentration of olmesartan and reduces t<sub>1/2</sub>. Administration of olmesartan medoxomil at least 4 hours prior to colesevelam hydrochloride decreased the drug interaction effect. Administering olmesartan medoxomil at least 4 hours before the colesevelam hydrochloride dose should be considered (see section 5.2).

Additional information

After treatment with antacid (aluminium magnesium hydroxide), a modest reduction in bioavailability of olmesartan was observed.

Olmesartan medoxomil had no significant effect on the pharmacokinetics or pharmacodynamics of warfarin or the pharmacokinetics of digoxin. Coadministration of olmesartan medoxomil with pravastatin had no clinically relevant effects on the pharmacokinetics of either component in healthy subjects.

Olmesartan had no clinically relevant inhibitory effects on human cytochrome P450 enzymes 1A1/2, 2A6, 2C8/9, 2C19, 2D6, 2E1 and 3A4 *in vitro*, and had no or minimal inducing effects on rat cytochrome P450 activities. No clinically relevant interactions between olmesartan and medicinal products metabolised by the above cytochrome P450 enzymes are expected.

Potential interactions related to the amlodipine component of Olmesartan /Amlodipine Mylan:*Effects of other medicinal products on amlodipine*CYP3A4 inhibitors:

Concomitant use of amlodipine with strong or moderate CYP3A4 inhibitors (protease inhibitors, azole antifungals, macrolides like erythromycin or clarithromycin, verapamil or diltiazem) may give rise to significant increase in amlodipine exposure. The clinical translation of these PK variations may be more pronounced in the elderly. There is an increased risk of hypotension. Close observation of patients is recommended and dose adjustment may thus be required.

CYP3A4 inducers:

Upon co-administration of known inducers of the CYP3A4, the plasma concentration of amlodipine may vary. Therefore, blood pressure should be monitored and dose regulation considered both during and after concomitant medication particularly with strong CYP3A4 inducers (e.g. rifampicin, *Hypericum perforatum*).

Administration of amlodipine with grapefruit or grapefruit juice is not recommended as bioavailability may be increased in some patients resulting in increased blood pressure lowering effects.

Dantrolene (infusion): In animals, lethal ventricular fibrillation and cardiovascular collapse are observed in association with hyperkalaemia after administration of verapamil and intravenous dantrolene. Due to risk of hyperkalaemia, it is recommended that the co-administration of calcium channel blockers such as amlodipine be avoided in patients susceptible to malignant hyperthermia and in the management of malignant hyperthermia.

*Effects of amlodipine on other medicinal products*

The blood pressure lowering effects of amlodipine adds to the blood pressure-lowering effects of other antihypertensive agents.

In clinical interaction studies, amlodipine did not affect the pharmacokinetics of atorvastatin, digoxin or warfarin.

#### Simvastatin:

Co-administration of multiple doses of 10 mg of amlodipine with 80 mg simvastatin resulted in a 77% increase in exposure to simvastatin compared to simvastatin alone. Limit the dose of simvastatin in patients on amlodipine to 20 mg daily.

#### Tacrolimus:

There is a risk of increased tacrolimus blood levels when co-administered with amlodipine but the pharmacokinetic mechanism of this interaction is not fully understood. In order to avoid toxicity of tacrolimus, administration of amlodipine in a patient treated with tacrolimus requires monitoring of tacrolimus blood levels and dose adjustment of tacrolimus when appropriate.

#### Cyclosporine:

In a prospective study in renal transplant patients, an average 40% increase in trough cyclosporine levels was observed when used concomitantly with amlodipine. The co-administration of Olmesartan /Amlodipine Mylan with cyclosporine may increase exposure to cyclosporine. Monitor trough cyclosporine levels during concomitant use and cyclosporine dose reductions should be made as necessary.

### **4.6 Fertility, pregnancy and lactation**

#### Pregnancy (see section 4.3)

There are no data about the use of olmesartan medoxomil/amlodipine in pregnant patients. Animal reproductive toxicity studies with olmesartan medoxomil/amlodipine have not been performed.

#### Olmesartan medoxomil

The use of angiotensin II antagonists is not recommended during the first trimester of pregnancy (see section 4.4). The use of angiotensin II antagonists is contraindicated during the 2<sup>nd</sup> and 3<sup>rd</sup> trimesters of pregnancy (see sections 4.3 and 4.4).

Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Whilst there is no controlled epidemiological data on the risk with angiotensin II antagonists, similar risks may exist for this class of drugs. Unless continued angiotensin II antagonists therapy is considered essential, patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with angiotensin II antagonists should be stopped immediately, and, if appropriate, alternative therapy should be started.

Exposure to angiotensin II antagonists therapy during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia) (see section 5.3).

Should exposure to angiotensin II antagonists have occurred from the second trimester on, ultrasound check of renal function and skull is recommended. Infants whose mothers have taken angiotensin II antagonists should be closely observed for hypotension (see sections 4.3 and 4.4).

#### Amlodipine

Data on a limited number of exposed pregnancies do not indicate that amlodipine or other calcium receptor antagonists have a harmful effect on the health of the fetus. However, there may be a risk of prolonged delivery.

As a consequence, Olmesartan /Amlodipine Mylan is not recommended during the first trimester of pregnancy and is contraindicated during the second and third trimesters of pregnancy (see sections 4.3 and 4.4).

#### Breast-feeding

Olmesartan is excreted into the milk of lactating rats. However, it is not known whether olmesartan passes into human milk. Amlodipine is excreted in human milk. The proportion of the maternal dose received by the infant has been estimated with an interquartile range of 3-7 %, with a maximum of 15%. The effect of amlodipine on infants is unknown.

During breast-feeding, Olmesartan /Amlodipine Mylan is not recommended and alternative treatments with better established safety profiles during breast-feeding are preferable, especially while nursing a newborn or preterm infant.

#### Fertility

Reversible biochemical changes in the head of spermatozoa have been reported in some patients treated by calcium channel blockers. Clinical data are insufficient regarding the potential effect of amlodipine on fertility. In one rat study, adverse effects were found on male fertility (see section 5.3).

#### 4.7 Effects on ability to drive and use machines

Olmesartan /Amlodipine Mylan can have minor or moderate influence on the ability to drive and use machines.

Dizziness, headache, nausea or fatigue may occasionally occur in patients taking antihypertensive therapy, which may impair the ability to react. Caution is recommended especially at the start of treatment.

#### 4.8 Undesirable effects

The most commonly reported adverse reactions during treatment with olmesartan medoxomil/amlodipine are peripheral oedema (11.3%), headache (5.3%) and dizziness (4.5%).

Adverse reactions from olmesartan medoxomil/amlodipine in clinical trials, post-authorisation safety studies and spontaneous reporting are summarised in the below table as well as adverse reactions from the individual components olmesartan medoxomil and amlodipine based on the known safety profile of these substances.

The following terminologies have been used in order to classify the occurrence of adverse reactions, as applicable:

Very common ( $\geq 1/10$ )

Common ( $\geq 1/100$  to  $< 1/10$ )

Uncommon ( $\geq 1/1,000$  to  $< 1/100$ )

Rare ( $\geq 1/10,000$  to  $< 1/1,000$ )

Very rare ( $< 1/10,000$ )

Not known (cannot be estimated from the available data)

| MedDRA System Organ Class            | Adverse reactions                        | Frequency                         |            |                           |
|--------------------------------------|--|-----------------------------------|------------|---------------------------|
|                                      |  | Olmesartan/Amlodipine combination | Olmesartan | Amlodipine                |
| Blood and lymphatic system disorders | Leukocytopenia                           |                                   |            | Very rare                 |
|                                      | Thrombocytopenia                         |                                   | Uncommon   | Very rare                 |
| Immune system disorders              | Allergic reaction /Drug hypersensitivity | Rare                              |            | Very rare                 |
|                                      | Anaphylactic reaction                    |                                   | Uncommon   |                           |
| Metabolism and nutrition disorders   | Hyperglycaemia                           |                                   |            | Very rare                 |
|                                      | Hyperkalaemia                            | Uncommon                          | Rare       |                           |
|                                      | Hypertriglyceridaemia                    |                                   | Common     |                           |
|                                      | Hyperuricaemia                           |                                   | Common     |                           |
| Psychiatric disorders                | Confusion                                |                                   |            | Rare                      |
|                                      | Depression                               |                                   |            | Uncommon                  |
|                                      | Insomnia                                 |                                   |            | Uncommon                  |
|                                      | Irritability                             |                                   |            | Uncommon                  |
|                                      | Libido decreased                         | Uncommon                          |            |                           |
|                                      | Mood changes (including anxiety)         |                                   |            | Uncommon                  |
| Nervous system disorders             | Dizziness                                | Common                            | Common     | Common                    |
|                                      | Dysgeusia                                |                                   |            | Uncommon                  |
|                                      | Headache                                 | Common                            | Common     | Common (especially at the |

|  |   |          |          |   |
|--|---|----------|----------|---|
|  |   |          |          | beginning of treatment)                         |
|  | Hypertonia  |          |          | Very rare                                       |
|  | Hypoaesthesia   | Uncommon |          | Uncommon  |
|  | Lethargy  | Uncommon |          |   |
|  | Paraesthesia  | Uncommon |          | Uncommon  |
|  | Peripheral neuropathy   |          |          | Very rare                                       |
|  | Postural dizziness  | Uncommon |          |   |
|  | Sleep disorder  |          |          | Uncommon  |
|  | Somnolence  |          |          | Common  |
|  | Syncope   | Rare     |          | Uncommon  |
|  | Tremor  |          |          | Uncommon  |
|  | Extrapyramidal disorder   |          |          | Not known                                       |
| <b>Eye disorders</b>                                   | Visual disturbance (including diplopia)   |          |          | Common  |
| <b>Ear and labyrinth disorders</b>                     | Tinnitus  |          |          | Uncommon  |
|  | Vertigo   | Uncommon | Uncommon |   |
| <b>Cardiac disorders</b>                               | Angina pectoris   |          | Uncommon | Uncommon (incl. aggravation of angina pectoris) |
|  | Arrhythmia (including bradycardia, ventricular tachycardia and atrial fibrillation) |          |          | Uncommon  |
|  | Myocardial infarction   |          |          | Very rare                                       |
|  | Palpitations  | Uncommon |          | Common  |
|  | Tachycardia   | Uncommon |          |   |
| <b>Vascular disorders</b>                              | Hypotension   | Uncommon | Rare     | Uncommon  |
|  | Orthostatic hypotension   | Uncommon |          |   |
|  | Flushing  | Rare     |          | Common  |
|  | Vasculitis  |          |          | Very rare                                       |
| <b>Respiratory, thoracic and mediastinal disorders</b> | Bronchitis  |          | Common   |   |
|  | Cough   | Uncommon | Common   | Uncommon  |
|  | Dyspnoea  | Uncommon |          | Common  |
|  | Pharyngitis   |          | Common   |   |
|  | Rhinitis  |          | Common   | Uncommon  |

| MedDRA System Organ Class         | Adverse reactions   | Frequency                         |            |            |
|-----------------------------------|---|-----------------------------------|------------|------------|
|                                   |   | Olmesartan/Amlodipine combination | Olmesartan | Amlodipine |
| <b>Gastrointestinal disorders</b> | Abdominal pain  |                                   | Common     | Common     |
|                                   | Altered bowel habits (including diarrhoea and constipation) |                                   |            | Common     |
|                                   | Constipation  | Uncommon                          |            |            |
|                                   | Diarrhoea   | Uncommon                          | Common     |            |
|                                   | Dry mouth   | Uncommon                          |            | Uncommon   |
|                                   | Dyspepsia   | Uncommon                          | Common     | Common     |
|                                   | Gastritis   |                                   |            | Very rare  |
|                                   | Gastroenteritis   |                                   | Common     |            |
|                                   | Gingival hyperplasia  |                                   |            | Very rare  |
|                                   | Nausea  | Uncommon                          | Common     | Common     |
|                                   | Pancreatitis  |                                   |            | Very rare  |
|                                   | Upper abdominal pain  | Uncommon                          |            |            |
|                                   | Vomiting  | Uncommon                          | Uncommon   | Uncommon   |



|                                |  |  |           |  |
|--------------------------------|--|--|-----------|--|
|                                | Sprue-like enteropathy (see section 4.4) |  | Very rare |  |
| <b>Hepatobiliary disorders</b> | Hepatic enzymes increased                |  | Common    | Very rare (mostly consistent with cholestasis) |
|                                | Hepatitis                                |  |           | Very rare                                      |
|                                | Jaundice                                 |  |           | Very rare                                      |

| <b>MedDRA System Organ Class</b>              | <b>Adverse reactions</b>   | <b>Frequency</b>                         |                   |                   |
|---|----------------------------|--|-------------------|-------------------|
|   |                            | <b>Olmesartan/Amlodipine combination</b> | <b>Olmesartan</b> | <b>Amlodipine</b> |
| <b>Skin and subcutaneous tissue disorders</b> | Alopecia                   |  |                   | Uncommon          |
|   | Angioneurotic oedema       |  | Rare              | Very rare         |
|   | Allergic dermatitis        |  | Uncommon          |                   |
|   | Erythema multiforme        |  |                   | Very rare         |
|   | Exanthema                  |  | Uncommon          | Uncommon          |
|   | Exfoliative dermatitis     |  |                   | Very rare         |
|   | Hyperhidrosis              |  |                   | Uncommon          |
|   | Photosensitivity           |  |                   | Very rare         |
|   | Pruritus                   |  | Uncommon          | Uncommon          |
|   | Purpura                    |  |                   | Uncommon          |
|   | Quincke oedema             |  |                   | Very rare         |
|   | Rash                       | Uncommon                                 | Uncommon          | Uncommon          |
|   | Skin discoloration         |  |                   | Uncommon          |
|   | Stevens-Johnson syndrome   |  |                   | Very rare         |
|   | Toxic epidermal necrolysis |  |                   | Not known         |
| Urticaria                                     | Rare                       | Uncommon                                 | Uncommon          |                   |

| <b>MedDRA System Organ Class</b>                       | <b>Adverse reactions</b>       | <b>Frequency</b>                         |                   |                   |
|--|--------------------------------|--|-------------------|-------------------|
|  |                                | <b>Olmesartan/Amlodipine combination</b> | <b>Olmesartan</b> | <b>Amlodipine</b> |
| <b>Musculoskeletal and connective tissue disorders</b> | Ankle swelling                 |  |                   | Common            |
|  | Arthralgia                     |  |                   | Uncommon          |
|  | Arthritis                      |  | Common            |                   |
|  | Back pain                      | Uncommon                                 | Common            | Uncommon          |
|  | Muscle spasm                   | Uncommon                                 | Rare              | Common            |
|  | Myalgia                        |  | Uncommon          | Uncommon          |
|  | Pain in extremity              | Uncommon                                 |                   |                   |
|  | Skeletal pain                  |  | Common            |                   |
| <b>Renal and urinary disorders</b>                     | Acute renal failure            |  | Rare              |                   |
|  | Haematuria                     |  | Common            |                   |
|  | Increased urinary frequency    |  |                   | Uncommon          |
|  | Micturition disorder           |  |                   | Uncommon          |
|  | Nocturia                       |  |                   | Uncommon          |
|  | Pollakiuria                    | Uncommon                                 |                   |                   |
|  | Renal insufficiency            |  | Rare              |                   |
|  | Urinary tract infection        |  | Common            |                   |
| <b>Reproductive system and breast disorders</b>        | Erectile dysfunction/impotence | Uncommon                                 |                   | Uncommon          |
|  | Gynecomastia                   |  |                   | Uncommon          |

| MedDRA<br>System Organ Class                            | Adverse<br>reactions                          | Frequency                            |            |             |
|---|---|--------------------------------------|------------|-------------|
|   |   | Olmesartan/Amlodipine<br>combination | Olmesartan | Amlodipine  |
| General disorders and administration site<br>conditions | Asthenia                                      | Uncommon                             | Uncommon   | Common      |
|   | Chest pain                                    |                                      | Common     | Uncommon    |
|   | Face oedema                                   | Rare                                 | Uncommon   |             |
|   | Fatigue                                       | Common                               | Common     | Common      |
|   | Influenza-like<br>symptoms                    |                                      | Common     |             |
|   | Lethargy                                      |                                      | Rare       |             |
|   | Malaise                                       |                                      | Uncommon   | Uncommon    |
|   | Oedema  | Common                               |            | Very common |
|   | Pain  |                                      | Common     | Uncommon    |
|   | Peripheral<br>oedema                          | Common                               | Common     |             |
|   | Pitting oedema                                | Common                               |            |             |
| Investigations  | Blood<br>creatinine<br>increased              | Uncommon                             | Rare       |             |
|   | Blood creatine<br>phosphokinase<br>increased  |                                      | Common     |             |
|   | Blood<br>potassium<br>decreased               | Uncommon                             |            |             |
|   | Blood urea<br>increased                       |                                      | Common     |             |
|   | Blood uric acid<br>increased                  | Uncommon                             |            |             |
|   | Gamma<br>glutamyl<br>transferase<br>increased | Uncommon                             |            |             |
|   | Weight<br>decrease                            |                                      |            | Uncommon    |
|   | Weight<br>increase                            |                                      |            | Uncommon    |

Single cases of rhabdomyolysis have been reported in temporal association with the intake of angiotensin II receptor blockers. Single cases of extrapyramidal syndrome have been reported in patients treated with amlodipine.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRC Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: [www.hpra.ie](http://www.hpra.ie); e-mail: [medsafety@hpra.ie](mailto:medsafety@hpra.ie).

#### **4.9 Overdose**

##### Symptoms

There is no experience of overdose with olmesartan medoxomil/amlodipine. The most likely effects of olmesartan medoxomil overdosage are hypotension and tachycardia; bradycardia could be encountered if parasympathetic (vagal) stimulation occurred. Amlodipine overdosage can be expected to lead to excessive peripheral vasodilatation with marked hypotension and possibly a reflex tachycardia. Marked and potentially prolonged systemic hypotension up to and including shock with fatal outcome has been reported.

##### Management

If intake is recent, gastric lavage may be considered. In healthy subjects, the administration of activated charcoal immediately or up to 2 hours after ingestion of amlodipine has been shown to reduce substantially the absorption of amlodipine.

Clinically significant hypotension due to an overdose of olmesartan medoxomil/amlodipine requires active support of the cardiovascular system, including close monitoring of heart and lung function, elevation of the extremities, and attention to circulating fluid volume and urine output. A vasoconstrictor may be helpful in restoring vascular tone and blood pressure, provided that there is no contraindication to its use. Intravenous calcium gluconate may be beneficial in reversing the effects of calcium channel blockade.

Since amlodipine is highly protein-bound, dialysis is not likely to be of benefit. The dialysability of olmesartan is unknown.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Angiotensin II antagonists and calcium channel blockers, ATC code C09DB02.

#### Mechanism of action

Olmesartan /Amlodipine Mylan is a combination of an angiotensin II receptor antagonist, olmesartan medoxomil, and a calcium channel blocker, amlodipine besilate. The combination of these active ingredients has an additive antihypertensive effect, reducing blood pressure to a greater degree than either component alone.

#### Clinical efficacy and safety

##### *Olmesartan medoxomil/amlodipine*

In an 8-week, double-blind, randomised, placebo-controlled factorial design study in 1940 patients (71% Caucasian and 29% non-Caucasian patients), treatment with each combination dose of olmesartan medoxomil/amlodipine resulted in significantly greater reductions in diastolic and systolic blood pressures than the respective monotherapy components. The mean change in systolic/diastolic blood pressure was dose-dependent: -24/-14 mmHg (20 mg/5 mg combination), -25/-16 mmHg (40 mg/5 mg combination) and -30/-19 mmHg (40 mg/10 mg combination).

Olmesartan medoxomil/amlodipine 40 mg/5 mg reduced seated systolic/diastolic blood pressure by an additional 2.5/1.7 mmHg over olmesartan medoxomil/amlodipine 20 mg/5 mg. Similarly olmesartan medoxomil/amlodipine 40 mg/10 mg reduced seated systolic/diastolic blood pressure by an additional 4.7/3.5 mmHg over olmesartan medoxomil/amlodipine 40 mg/5 mg.

The proportions of patients reaching blood pressure goal (< 140/90 mmHg for non-diabetic patients and < 130/80 mmHg for diabetic patients) were 42.5%, 51.0% and 49.1% for olmesartan medoxomil/amlodipine 20 mg/5 mg, 40 mg/5 mg and 40 mg/10 mg respectively.

The majority of the antihypertensive effect of olmesartan medoxomil/amlodipine was generally achieved within the first 2 weeks of therapy.

A second double-blind, randomised, placebo-controlled study evaluated the effectiveness of adding amlodipine to the treatment in Caucasian patients whose blood pressure was inadequately controlled by 8 weeks of monotherapy with 20 mg olmesartan medoxomil.

In patients who continued to receive only 20 mg olmesartan medoxomil, systolic/diastolic blood pressure was reduced by -10.6/-7.8 mmHg after a further 8 weeks. The addition of 5 mg amlodipine for 8 weeks resulted in a reduction in systolic/diastolic blood pressure of -16.2/-10.6 mmHg ( $p = 0.0006$ ).

The proportion of patients reaching blood pressure goal (< 140/90 mmHg for non-diabetic patients and < 130/80 mmHg for diabetic patients) was 44.5% for the 20 mg/5 mg combination compared to 28.5% for 20 mg olmesartan medoxomil.

A further study evaluated the addition of various doses of olmesartan medoxomil in Caucasian patients whose blood pressure was not adequately controlled by 8 weeks of monotherapy with 5 mg amlodipine.

In patients who continued to receive only 5 mg amlodipine, systolic/diastolic blood pressure was reduced by -9.9/-5.7 mmHg after a further 8 weeks. The addition of 20 mg olmesartan medoxomil resulted in a reduction in systolic/diastolic blood

pressure of -15.3/-9.3 mmHg and the addition of 40 mg olmesartan medoxomil resulted in a reduction in systolic/diastolic blood pressure of -16.7/-9.5 mmHg ( $p < 0.0001$ ).

The proportions of patients reaching blood pressure goal ( $< 140/90$  mmHg for non-diabetic patients and  $< 130/80$  mmHg for diabetic patients) was 29.9% for the group who continued to receive 5 mg amlodipine alone, 53.5% for olmesartan medoxomil/amlodipine 20 mg/5 mg and 50.5% for olmesartan medoxomil/amlodipine 40 mg/5 mg.

Randomised data in uncontrolled hypertensive patients, comparing the use of medium dose olmesartan medoxomil/amlodipine combination therapy versus escalation to top dose monotherapy of amlodipine or olmesartan, are not available.

The three studies performed confirmed that the blood pressure lowering effect of olmesartan medoxomil/amlodipine once daily was maintained throughout the 24-hour dose interval, with trough-to-peak ratios of 71% to 82% for systolic and diastolic response and with 24-hour effectiveness being confirmed by ambulatory blood pressure monitoring.

The antihypertensive effect of olmesartan medoxomil/amlodipine was similar irrespective of age and gender, and was similar in patients with and without diabetes.

In two open-label, non-randomised extension studies, sustained efficacy using olmesartan medoxomil/amlodipine 40 mg/5 mg was demonstrated at one year for 49 - 67% of patients.

### *Olmesartan medoxomil*

The olmesartan medoxomil component of Olmesartan /Amlodipine Mylan is a selective angiotensin II type 1 ( $AT_1$ ) receptor antagonist. Olmesartan medoxomil is rapidly converted to the pharmacologically active metabolite, olmesartan. Angiotensin II is the primary vasoactive hormone of the renin-angiotensin-aldosterone system and plays a significant role in the pathophysiology of hypertension. The effects of angiotensin II include vasoconstriction, stimulation of the synthesis and release of aldosterone, cardiac stimulation and renal reabsorption of sodium. Olmesartan blocks the vasoconstrictor and aldosterone-secreting effects of angiotensin II by blocking its binding to the  $AT_1$  receptor in tissues including vascular smooth muscle and the adrenal gland. The action of olmesartan is independent of the source or route of synthesis of angiotensin II. The selective antagonism of the angiotensin II ( $AT_1$ ) receptors by olmesartan results in increases in plasma renin levels and angiotensin I and II concentrations, and some decrease in plasma aldosterone concentrations.

In hypertension, olmesartan medoxomil causes a dose-dependent, long-lasting reduction in arterial blood pressure. There has been no evidence of first-dose hypotension, of tachyphylaxis during long-term treatment, or of rebound hypertension after abrupt cessation of therapy.

Following once daily administration to patients with hypertension, olmesartan medoxomil produces an effective and smooth reduction in blood pressure over the 24 hour dose interval. Once daily dosing produced similar decreases in blood pressure as twice daily dosing at the same total daily dose.

With continuous treatment, maximum reductions in blood pressure are achieved by 8 weeks after the initiation of therapy, although a substantial proportion of the blood pressure lowering effect is already observed after 2 weeks of treatment.

The effect of olmesartan medoxomil on mortality and morbidity is not yet known.

The Randomised Olmesartan and Diabetes Microalbuminuria Prevention (ROADMAP) study in 4447 patients with type 2 diabetes, normo-albuminuria and at least one additional cardiovascular risk factor, investigated whether treatment with olmesartan could delay the onset of microalbuminuria. During the median follow-up duration of 3.2 years, patients received either olmesartan or placebo in addition to other antihypertensive agents, except ACE inhibitors or ARBs.

For the primary endpoint, the study demonstrated a significant risk reduction in the time to onset of microalbuminuria, in favour of olmesartan. After adjustment for BP differences this risk reduction was no longer statistically significant. 8.2% (178 of 2160) of the patients in the olmesartan group and 9.8% (210 of 2139) in the placebo group developed microalbuminuria.

For the secondary endpoints, cardiovascular events occurred in 96 patients (4.3%) with olmesartan and in 94 patients (4.2%) with placebo. The incidence of cardiovascular mortality was higher with olmesartan compared to placebo treatment (15 patients (0.7%) vs. 3 patients (0.1%)), despite similar rates for non-fatal stroke (14 patients (0.6%) vs. 8 patients (0.4%)), non-fatal myocardial infarction (17 patients (0.8%) vs. 26 patients (1.2%)) and non-cardiovascular mortality (11 patients (0.5%)).

vs. 12 patients (0.5%)). Overall mortality with olmesartan was numerically increased (26 patients (1.2%) vs. 15 patients (0.7%)), which was mainly driven by a higher number of fatal cardiovascular events.

The Olmesartan Reducing Incidence of End-stage Renal Disease in Diabetic Nephropathy Trial (ORIENT) investigated the effects of olmesartan on renal and cardiovascular outcomes in 577 randomized Japanese and Chinese type 2 diabetic patients with overt nephropathy. During a median follow-up of 3.1 years, patients received either olmesartan or placebo in addition to other antihypertensive agents including ACE inhibitors.

The primary composite endpoint (time to first event of the doubling of serum creatinine, end-stage renal disease, all-cause death) occurred in 116 patients in the olmesartan group (41.1%) and 129 patients in the placebo group (45.4%) (HR 0.97 (95% CI 0.75 to 1.24);  $p=0.791$ ). The composite secondary cardiovascular endpoint occurred in 40 olmesartan-treated patients (14.2%) and 53 placebo-treated patients (18.7%). This composite cardiovascular endpoint included cardiovascular death in 10 (3.5%) patients receiving olmesartan versus 3 (1.1%) receiving placebo, overall mortality 19 (6.7%) versus 20 (7.0%), non-fatal stroke 8 (2.8%) versus 11 (3.9%) and non-fatal myocardial infarction 3 (1.1%) versus 7 (2.5%), respectively.

### *Amlodipine*

The amlodipine component of Olmesartan /Amlodipine Mylan is a calcium channel blocker that inhibits the transmembrane influx of calcium ions through the potential-dependent L-type channels into the heart and smooth muscle. Experimental data indicate that amlodipine binds to both dihydropyridine and non-dihydropyridine binding sites. Amlodipine is relatively vessel-selective, with a greater effect on vascular smooth muscle cells than on cardiac muscle cells. The antihypertensive effect of amlodipine derives from a direct relaxant effect on arterial smooth muscle, which leads to a lowering of peripheral resistance and hence of blood pressure.

In hypertensive patients, amlodipine causes a dose-dependent, long-lasting reduction in arterial blood pressure. There has been no evidence of first-dose hypotension, of tachyphylaxis during long-term treatment, or of rebound hypertension after abrupt cessation of therapy.

Following administration of therapeutic doses to patients with hypertension, amlodipine produces an effective reduction in blood pressure in the supine, sitting and standing positions. Chronic use of amlodipine is not associated with significant changes in heart rate or plasma catecholamine levels. In hypertensive patients with normal renal function, therapeutic doses of amlodipine reduce renal vascular resistance and increase glomerular filtration rate and effective renal plasma flow, without changing filtration fraction or proteinuria.

In haemodynamic studies in patients with heart failure and in clinical studies based on exercise tests in patients with NYHA class II-IV heart failure, amlodipine was found not to cause any clinical deterioration, as measured by exercise tolerance, left ventricular ejection fraction and clinical signs and symptoms.

A placebo-controlled study (PRAISE) designed to evaluate patients with NYHA class III-IV heart failure receiving digoxin, diuretics and ACE inhibitors has shown that amlodipine did not lead to an increase in risk of mortality or combined mortality and morbidity in patients with heart failure.

In a follow-up, long-term, placebo controlled study (PRAISE-2) of amlodipine in patients with NYHA III and IV heart failure without clinical symptoms or objective findings suggestive of underlying ischaemic disease, on stable doses of ACE inhibitors, digitalis, and diuretics, amlodipine had no effect on total or cardiovascular mortality. In this same population amlodipine was associated with increased reports of pulmonary oedema despite no significant difference in the incidence of worsening heart failure as compared to placebo.

### Treatment to prevent heart attack trial (ALLHAT)

A randomized double-blind morbidity-mortality study called the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT) was performed to compare newer drug therapies: amlodipine 2.5-10 mg/d (calcium channel blocker) or lisinopril 10-40 mg/d (ACE-inhibitor) as first-line therapies to that of the thiazide-diuretic, chlorthalidone 12.5-25 mg/d in mild to moderate hypertension.

A total of 33,357 hypertensive patients aged 55 or older were randomized and followed for a mean of 4.9 years. The patients had at least one additional CHD risk factor, including: previous myocardial infarction or stroke (> 6 months prior to enrolment) or documentation of other atherosclerotic CVD (overall 51.5%), type 2 diabetes (36.1%), HDL-C < 35 mg/dL (11.6%), left ventricular hypertrophy diagnosed by electrocardiogram or echocardiography (20.9%), current cigarette smoking (21.9%).

The primary endpoint was a composite of fatal CHD or non-fatal myocardial infarction. There was no significant difference in the primary endpoint between amlodipine-based therapy and chlorthalidone-based therapy: RR 0.98 95% CI (0.90-1.07)  $p=0.65$ . Among secondary endpoints, the incidence of heart failure (component of a composite combined cardiovascular endpoint) was significantly higher in the amlodipine group as compared to the chlorthalidone group (10.2% vs. 7.7%, RR 1.38, 95% CI [1.25-1.52]  $p<0.001$ ). However, there was no significant difference in all-cause mortality between amlodipine-based therapy and chlorthalidone-based therapy (RR 0.96 95% CI [0.89-1.02]  $p=0.20$ ).

*Other information:*

Two large randomised, controlled trials (ONTARGET (ONgoing Telmisartan Alone and in combination with Ramipril Global Endpoint Trial) and VA NEPHRON-D (The Veterans Affairs Nephropathy in Diabetes)) have examined the use of the combination of an ACE-inhibitor with an angiotensin II receptor blocker.

ONTARGET was a study conducted in patients with a history of cardiovascular or cerebrovascular disease, or type 2 diabetes mellitus accompanied by evidence of end-organ damage. VA NEPHRON-D was a study in patients with type 2 diabetes mellitus and diabetic nephropathy.

These studies have shown no significant beneficial effect on renal and/or cardiovascular outcomes and mortality, while an increased risk of hyperkalaemia, acute kidney injury and/or hypotension as compared to monotherapy was observed. Given their similar pharmacodynamic properties, these results are also relevant for other ACE-inhibitors and angiotensin II receptor blockers.

ACE-inhibitors and angiotensin II receptor blockers should therefore not be used concomitantly in patients with diabetic nephropathy.

ALTITUDE (Aliskiren Trial in Type 2 Diabetes Using Cardiovascular and Renal Disease Endpoints) was a study designed to test the benefit of adding aliskiren to a standard therapy of an ACE-inhibitor or an angiotensin II receptor blocker in patients with type 2 diabetes mellitus and chronic kidney disease, cardiovascular disease, or both. The study was terminated early because of an increased risk of adverse outcomes. Cardiovascular death and stroke were both numerically more frequent in the aliskiren group than in the placebo group and adverse events and serious adverse events of interest (hyperkalaemia, hypotension and renal dysfunction) were more frequently reported in the aliskiren group than in the placebo group.

## 5.2 Pharmacokinetic properties

### *Olmesartan medoxomil/amlodipine*

Following oral intake of olmesartan medoxomil/amlodipine, peak plasma concentrations of olmesartan and amlodipine are reached at 1.5 – 2 h and 6 – 8 hours, respectively. The rate and extent of absorption of the two active substances from olmesartan medoxomil/amlodipine are equivalent to the rate and extent of absorption following intake of the two components as separate tablets. Food does not affect the bioavailability of olmesartan and amlodipine from Olmesartan /Amlodipine Mylan.

### *Olmesartan medoxomil*

#### Absorption and distribution

Olmesartan medoxomil is a prodrug. It is rapidly converted to the pharmacologically active metabolite, olmesartan, by esterases in the gut mucosa and in portal blood during absorption from the gastrointestinal tract. No intact olmesartan medoxomil or intact side chain medoxomil moiety have been detected in plasma or excreta. The mean absolute bioavailability of olmesartan from a tablet formulation was 25.6%.

The mean peak plasma concentration ( $C_{max}$ ) of olmesartan is reached within about 2 hours after oral dosing with olmesartan medoxomil, and olmesartan plasma concentrations increase approximately linearly with increasing single oral doses up to about 80 mg.

Food had minimal effect on the bioavailability of olmesartan and therefore olmesartan medoxomil may be administered with or without food.

No clinically relevant gender-related differences in the pharmacokinetics of olmesartan have been observed.

Olmesartan is highly bound to plasma protein (99.7%), but the potential for clinically significant protein binding displacement interactions between olmesartan and other highly bound coadministered active substances is low (as confirmed by the lack of

a clinically significant interaction between olmesartan medoxomil and warfarin). The binding of olmesartan to blood cells is negligible. The mean volume of distribution after intravenous dosing is low (16 – 29 L).

#### Biotransformation and elimination

Total plasma clearance of olmesartan was typically 1.3 L/h (CV, 19%) and was relatively slow compared to hepatic blood flow (ca 90 L/h). Following a single oral dose of <sup>14</sup>C-labelled olmesartan medoxomil, 10% – 16% of the administered radioactivity was excreted in the urine (the vast majority within 24 hours of dose administration) and the remainder of the recovered radioactivity was excreted in the faeces. Based on the systemic availability of 25.6%, it can be calculated that absorbed olmesartan is cleared by both renal excretion (ca 40%) and hepato-biliary excretion (ca 60%). All recovered radioactivity was identified as olmesartan. No other significant metabolite was detected. Enterohepatic recycling of olmesartan is minimal. Since a large proportion of olmesartan is excreted via the biliary route, use in patients with biliary obstruction is contraindicated (see section 4.3).

The terminal elimination half life of olmesartan is between 10 and 15 hours after multiple oral dosing. Steady state is reached after the first few doses and no further accumulation is evident after 14 days of repeated dosing. Renal clearance is approximately 0.5 – 0.7 L/h and is independent of dose.

#### Drug interactions

Bile acid sequestering agent colesevelam:

Concomitant administration of 40 mg olmesartan medoxomil and 3750 mg colesevelam hydrochloride in healthy subjects resulted in 28% reduction in  $C_{max}$  and 39% reduction in AUC of olmesartan. Lesser effects, 4% and 15% reduction in  $C_{max}$  and AUC respectively, were observed when olmesartan medoxomil was administered 4 hours prior to colesevelam hydrochloride. Elimination half life of olmesartan was reduced by 50 – 52% irrespectively of whether administered concomitantly or 4 hours prior to colesevelam hydrochloride (see section 4.5).

#### *Amlodipine*

#### Absorption and distribution

After oral administration of therapeutic doses, amlodipine is well absorbed with peak blood levels between 6-12 hours post dose. Absolute bioavailability has been estimated to be between 64 and 80%. The volume of distribution is approximately 21 l/kg. In vitro studies have shown that approximately 97.5% of circulating amlodipine is bound to plasma proteins. The absorption of amlodipine is unaffected by the concomitant intake of food.

#### Biotransformation and elimination

The terminal plasma elimination half life is about 35-50 hours and is consistent with once daily dosing. Amlodipine is extensively metabolised by the liver to inactive metabolites with 10% of the parent compound and 60% of metabolites excreted in the urine.

#### *Olmesartan medoxomil and amlodipine*

#### Special populations

##### Paediatric population (age below 18 years):

No pharmacokinetic data in paediatric patients are available.

##### Elderly (age 65 years or over):

In hypertensive patients, the olmesartan AUC at steady state is increased by ca 35% in elderly people (65 – 75 years old) and by ca 44% in very old people ( $\geq 75$  years old) compared with the younger age group (see section 4.2). This may be at least in part related to a mean decrease in renal function in this group of patients. The recommended dosage regimen for elderly people is, however, the same, although caution should be exercised when increasing the dosage.

The time to reach peak plasma concentrations of amlodipine is similar in older and younger subjects. Amlodipine clearance tends to be decreased with resulting increases in AUC and elimination half life in older people. Increases in AUC and elimination half life in patients with congestive heart failure were as expected for the patient age group in this study (see section 4.4).

##### Renal impairment:

In renally impaired patients, the olmesartan AUC at steady state increased by 62%, 82% and 179% in patients with mild, moderate and severe renal impairment, respectively, compared to healthy controls (see sections 4.2 and 4.4).

Amlodipine is extensively metabolised to inactive metabolites. Ten percent of the substance is excreted unchanged in the urine. Changes in amlodipine plasma concentration are not correlated with the degree of renal impairment. In these patients, amlodipine may be administered at the normal dosage. Amlodipine is not dialysable.

#### Hepatic impairment:

After single oral administration, olmesartan AUC values are 6% and 65% higher in mildly and moderately hepatically impaired patients, respectively, than in their corresponding matched healthy controls. The unbound fraction of olmesartan at 2 hours post-dose in healthy subjects, in patients with mild hepatic impairment and in patients with moderate hepatic impairment is 0.26%, 0.34% and 0.41%, respectively. Following repeated dosing in patients with moderate hepatic impairment, olmesartan mean AUC is again about 65% higher than in matched healthy controls. Olmesartan mean  $C_{max}$  values are similar in hepatically-impaired and healthy subjects. Olmesartan medoxomil has not been evaluated in patients with severe hepatic impairment (see sections 4.2 and 4.4).

Very limited clinical data are available regarding amlodipine administration in patients with hepatic impairment. The clearance of amlodipine is decreased and the half-life is prolonged in patients with impaired hepatic function, resulting in an increase in AUC of about 40% – 60% (see sections 4.2 and 4.4).

### **5.3 Preclinical safety data**

Based on the non-clinical toxicity profile of each substance, no exacerbation of toxicities for the combination is expected, because each substance has different targets, i.e. the kidneys for olmesartan medoxomil and the heart for amlodipine.

In a 3-month, repeat-dose toxicity study of orally administered olmesartan medoxomil/amlodipine in combination in rats the following alterations were observed: decreases in red blood cell count-related parameters and kidney changes both of which might be induced by the olmesartan medoxomil component; alterations in the intestines (luminal dilatation and diffuse mucosal thickening of the ileum and colon), the adrenals (hypertrophy of the glomerular cortical cells and vacuolation of the fascicular cortical cells), and hypertrophy of the ducts in the mammary glands which might be induced by the amlodipine component. These alterations neither augmented any of the previously reported and existing toxicity of the individual agents nor induced any new toxicity, and no toxicologically synergistic effects were observed.

#### *Olmesartan medoxomil*

In chronic toxicity studies in rats and dogs, olmesartan medoxomil showed similar effects to other AT<sub>1</sub> receptor antagonists and ACE inhibitors: raised blood urea (BUN) and creatinine; reduction in heart weight; reduction of red cell parameters (erythrocytes, haemoglobin, haematocrit); histological indications of renal damage (regenerative lesions of the renal epithelium, thickening of the basal membrane, dilatation of the tubules). These adverse effects caused by the pharmacological action of olmesartan medoxomil have also occurred in preclinical trials on other AT<sub>1</sub> receptor antagonists and ACE inhibitors and can be reduced by simultaneous oral administration of sodium chloride. In both species, increased plasma renin activity and hypertrophy/hyperplasia of the juxtaglomerular cells of the kidney were observed. These changes, which are a typical effect of the class of ACE inhibitors and other AT<sub>1</sub> receptor antagonists, would appear to have no clinical relevance.

Like other AT<sub>1</sub> receptor antagonists olmesartan medoxomil was found to increase the incidence of chromosome breaks in cell cultures in vitro. No relevant effects were observed in several in vivo studies using olmesartan medoxomil at very high oral doses of up to 2000 mg/kg. The overall data of a comprehensive genotoxicity testing program suggest that olmesartan is very unlikely to exert genotoxic effects under conditions of clinical use.

Olmesartan medoxomil was not carcinogenic, in a 2-year study in rats nor in two 6-month carcinogenicity studies in transgenic mice.

In reproductive studies in rats, olmesartan medoxomil did not affect fertility and there was no evidence of a teratogenic effect. In common with other angiotensin II antagonists, survival of offspring was reduced following exposure to olmesartan medoxomil and pelvic dilatation of the kidney was seen after exposure of the dams in late pregnancy and lactation. In common with other antihypertensive agents, olmesartan medoxomil was shown to be more toxic to pregnant rabbits than to pregnant rats, however, there was no indication of a fetotoxic effect.

#### *Amlodipine*

-

#### Reproductive toxicology



Reproductive studies in rats and mice have shown delayed date of delivery, prolonged duration of labour and decreased pup survival at dosages approximately 50 times greater than the maximum recommended dosage for humans based on mg/kg.

#### Impairment of fertility

There was no effect on the fertility of rats treated with amlodipine (males for 64 days and females 14 days prior to mating) at doses up to 10 mg/kg/day (8 times\* the maximum recommended human dose of 10 mg on a mg/m<sup>2</sup> basis). In another rat study in which male rats were treated with amlodipine besilate for 30 days at a dose comparable with the human dose based on mg/kg, decreased plasma follicle-stimulating hormone and testosterone were found as well as decreases in sperm density and in the number of mature spermatids and Sertoli cells.

#### Carcinogenesis, mutagenesis

Rats and mice treated with amlodipine in the diet for two years, at concentrations calculated to provide daily dosage levels of 0.5, 1.25, and 2.5 mg/kg/day showed no evidence of carcinogenicity. The highest dose (for mice, similar to, and for rats twice\* the maximum recommended clinical dose of 10 mg on a mg/m<sup>2</sup> basis) was close to the maximum tolerated dose for mice but not for rats.

Mutagenicity studies revealed no drug related effects at either the gene or chromosome levels.

\*Based on patient weight of 50 kg

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Microcrystalline cellulose  
Sodium laurilsulfate  
Citric acid monohydrate  
Hydroxypropylcellulose  
Croscarmellose sodium  
Maltose  
Mannitol  
Low-substituted hydroxypropylcellulose  
Magnesium stearate

#### Tablet coat

Polyvinyl alcohol  
Titanium dioxide (E171)  
Macrogol  
Talc

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

#### Blisters

3 years

#### Bottles

3 years

Once opened use within 100 days.

### **6.4 Special precautions for storage**

#### Blisters with desiccant

This medicinal product does not require any special storage conditions.

Blisters without desiccant

Do not store above 30°C.

Bottles

This medicinal product does not require any special storage conditions.

**6.5 Nature and contents of container**

Aluminium foil (laminated with desiccant) blister packs of 14, 28, 56 and 98 film-coated tablets.

Aluminium foil blister packs of 14, 28, 56 and 98 film-coated tablets.

HDPE bottle packs of 28, 56 and 98 film-coated tablets.

HDPE bottle (with integrated desiccant) packs of 28, 56 and 98 film-coated tablets.

Not all pack sizes may be marketed.

**6.6 Special precautions for disposal and other handling**

No special requirements for disposal.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

**7 MARKETING AUTHORISATION HOLDER**

McDermott Laboratories Ltd., T/A Gerard Laboratories

35/36 Baldoyle Industrial Estate

Grange Road

Dublin 13

Ireland

**8 MARKETING AUTHORISATION NUMBER**

PA0577/185/001

**9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 18<sup>th</sup> May 2018

**10 DATE OF REVISION OF THE TEXT**

July 2021