

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Pregabalin 25 mg capsules, hard

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each hard capsule contains 25 mg of pregabalin.

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Hard capsule

25 mg capsules: White cap / White body size '4' hard gelatin capsules imprinted with '138' on cap and 'J' on body with black ink, filled with white to off white powder.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Neuropathic pain

Pregabalin is indicated for the treatment of peripheral and central neuropathic pain in adults.

Epilepsy

Pregabalin is indicated as adjunctive therapy in adults with partial seizures with or without secondary generalisation.

Generalised Anxiety Disorder

Pregabalin is indicated for the treatment of Generalised Anxiety Disorder (GAD) in adults.

4.2 Posology and method of administration

Posology

The dose range is 150 to 600 mg per day given in either two or three divided doses.

Neuropathic pain

Pregabalin treatment can be started at a dose of 150 mg per day given as two or three divided doses. Based on individual patient response and tolerability, the dose may be increased to 300 mg per day after an interval of 3 to 7 days, and if needed, to a maximum dose of 600 mg per day after an additional 7-day interval.

Epilepsy

Pregabalin treatment can be started with a dose of 150 mg per day given as two or three divided doses. Based on individual patient response and tolerability, the dose may be increased to 300 mg per day after 1 week. The maximum dose of 600 mg per day may be achieved after an additional week.

Generalised Anxiety Disorder

The dose range is 150 to 600 mg per day given as two or three divided doses. The need for treatment should be reassessed regularly.

Pregabalin treatment can be started with a dose of 150 mg per day. Based on individual patient response and tolerability, the dose may be increased to 300 mg per day after 1 week. Following an additional week the dose may be increased to 450 mg per day. The maximum dose of 600 mg per day may be achieved after an additional week.

Discontinuation of pregabalin

In accordance with current clinical practice, if pregabalin has to be discontinued it is recommended this should be done gradually over a minimum of 1 week independent of the indication (see sections 4.4 and 4.8).

Renal impairment

Pregabalin is eliminated from the systemic circulation primarily by renal excretion as unchanged drug.

As pregabalin clearance is directly proportional to creatinine clearance (see section 5.2), dose reduction in patients with compromised renal function must be individualised according to creatinine clearance (CL_{cr}), as indicated in Table 1 determined using the following formula:

$$CL_{cr}(\text{ml/min}) = \left[\frac{1.23 \times [140 - \text{age (years)}] \times \text{weight (kg)}}{\text{serum creatinine } (\mu\text{mol/l})} \right] (\times 0.85 \text{ for female patients})$$

Pregabalin is removed effectively from plasma by haemodialysis (50% of drug in 4 hours). For patients receiving haemodialysis, the pregabalin daily dose should be adjusted based on renal function. In addition to the daily dose, a supplementary dose should be given immediately following every 4 hour haemodialysis treatment (see Table 1).

Table 1. Pregabalin dose adjustment based on renal function

| Creatinine clearance (CL _{cr}) (ml/min) | Total pregabalin daily dose * | | Dose regimen |
|--|-------------------------------|-----------------------|--------------------------|
| | Starting dose (mg/day) | Maximum dose (mg/day) | |
| ≥ 60 | 150 | 600 | BID or TID |
| ≥ 30 - < 60 | 75 | 300 | BID or TID |
| ≥ 15 - < 30 | 25 – 50 | 150 | Once Daily or BID |
| < 15 | 25 | 75 | Once Daily |
| Supplementary dosage following haemodialysis (mg) | | | |
| | 25 | 100 | Single dose [†] |

TID = Three divided doses

BID = Two divided doses

* Total daily dose (mg/day) should be divided as indicated by dose regimen to provide mg/dose

+ Supplementary dose is a single additional dose

Hepatic impairment

No dose adjustment is required for patients with hepatic impairment (see section 5.2).

Paediatric population

The safety and efficacy of pregabalin in children below the age of 12 years and in adolescents (12-17 years of age) have not been established. Currently available data are described in section 4.8, 5.1 and 5.2 but no recommendation on a posology can be made.

Elderly

Elderly patients may require a dose reduction of pregabalin due to a decreased renal function (see section 5.2)

Method of administration

This medicine may be taken with or without food.

This medicine is for oral use only.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Diabetic patients

In accordance with current clinical practice, some diabetic patients who gain weight on pregabalin treatment may need to adjust hypoglycaemic medicinal products.

Hypersensitivity reactions

There have been reports in the post marketing experience of hypersensitivity reactions, including cases of angioedema. Pregabalin should be discontinued immediately if symptoms of angioedema, such as facial, perioral, or upper airway swelling occur.

Dizziness, somnolence, loss of consciousness, confusion, and mental impairment

Pregabalin treatment has been associated with dizziness and somnolence, which could increase the occurrence of accidental injury (fall) in the elderly population. There have also been post marketing reports of loss of consciousness, confusion and mental impairment. Therefore, patients should be advised to exercise caution until they are familiar with the potential effects of the medicinal product.

Vision-related effects

In controlled trials, a higher proportion of patients treated with pregabalin reported blurred vision than did patients treated with placebo which resolved in a majority of cases with continued dosing. In the clinical studies where ophthalmologic testing was conducted, the incidence of visual acuity reduction and visual field changes was greater in pregabalin-treated patients than in placebo-treated patients; the incidence of fundoscopic changes was greater in placebo-treated patients (see section 5.1).

In the post marketing experience, visual adverse reactions have also been reported, including loss of vision, visual blurring or other changes of visual acuity, many of which were transient.

Discontinuation of pregabalin may result in resolution or improvement of these visual symptoms.

Renal failure

Cases of renal failure have been reported and in some cases discontinuation of pregabalin did show reversibility of this adverse reaction.

Withdrawal of concomitant anti-epileptic medicinal products

There are insufficient data for the withdrawal of concomitant anti-epileptic medicinal products, once seizure control with pregabalin in the add-on situation has been reached, in order to reach monotherapy on pregabalin.

Withdrawal symptoms

After discontinuation of short-term and long-term treatment with pregabalin, withdrawal symptoms have been observed. The following events have been mentioned: insomnia, headache, nausea, anxiety, diarrhoea, flu syndrome, nervousness, depression, pain, convulsion, hyperhidrosis and dizziness. The occurrence of withdrawal symptoms following discontinuation of pregabalin may indicate drug dependence (see section 4.8). The patient should be informed about this at the start of the treatment. If pregabalin should be discontinued, it is recommended this should be done gradually over a minimum of 1 week independent of the indication (see section 4.2).

Convulsions, including status epilepticus and grand mal convulsions, may occur during pregabalin use or shortly after discontinuing pregabalin.

Concerning discontinuation of long-term treatment of pregabalin, data suggest that the incidence and severity of withdrawal symptoms may be dose-related.

Congestive heart failure

There have been postmarketing reports of congestive heart failure in some patients receiving pregabalin. These reactions are mostly seen in elderly cardiovascular compromised patients during pregabalin treatment for a neuropathic indication. Pregabalin should be used with caution in these patients. Discontinuation of pregabalin may resolve the reaction.

Treatment of central neuropathic pain due to spinal cord injury

In the treatment of central neuropathic pain due to spinal cord injury the incidence of adverse reactions in general, central nervous system adverse reactions and especially somnolence was increased. This may be attributed to an additive effect due to concomitant medicinal products (e.g. anti-spasticity agents) needed for this condition. This should be considered when prescribing pregabalin in this condition.

Respiratory depression

There have been reports of severe respiratory depression in relation to pregabalin use. Patients with compromised respiratory function, respiratory or neurological disease, renal impairment, concomitant use of CNS depressants and the elderly may be at higher risk of experiencing this severe adverse reaction. Dose adjustments may be necessary in these patients (see section 4.2).

Suicidal ideation and behaviour

Suicidal ideation and behaviour have been reported in patients treated with anti-epileptic agents in several indications. A meta-analysis of randomised placebo controlled studies of anti-epileptic drugs has also shown a small increased risk of suicidal ideation and behaviour. The mechanism of this risk is not known and the available data do not exclude the possibility of an increased risk for pregabalin.

Therefore patients should be monitored for signs of suicidal ideation and behaviours and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.

Reduced lower gastrointestinal tract function

There are post marketing reports of events related to reduced lower gastrointestinal tract function (e.g. intestinal obstruction, paralytic ileus, constipation) when pregabalin was co-administered with medications that have the potential to produce constipation, such as opioid analgesics. When pregabalin and opioids will be used in combination, measures to prevent constipation may be considered (especially in female patients and elderly).

Concomitant use with opioids

Caution is advised when prescribing pregabalin concomitantly with opioids due to risk of CNS depression (see section 4.5). In a case-control study of opioid users, those patients who took pregabalin concomitantly with an opioid had an increased risk for opioid-related death compared to opioid use alone (adjusted odds ratio [aOR], 1.68 [95% CI, 1.19 – 2.36]). This increased risk was observed at low doses of pregabalin (≤ 300 mg, aOR 1.52 [95% CI, 1.04 – 2.22]) and there was a trend for a greater risk at high doses of pregabalin (> 300 mg, aOR 2.51 [95% CI 1.24 – 5.06]).

Misuse, abuse potential or dependence

Pregabalin can cause drug dependence, which may occur at therapeutic doses. Cases of abuse and misuse have been reported. Patients with a history of substance abuse may be at higher risk for pregabalin misuse, abuse and dependence, and pregabalin should be used with caution in such patients. Before prescribing Pregabalin, the patient's risk of misuse, abuse or dependence should be carefully evaluated.

Patients treated with pregabalin should be monitored for symptoms of pregabalin misuse, abuse or dependence, such as development of tolerance, dose escalation and drug-seeking behaviour.

Encephalopathy

Cases of encephalopathy have been reported, mostly in patients with underlying conditions that may precipitate encephalopathy.

Severe cutaneous adverse reactions (SCARs) including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), which can be life-threatening or fatal, have been reported rarely in association with pregabalin treatment. At the time of prescription patients should be advised of the signs and symptoms and monitored closely for skin reactions. If signs and symptoms suggestive of these reactions appear, pregabalin should be withdrawn immediately and an alternative treatment considered (as appropriate).

4.5 Interaction with other medicinal products and other forms of interaction

Since pregabalin is predominantly excreted unchanged in the urine, undergoes negligible metabolism in humans ($< 2\%$ of a dose recovered in urine as metabolites), does not inhibit drug metabolism *in vitro*, and is not bound to plasma proteins, it is unlikely to produce, or be subject to, pharmacokinetic interactions.

In vivo studies and population pharmacokinetic analysis

Accordingly, in *in vivo* studies no clinically relevant pharmacokinetic interactions were observed between pregabalin and phenytoin, carbamazepine, valproic acid, lamotrigine, gabapentin, lorazepam, oxycodone or ethanol. Population pharmacokinetic analysis indicated that oral antidiabetics, diuretics, insulin, phenobarbital, tiagabine and topiramate had no clinically significant effect on pregabalin clearance.

Oral contraceptives, norethisterone and/or ethinyl oestradiol

Co-administration of pregabalin with the oral contraceptives norethisterone and/or ethinyl oestradiol does not influence the steady-state pharmacokinetics of either substance.

Central nervous system influencing medical products

Pregabalin may potentiate the effects of ethanol and lorazepam. In the post marketing experience, there are reports of respiratory failure, coma and deaths in patients taking pregabalin and opioids and/or other central nervous system (CNS) depressant medicinal products. Pregabalin appears to be additive in the impairment of cognitive and gross motor function caused by oxycodone.

Interactions and the elderly

No specific pharmacodynamic interaction studies were conducted in elderly volunteers. Interaction studies have only been performed in adults.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential/Contraception

Women of childbearing potential have to use effective contraception during treatment (see section 4.4).

Pregnancy

Studies in animals have shown reproductive toxicity (see section 5.3).

Pregabalin has been shown to cross the placenta in rats (see section 5.2). Pregabalin may cross the human placenta.

Major congenital malformations

Data from a Nordic observational study of more than 2700 pregnancies exposed to pregabalin in the first trimester showed a higher prevalence of major congenital malformations (MCM) among the paediatric population (live or stillborn) exposed to pregabalin compared to the unexposed population (5.9% vs. 4.1%).

The risk of MCM among the paediatric population exposed to pregabalin in the first trimester was slightly higher compared to unexposed population (adjusted prevalence ratio and 95% confidence interval: 1.14 (0.96-1.35)), and compared to population exposed to lamotrigine (1.29 (1.01–1.65)) or to duloxetine (1.39 (1.07–1.82)).

The analyses on specific malformations showed higher risks for malformations of the nervous system, the eye, orofacial clefts, urinary malformations and genital malformations, but numbers were small and estimates imprecise.

This medicine should not be used during pregnancy unless clearly necessary (if the benefit to the mother clearly outweighs the potential risk to the foetus).

Breast-feeding

Pregabalin is excreted into human milk (see section 5.2). The effect of pregabalin on newborns/infants is unknown. A decision must be made whether to discontinue breast-feeding or to discontinue pregabalin therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

Fertility

There are no clinical data on the effects of pregabalin on female fertility.

In a clinical trial to assess the effect of pregabalin on sperm motility, healthy male subjects were exposed to pregabalin at a dose of 600 mg/day. After 3 months of treatment, there were no effects on sperm motility.

A fertility study in female rats has shown adverse reproductive effects. Fertility studies in male rats have shown adverse reproductive and developmental effects. The clinical relevance of these findings is unknown (see section 5.3).

4.7 Effects on ability to drive and use machines

Pregabalin may have minor or moderate influence on the ability to drive and use machines. Pregabalin may cause dizziness and somnolence and therefore may influence the ability to drive or use machines. Patients are advised not to drive, operate complex machinery or engage in other potentially hazardous activities until it is known whether this medicinal product affects their ability to perform these activities.

4.8 Undesirable effects

The pregabalin clinical programme involved over 8900 patients exposed to pregabalin, of whom over 5600 were in double-blind placebo-controlled trials. The most commonly reported adverse reactions were dizziness and somnolence. Adverse reactions were usually mild to moderate in intensity. In all controlled studies, the discontinuation rate due to adverse reactions was 12% for patients receiving pregabalin and 5% for patients receiving placebo. The most common adverse reactions resulting in discontinuation from pregabalin treatment groups were dizziness and somnolence.

In table 2 below all adverse reactions, which occurred at an incidence greater than placebo and in more than one patient, are listed by class and frequency (very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data).

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

The adverse reactions listed may also be associated with the underlying disease and / or concomitant medicinal products.

In the treatment of central neuropathic pain due to spinal cord injury the incidence of adverse reactions in general, CNS adverse reactions and especially somnolence was increased (see section 4.4).

Additional reactions reported from post marketing experience are included in italics in the list below.

Table 2. Pregabalin Adverse Drug Reactions

| System Organ Class | Adverse drug reactions |
|---|---|
| Infections and infestations | |
| Common | Nasopharyngitis |
| Blood and lymphatic system disorders | |
| Uncommon | Neutropaenia |
| Immune system disorders | |
| Uncommon | <i>Hypersensitivity</i> |
| Rare | <i>Angioedema, allergic reaction</i> |
| Metabolism and nutrition disorders | |
| Common | Appetite increased |
| Uncommon | Anorexia, hypoglycaemia |
| Psychiatric disorders | |
| Common | Euphoric mood, confusion, irritability, disorientation, insomnia, libido decreased |
| Uncommon | Hallucination, panic attack, restlessness, agitation, depression, depressed mood, elevated mood, <i>aggression</i> , mood swings, depersonalisation, word finding difficulty, abnormal dreams, libido increased, anorgasmia, apathy |
| Rare | Disinhibition |
| Not known | <i>Drug dependence</i> |
| Nervous system disorders | |
| Very Common | Dizziness, somnolence, headache |
| Common | Ataxia, coordination abnormal, tremor, dysarthria, amnesia, memory impairment, disturbance in attention, paraesthesia, hypoaesthesia, sedation, balance disorder, lethargy |
| Uncommon | Syncope, stupor, myoclonus, <i>loss of consciousness</i> , psychomotor hyperactivity, dyskinesia, dizziness postural, intention tremor, nystagmus, cognitive disorder, <i>mental impairment</i> , speech disorder, hyporeflexia, hyperaesthesia, burning sensation, ageusia, <i>malaise</i> |
| Rare | <i>Convulsions</i> , parosmia, hypokinesia, dysgraphia, Parkinsonism |
| Eye disorders | |
| Common | Vision blurred, diplopia |
| Uncommon | Peripheral vision loss, visual disturbance, eye swelling, visual field defect, visual acuity reduced, eye pain, asthenopia, photopsia, dry eye, lacrimation increased, eye irritation |
| Rare | <i>Vision loss, keratitis</i> , oscillopsia, altered visual depth perception, |

| | |
|---|--|
| | mydriasis, strabismus, visual brightness |
| Ear and labyrinth disorders | |
| Common | Vertigo |
| Uncommon | Hyperacusis |
| Cardiac disorders | |
| Uncommon | Tachycardia, atrioventricular block first degree, sinus bradycardia, <i>congestive heart failure</i> |
| Rare | <i>QT prolongation, sinus tachycardia, sinus arrhythmia</i> |
| Vascular disorders | |
| Uncommon | Hypotension, hypertension, hot flushes, flushing, peripheral coldness |
| Respiratory, thoracic and mediastinal disorders | |
| Uncommon | Dyspnoea, epistaxis, cough, nasal congestion, rhinitis, snoring, nasal dryness |
| Rare | <i>Pulmonary oedema, throat tightness,</i> |
| Not known | Respiratory depression |
| Gastrointestinal disorders | |
| Common | Vomiting, <i>nausea</i> , constipation, <i>diarrhoea</i> , flatulence, abdominal distension, dry mouth |
| Uncommon | Gastrooesophageal reflux disease, salivary hypersecretion, hypoaesthesia oral |
| Rare | <i>Ascites, pancreatitis, swollen tongue, dysphagia</i> |
| Hepatobiliary disorders | |
| Uncommon | Elevated liver enzymes* |
| Rare | Jaundice |
| Very rare | Hepatic failure, hepatitis |
| Skin and subcutaneous tissue disorders | |
| Uncommon | Rash papular, urticaria, hyperhidrosis, <i>pruritus</i> |
| Rare | <i>Stevens Johnson syndrome, cold sweat, Toxic Epidermal Necrolysis</i> |
| Musculoskeletal and connective tissue disorders | |
| Common | Muscle cramp, arthralgia, back pain, pain in limb, cervical spasm |
| Uncommon | Joint swelling, myalgia, muscle twitching, neck pain, muscle stiffness |
| Rare | Rhabdomyolysis |
| Renal and urinary disorders | |
| Uncommon | Urinary incontinence, dysuria |
| Rare | Renal failure, oliguria, <i>urinary retention</i> |
| Reproductive system and breast disorders | |
| Common | Erectile dysfunction |
| Uncommon | Sexual dysfunction, ejaculation delayed, dysmenorrhoea, breast pain |
| Rare | Amenorrhoea, breast discharge, breast enlargement, <i>gynaecomastia</i> |
| General disorders and administration site conditions | |
| Common | Oedema peripheral, oedema, gait abnormal, fall, feeling drunk, feeling abnormal, fatigue |
| Uncommon | Generalised oedema, <i>face oedema</i> , chest tightness, pain, pyrexia, thirst, chills, asthenia |
| Investigations | |
| Common | Weight increased |

| | |
|----------|--|
| Uncommon | Blood creatine phosphokinase increased, blood glucose increased, platelet count decreased, blood creatinine increased, blood potassium decreased, weight decreased |
| Rare | White blood cell count decreased |

* Alanine aminotransferase increased (ALT) and aspartate aminotransferase increased (AST).

After discontinuation of short-term and long-term treatment with pregabalin withdrawal symptoms have been observed. The following reactions have been reported: insomnia, headache, nausea, anxiety, diarrhoea, flu syndrome, convulsions, nervousness, depression, pain, hyperhidrosis and dizziness. These symptoms may indicate drug dependence. The patient should be informed about this at the start of the treatment.

Concerning discontinuation of long-term treatment of pregabalin, data suggest that the incidence and severity of withdrawal symptoms may be dose-related (see section 4.4).

Paediatric population

The pregabalin safety profile observed in five paediatric studies in patients with partial seizures with or without secondary generalization (12-week efficacy and safety study in patients 4 to 16 years of age, n=295; 14 day efficacy and safety study in patients 1 month to younger than 4 years of age, n=175; pharmacokinetic and tolerability study, n=65; and two 1 year open label follow on safety study, n=54 and n=431) was similar to that observed in the adult studies of patients with epilepsy. The most common adverse events observed in the 12-week study with pregabalin treatment were somnolence, pyrexia, upper respiratory tract infection, increased appetite, weight increased, and nasopharyngitis. The most common adverse events observed in the 14 day study with pregabalin treatment were somnolence, upper respiratory tract infection, and pyrexia (see sections 4.2, 5.1 and 5.2).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via

HPRA Pharmacovigilance

Website: www.hpra.ie

4.9 Overdose

In the post marketing experience, the most commonly reported adverse reactions observed when pregabalin was taken in overdose included somnolence, confusional state, agitation, and restlessness. Seizures were also reported.

In rare occasions, cases of coma have been reported.

Treatment of pregabalin overdose should include general supportive measures and may include haemodialysis if necessary (see section 4.2 Table 1).

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Anti-epileptics, other anti-epileptics

ATC code: N03AX16

The active substance, pregabalin, is a gamma-aminobutyric acid analogue [(S)-3-(aminomethyl)-5-methylhexanoic acid].

Mechanism of action

Pregabalin binds to an auxiliary subunit (α_2 - δ protein) of voltage-gated calcium channels in the central nervous system.

Clinical efficacy and safety

Neuropathic pain

Efficacy has been shown in trials in diabetic neuropathy, post herpetic neuralgia and spinal cord injury. Efficacy has not been studied in other models of neuropathic pain.

Pregabalin has been studied in 10 controlled clinical trials of up to 13 weeks with twice a day dosing (BID) and up to 8 weeks with three times a day (TID) dosing. Overall, the safety and efficacy profiles for BID and TID dosing regimens were similar.

In clinical trials up to 12 weeks for both peripheral and central neuropathic pain, a reduction in pain was seen by week 1 and was maintained throughout the treatment period.

In controlled clinical trials in peripheral neuropathic pain 35% of the pregabalin treated patients and 18% of the patients on placebo had a 50% improvement in pain score. For patients not experiencing somnolence, such an improvement was observed in 33% of patients treated with pregabalin and 18% of patients on placebo. For patients who experienced somnolence the responder rates were 48% on pregabalin and 16% on placebo.

In the controlled clinical trial in central neuropathic pain 22% of the pregabalin treated patients and 7% of the patients on placebo had a 50% improvement in pain score.

Epilepsy

Adjunctive Treatment

Pregabalin has been studied in 3 controlled clinical trials of 12 week duration with either BID or TID dosing. Overall, the safety and efficacy profiles for BID and TID dosing regimens were similar.

A reduction in seizure frequency was observed by Week 1.

Paediatric population

The efficacy and safety of pregabalin as adjunctive treatment for epilepsy in paediatric patients below the age of 12 and adolescents has not been established. The adverse events observed in a pharmacokinetic and tolerability study that enrolled patients from 3 months to 16 years of age (n=65) with partial onset seizures were similar to those observed in adults. Results of a 12-week placebo-controlled study of 295 paediatric patients aged 4 to 16 years and a 14-day placebo controlled study of 175 paediatric patients aged 1 month to younger than 4 years of age performed to evaluate the efficacy and safety of pregabalin as adjunctive therapy for the treatment of partial onset seizures and two 1 year open label safety studies in 54 and 431 paediatric patients respectively, from 3 months to 16 years of age with epilepsy indicate that the adverse events of pyrexia and upper respiratory infections were observed more frequently than in adult studies of patients with epilepsy (see sections 4.2, 4.8 and 5.2).

In the 12-week placebo-controlled study, paediatric patients (4 to 16 years of age) were assigned to pregabalin 2.5 mg/kg/day (maximum, 150 mg/day), pregabalin 10 mg/kg/day (maximum, 600 mg/day), or placebo. The percentage of subjects with at least a 50% reduction in partial onset seizures as compared to baseline was 40.6% of subjects treated with pregabalin 10 mg/kg/day (p=0.0068 versus placebo), 29.1% of subjects treated with pregabalin 2.5 mg/kg/day (p=0.2600 versus placebo) and 22.6% of those receiving placebo.

In the 14-day placebo-controlled study, paediatric patients (1 month to younger than 4 years of age) were assigned to pregabalin 7 mg/kg/day, pregabalin 14 mg/kg/day, or placebo. Median 24-hour seizure frequencies at baseline and at the final visit were 4.7 and 3.8 for pregabalin 7 mg/kg/day, 5.4 and 1.4 for pregabalin 14 mg/kg/day, and 2.9 and 2.3 for placebo, respectively. Pregabalin 14 mg/kg/day significantly reduced the log-transformed partial onset seizure frequency versus placebo (p=0.0223); pregabalin 7 mg/kg/day did not show improvement relative to placebo.

In a 12 week placebo-controlled study in subjects with Primary Generalized Tonic-Clonic (PGTC) seizures 219 subjects (aged 5 to 65 years, of which 66 were aged 5 to 16 years) were assigned to pregabalin 5 mg/kg/day (maximum 300 mg/day), 10 mg/kg/day (maximum 600 mg/day) or placebo as adjunctive therapy. The percentage of subjects with at least a 50% reduction in PGTC seizure rate was 41.3%, 38.9% and 41.7% for pregabalin 5 mg/kg/day, pregabalin 10 mg/kg/day and placebo respectively.

Monotherapy (newly diagnosed patients)

Pregabalin has been studied in 1 controlled clinical trial of 56 week duration with BID dosing. Pregabalin did not achieve non-inferiority to lamotrigine based on the 6-month seizure freedom endpoint. Pregabalin and lamotrigine were similarly safe and well tolerated.

Generalised Anxiety Disorder

Pregabalin has been studied in 6 controlled trials of 4-6 week duration, an elderly study of 8 week duration and a long-term relapse prevention study with a double blind relapse prevention phase of 6 months duration.

Relief of the symptoms of GAD as reflected by the Hamilton Anxiety Rating Scale (HAM-A) was observed by Week 1.

In controlled clinical trials (4-8 week duration) 52% of the pregabalin treated patients and 38% of the patients on placebo had at least a 50% improvement in HAM-A total score from baseline to endpoint.

In controlled trials, a higher proportion of patients treated with pregabalin reported blurred vision than did patients treated with placebo which resolved in a majority of cases with continued dosing.

Ophthalmologic testing (including visual acuity testing, formal visual field testing and dilated fundoscopic examination) was conducted in over 3600 patients within controlled clinical trials. In these patients, visual acuity was reduced in 6.5% of patients treated with pregabalin, and 4.8% of placebo-treated patients. Visual field changes were detected in 12.4% of pregabalin-treated, and 11.7% of placebo-treated patients. Fundoscopic changes were observed in 1.7% of pregabalin-treated and 2.1% of placebo-treated patients.

5.2 Pharmacokinetic properties

Pharmacotherapeutic group: Anti-epileptics, other anti-epileptics

ATC code: N03AX16

The active substance, pregabalin, is a gamma-aminobutyric acid analogue [(S)-3-(aminomethyl)-5-methylhexanoic acid].

Mechanism of action

Pregabalin binds to an auxiliary subunit ($\alpha_2\text{-}\delta$ protein) of voltage-gated calcium channels in the central nervous system.

Clinical efficacy and safety

Neuropathic pain

Efficacy has been shown in trials in diabetic neuropathy, post herpetic neuralgia and spinal cord injury. Efficacy has not been studied in other models of neuropathic pain.

Pregabalin has been studied in 10 controlled clinical trials of up to 13 weeks with twice a day dosing (BID) and up to 8 weeks with three times a day (TID) dosing. Overall, the safety and efficacy profiles for BID and TID dosing regimens were similar.

In clinical trials up to 12 weeks for both peripheral and central neuropathic pain, a reduction in pain was seen by week 1 and was maintained throughout the treatment period.

In controlled clinical trials in peripheral neuropathic pain 35% of the pregabalin treated patients and 18% of the patients on placebo had a 50% improvement in pain score. For patients not experiencing somnolence, such an improvement was observed in 33% of patients treated with pregabalin and 18% of patients on placebo. For patients who experienced somnolence the responder rates were 48% on pregabalin and 16% on placebo.

In the controlled clinical trial in central neuropathic pain 22% of the pregabalin treated patients and 7% of the patients on placebo had a 50% improvement in pain score.

Epilepsy

Adjunctive Treatment

Pregabalin has been studied in 3 controlled clinical trials of 12 week duration with either BID or TID dosing. Overall, the safety and efficacy profiles for BID and TID dosing regimens were similar.

A reduction in seizure frequency was observed by Week 1.

Paediatric population

The efficacy and safety of pregabalin as adjunctive treatment for epilepsy in paediatric patients below the age of 12 and adolescents has not been established. The adverse events observed in a pharmacokinetic and tolerability study that enrolled patients from 3 months to 16 years of age (n=65) with partial onset seizures were similar to those observed in adults. Results of a 12-week placebo-controlled study of 295 paediatric patients aged 4 to 16 years and a 14-day placebo controlled study of 175 paediatric patients aged 1 month to younger than 4 years of age performed to evaluate the efficacy and safety of pregabalin as adjunctive therapy for the treatment of partial onset seizures and two 1 year open label safety studies in 54 and 431 paediatric patients respectively, from 3 months to 16 years of age with epilepsy indicate that the adverse events of pyrexia and upper respiratory infections were observed more frequently than in adult studies of patients with epilepsy (see sections 4.2, 4.8 and 5.2).

In the 12-week placebo-controlled study, paediatric patients (4 to 16 years of age) were assigned to pregabalin 2.5 mg/kg/day (maximum, 150 mg/day), pregabalin 10 mg/kg/day (maximum, 600 mg/day), or placebo. The percentage of subjects with at least a 50% reduction in partial onset seizures as compared to baseline was 40.6% of subjects treated with pregabalin 10 mg/kg/day (p=0.0068 versus placebo), 29.1% of subjects treated with pregabalin 2.5 mg/kg/day (p=0.2600 versus placebo) and 22.6% of those receiving placebo.

In the 14-day placebo-controlled study, paediatric patients (1 month to younger than 4 years of age) were assigned to pregabalin 7 mg/kg/day, pregabalin 14 mg/kg/day, or placebo. Median 24-hour seizure frequencies at baseline and at the final visit were 4.7 and 3.8 for pregabalin 7 mg/kg/day, 5.4 and 1.4 for pregabalin 14 mg/kg/day, and 2.9 and 2.3 for placebo, respectively. Pregabalin 14 mg/kg/day significantly reduced the log-transformed partial onset seizure frequency versus placebo ($p=0.0223$); pregabalin 7 mg/kg/day did not show improvement relative to placebo.

Monotherapy (newly diagnosed patients)

Pregabalin has been studied in 1 controlled clinical trial of 56 week duration with BID dosing. Pregabalin did not achieve non-inferiority to lamotrigine based on the 6-month seizure freedom endpoint. Pregabalin and lamotrigine were similarly safe and well tolerated.

Generalised Anxiety Disorder

Pregabalin has been studied in 6 controlled trials of 4-6 week duration, an elderly study of 8 week duration and a long-term relapse prevention study with a double blind relapse prevention phase of 6 months duration.

Relief of the symptoms of GAD as reflected by the Hamilton Anxiety Rating Scale (HAM-A) was observed by Week 1.

In controlled clinical trials (4-8 week duration) 52% of the pregabalin treated patients and 38% of the patients on placebo had at least a 50% improvement in HAM-A total score from baseline to endpoint.

In controlled trials, a higher proportion of patients treated with pregabalin reported blurred vision than did patients treated with placebo which resolved in a majority of cases with continued dosing.

Ophthalmologic testing (including visual acuity testing, formal visual field testing and dilated funduscopic examination) was conducted in over 3600 patients within controlled clinical trials. In these patients, visual acuity was reduced in 6.5% of patients treated with pregabalin, and 4.8% of placebo-treated patients. Visual field changes were detected in 12.4% of pregabalin-treated, and 11.7% of placebo-treated patients. Funduscopic changes were observed in 1.7% of pregabalin-treated and 2.1% of placebo-treated patients.

5.3 Preclinical safety data

In conventional safety pharmacology studies in animals, pregabalin was well-tolerated at clinically relevant doses. In repeated dose toxicity studies in rats and monkeys CNS effects were observed, including hypoactivity, hyperactivity and ataxia. An increased incidence of retinal atrophy commonly observed in aged albino rats was seen after long-term exposure to pregabalin at exposures ≥ 5 times the mean human exposure at the maximum recommended clinical dose.

Pregabalin was not teratogenic in mice, rats or rabbits. Foetal toxicity in rats and rabbits occurred only at exposures sufficiently above human exposure. In prenatal/postnatal toxicity studies, pregabalin induced offspring developmental toxicity in rats at exposures > 2 times the maximum recommended human exposure.

Adverse effects on fertility in male and female rats were only observed at exposures sufficiently in excess of therapeutic exposure. Adverse effects on male reproductive organs and sperm parameters were reversible and occurred only at exposures sufficiently in excess of therapeutic exposure or were associated with spontaneous degenerative processes in male reproductive organs in the rat. Therefore the effects were considered of little or no clinical relevance.

Pregabalin is not genotoxic based on results of a battery of *in vitro* and *in vivo* tests.

Two-year carcinogenicity studies with pregabalin were conducted in rats and mice. No tumours were observed in rats at exposures up to 24 times the mean human exposure at the maximum recommended clinical dose of 600 mg/day. In mice, no increased incidence of tumours was found at exposures similar to the mean human exposure, but an increased incidence of haemangiosarcoma was observed at higher exposures. The non-genotoxic mechanism of pregabalin-induced tumour formation in mice involves platelet changes and associated endothelial cell proliferation. These platelet changes were not present in rats or in humans based on short-term and limited long-term clinical data. There is no evidence to suggest an associated risk to humans.

In juvenile rats the types of toxicity do not differ qualitatively from those observed in adult rats. However, juvenile rats are more sensitive. At therapeutic exposures, there was evidence of CNS clinical signs of hyperactivity and bruxism and some changes in growth (transient body weight gain suppression). Effects on the oestrus cycle were observed at 5-fold the human therapeutic exposure. Reduced acoustic startle response was observed in juvenile rats 1-2 weeks after exposure at > 2 times the human therapeutic exposure. Nine weeks after exposure, this effect was no longer observable.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content:

Mannitol
Maize starch
Talc

Capsule Shell:

Gelatin
Titanium dioxide (E171)

Printing ink:

Shellac (E904)
Propylene glycol (E1520)
Strong ammonia solution (E527)
Black iron oxide (E172)
Potassium hydroxide (E525)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

Clear PVC-Alu or Alu-Alu blister packs containing 14, 21, 56, 70, 84, 100 or 112 hard capsules.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

No special requirements for disposal.

7 MARKETING AUTHORISATION HOLDER

Brillpharma (Ireland) Limited
Inniscarra
Main Street
Rathcoole
Dublin
Ireland

8 MARKETING AUTHORISATION NUMBER

PA22749/015/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 4th August 2017
Date of last renewal: 5th March 2022

10 DATE OF REVISION OF THE TEXT

July 2023