

Summary of Product Characteristics

1 NAME OF THE VETERINARY MEDICINAL PRODUCT

Porcilis Ery + Parvo suspension for injection for pigs

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each dose (2 ml) contains:

Active substances:

- Inactivated Porcine Parvovirus strain 014: ≥ 552 EU*
- Inactivated lysed antigen concentrate of *Erysipelothrix rhusiopathiae* strain M2 (serotype 2): ≥ 1 pig protective dose (ppd)**

* as determined in the final product by antigenic mass ELISA

** as measured in the Ph. Eur. potency test

Adjuvant:

dl- α -tocopherol: 150 mg

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Suspension for injection.

Aqueous white or nearly white liquid.

4 CLINICAL PARTICULARS

4.1 Target Species

Pigs (sows and gilts)

4.2 Indications for use, specifying the target species

For active immunization of sows and gilts to prevent clinical signs of Erysipelas disease caused by all relevant *Erysipelothrix (E.) rhusiopathiae* serotypes (serotype 1 and 2) and for protection against embryonal and fetal death caused by porcine parvovirus (PPV) infection.

E. rhusiopathiae:

Onset of immunity: 3 weeks

Duration of immunity: 6 months

Porcine parvovirus:

Duration of immunity: 12 months

4.3 Contraindications

None.

4.4 Special warnings for each target species

Vaccinate healthy animals only.

4.5 Special precautions for use

Special precautions for use in animals

Sick and weak animals should not be vaccinated.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

In case of accidental self-injection, seek medical advice immediately and show the package leaflet or label to the physician.

4.6 Adverse reactions (frequency and seriousness)

In laboratory studies and field trials:

Transient increases in body temperature (0.5°C) within 24 hours may very commonly occur. Mild transient local swelling (Ø 1-10mm) until 8 days after vaccination may very commonly occur. Transient reluctance to move may commonly occur.

In post marketing experience:

In very rare cases, a hypersensitivity reaction may occur.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

4.7 Use during pregnancy, lactation or lay

Can be used during pregnancy and lactation.

4.8 Interaction with other medicinal products and other forms of interactions

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

4.9 Amounts to be administered and administration route

Administer one dose of 2 ml by deep intramuscular injection behind the ear.

Before use, allow the vaccine to reach room temperature. Shake well before use.

Use sterile syringe and needles. Avoid introduction of contamination by multiple broaching.

Primary vaccination course:

Protection against *E. rhusiopathiae* and PPV should be achieved in gilts before first mating. A single injection not later than 2 weeks before mating is sufficient to protect the following pregnancy from damage due to PPV. For the induction of protection against Erysipelas a double vaccination as a basic vaccination is advised. This can be achieved with the single Erysipelas vaccine either 4 weeks before or 4 weeks after the application of the combined ERY-PARVO vaccine.

Due to possible interference with maternal antibodies the pigs should have reached the age of 6 months before vaccination to ensure efficacy against porcine parvovirus.

Revaccinations should be given once a year, supplemented with the administration of a single Erysipelas vaccine, 6 months post each Porcilis Ery + Parvo vaccination.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Reactions observed after administration of a double dose are not different from those observed after administration of a single dose.

4.11 Withdrawal period(s)

Zero Days.

5 PHARMACOLOGICAL or IMMUNOLOGICAL PROPERTIES

Pharmacotherapeutic group: Inactivated bacterial and inactivated viral vaccines

ATCvet code: QI09AL01

The active substances are a lysate of *E. rhusiopathiae* strain M2 (serotype 2) and inactivated porcine parvo virus strain 014. For the active immunization of sows and gilts, as an aid in the control of swine erysipelas and for the protection of their embryos and fetuses against porcine parvovirus infection.

The antigens are incorporated in an aqueous tocopherol based adjuvant in order to enhance a prolonged stimulation of immunity.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Polysorbate 80

Tris (hydroxymethyl) aminomethane

Sodium chloride

Simethicone

Water for injections

6.2 Major incompatibilities

Do not mix with any other veterinary medicinal product.

6.3 Shelf-life

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf life after first opening the immediate packaging: 10 hours

6.4 Special precautions for storage

Store in a refrigerator (2°C - 8°C). Do not freeze. Protect from light.

6.5 Nature and composition of immediate packaging

Vial of glass, hydrolytical class type II or type I (Ph. Eur.), or PET-flask closed with a halogenobutyl rubber stopper and sealed with a coded aluminium cap.

Package sizes:

1 vial of 20 ml (10 doses), 50 ml (25 doses), 100 ml (50 doses) or 250 ml (125 doses) packed in a cardboard box.

Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

Intervet Ireland Limited
Magna Drive
Magna Business Park, Citywest Road
Dublin 24
Ireland

8 MARKETING AUTHORISATION NUMBER(S)

VPA10996/097/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 4th May 2001
Date of last renewal: 7th July 2006

10 DATE OF REVISION OF THE TEXT

October 2019