

# Summary of Product Characteristics

## 1 NAME OF THE VETERINARY MEDICINAL PRODUCT

Bovilis BVD Suspension for injection for cattle

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each dose of 2 ml contains:

### Active component:

Inactivated cytopathogenic bovine viral diarrhoea (BVD) virus type 1 strain C-86, containing 50 ELISA Units (EU) and inducing at least 4.6 log<sub>2</sub> VN units\*

\*Mean virus neutralizing titre obtained in the potency test

### Adjuvant:

Aluminium 3+ (as Al-phosphate and Al-hydroxide): 6-9 mg

### Excipients:

Methyl parahydroxybenzoate: 3 mg (preservative)

For the full list of excipients, see section 6.1

## 3 PHARMACEUTICAL FORM

Suspension for injection

Red to pink-coloured turbid suspension

## 4 CLINICAL PARTICULARS

### 4.1 Target Species

Cattle (cows and heifers)

### 4.2 Indications for use, specifying the target species

For active immunisation of cows and heifers from eight months of age onwards to protect the foetus against transplacental infection with bovine viral diarrhoea virus.

### 4.3 Contraindications

None.

### 4.4 Special warnings for each target species

Vaccinate healthy animals only.

### 4.5 Special precautions for use

Special precautions for use in animals

Not applicable.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

In the case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the physician.

#### 4.6 Adverse reactions (frequency and seriousness)

In very rare cases a slight swelling may be observed for 14 days at the site of injection. Also in very rare cases transient mild pyrexia may occur. In very rare cases, hypersensitivity reactions including anaphylactic shock may occur. In the event of anaphylactic type reactions, appropriate treatment such as with antihistamine, corticosteroid or adrenaline is recommended. The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

#### 4.7 Use during pregnancy, lactation or lay

Can be used during pregnancy.

#### 4.8 Interaction with other medicinal products and other forms of interactions

Safety and efficacy data are available which demonstrate that for re-vaccination in cattle from 15 months of age onwards (i.e. those that have previously been vaccinated separately with Bovilis BVD and Bovilis IBR marker live) - this vaccine can be mixed and administered with Bovilis IBR marker live (in Member States where this veterinary medicinal product is authorised). The product literature of Bovilis IBR marker live should be consulted before administration of the mixed products. The adverse effects observed after administration of one dose or an overdose of the mixed vaccines are not different from those described for the vaccines administered separately.

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product except the product mentioned above. A decision to use this vaccine before or after any other veterinary medicinal product therefore needs to be made on a case by case basis.

#### 4.9 Amounts to be administered and administration route

Before using the vaccine allow it to reach ambient temperature (15°C - 25°C).

Shake well before use. Use sterile syringes and needles.

Intramuscular injection. 2 ml per animal.

All cattle can be vaccinated from an age of eight months onwards.

Foetal protection can be expected if the primary immunisation has been finalised 4 weeks before start of the gestation. Animals which are vaccinated later than 4 weeks before gestation or during the early gestation will not be protected against foetal infection.

##### Individual vaccination

###### *Basic immunisation*

Two vaccinations with an interval of 4 weeks. The second vaccination should be given not later than 4 weeks before the start of the gestation.

###### *Revaccination*

One vaccination 4 weeks before start of the next gestation.

##### Herd vaccination

###### *Basic immunisation*

Two vaccinations with an interval of 4 weeks. For use in cattle from eight months of age, all animals should be vaccinated.

###### *Revaccination*

One vaccination 6 months after basic vaccination with next re-vaccinations at an interval no greater than 12 months.

For revaccination, the vaccine may be used for reconstitution of Bovilis IBR marker live for use in cattle from 15 months of age (i.e. those that have previously been vaccinated separately with Bovilis BVD and Bovilis IBR marker live) and the following instructions should be used:

Bovilis IBR marker live	Bovilis BVD
5 doses	+ 10 ml
10 doses	+ 20 ml
25 doses	+ 50 ml
50 doses	+ 100 ml

A single dose (2 ml) of Bovilis BVD mixed with Bovilis IBR marker live is given intramuscularly.

Visual appearance after reconstitution of Bovilis IBR marker live in Bovilis BVD:  
As specified for Bovilis BVD alone.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

Reactions after a double dose are not different from those after the single dose.

#### **4.11 Withdrawal period(s)**

Zero days.

### **5 PHARMACOLOGICAL or IMMUNOLOGICAL PROPERTIES**

Pharmacotherapeutic group: inactivated bovine viral diarrhoea vaccine.

ATC-vet code: QI02AA01.

Bovilis BVD is an adjuvanted aqueous inactivated viral vaccine for active immunisation of cows and heifers against transplacental infection with bovine viral diarrhoea virus.

### **6 PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Aluminium phosphate  
Aluminium hydroxide  
Methylparahydroxybenzoate  
Propylene glycol  
Tromethamine  
Tissue culture medium  
Hydrochloric acid solution or tromethamine solution  
Water for injections

#### **6.2 Major incompatibilities**

Do not mix with any other veterinary medicinal product, except with Bovilis IBR marker live (for revaccination only).

#### **6.3 Shelf-life**

Shelf life of the veterinary medicinal product as packaged for sale: 2 years.

Shelf life after first opening the immediate packaging: 10 hours.

Shelf life after mixing with Bovilis IBR marker live: 3 hours (at room temperature).

#### **6.4 Special precautions for storage**

Store in a refrigerator (2 °C - 8 °C). Do not freeze.

## **6.5 Nature and composition of immediate packaging**

Vials of glass (hydrolytic type I, Ph. Eur.) or plastic (polyethylene-terephthalate, PET) closed with a rubber (halogenobutyl) stopper and an aluminium cap.

Package sizes:

Carton box containing 1 glass or plastic vial of 2 ml (1 dose)

Carton box containing 1 glass or plastic vial of 10 ml (5 doses)

Carton box containing 1 glass or plastic vial of 20 ml (10 doses)

Carton box containing 1 glass or plastic vial of 50 ml (25 doses)

Carton box containing 1 glass or plastic vial of 100 ml (50 doses)

Carton box containing 1 glass or plastic vial of 250 ml (125 doses)

Not all pack sizes may be marketed.

## **6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **7 MARKETING AUTHORISATION HOLDER**

Intervet Ireland Limited  
Magna Drive  
Magna Business Park, Citywest Road  
Dublin 24  
Ireland

## **8 MARKETING AUTHORISATION NUMBER(S)**

VPA10996/271/001

## **9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 30<sup>th</sup> August 1999

Date of latest renewal: 22<sup>nd</sup> June 2009

## **10 DATE OF REVISION OF THE TEXT**

July 2020