

Summary of Product Characteristics

1 NAME OF THE VETERINARY MEDICINAL PRODUCT

Trimectin 1.0% w/v Solution for Injection

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Active Substances:

Ivermectin 1.0% w/v

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Solution for injection.

A clear, colourless to slightly yellow coloured solution.

4 CLINICAL PARTICULARS

4.1 Target Species

Cattle and pigs.

4.2 Indications for use, specifying the target species

For the effective treatment and control of the following harmful parasites of cattle and pigs:

Cattle:

Gastro-intestinal roundworms (adult and fourth stage larvae):

Ostertagia spp. (including inhibited *O. ostertagi*)

Haemonchus placei

Trichostrongylus axei

Trichostrongylus colubriformis

Cooperia spp.

Bunostomum phlebotomum

Oesophagostomum radiatum

Strongyloides papillosus (adult)

Nematodirus helvetianus (adult)

N. spathiger (adult)

Lungworms (adult and fourth stage larvae):

Dictyocaulus viviparus.

Eyeworms (adult):

Thelazia spp.

Warbles (parasitic stages):

Hypoderma bovis

H. lineatum.

Mange mites:

Psoroptes bovis

Sarcoptes scabiei var. *bovis*.

Sucking lice:

Linognathus vituli

Haematopinus eurysternus
Solenopotes capillatus.

May also be used as an aid in the control of the mange mite *Chorioptes bovis* but complete elimination may not occur.

Persistent activity

Treatment at the recommended dose rate controls re-infection with *Ostertagia spp.* and *Cooperia spp.* acquired up to 7 days after treatment and *Dictyocaulus viviparus* acquired up to 14 days after treatment.

Pigs:

Gastrointestinal roundworms: (Adults and fourth stage larvae):

Ascaris suum
Hyostrongylus rubidus
Oesophagostomum spp.
Strongyloides ransomi (adult)

Lungworms:

Metastrongylus spp. (adult)

Lice:

Haematopinus suis

Mange mites:

Sarcoptes scabiei var. suis

4.3 Contraindications

Do not use in case of known hypersensitivity to the active ingredient.
Do not administer by the intravenous or intramuscular route.

4.4 Special warnings for each target species

None.

4.5 Special precautions for use

(i) Special Precautions for use in Animals

The product has been formulated specifically for use in cattle and pigs. It should not be used in other species as severe adverse reactions, including fatalities in dogs, may occur.

Frequent and repeated use may lead to the development of resistance. It is important that the correct dose is given in order to minimise the risk of resistance. To avoid under-dosing, animals should be grouped according to their bodyweight and dosed according to the dose of the heaviest animal in the group.

(ii) Special Precautions to be taken by the Person Administering the Product to Animals

Take care to avoid self-administration: the product may cause local irritation and/or pain at the site of injection.
Direct contact of the product with the skin should be kept to a minimum.
Do not eat, drink or smoke while handling the product.
Wash hands after use.

(iii) Other precautions

When using the 250 ml and 500 ml pack sizes, use only automatic syringe equipment. For the 50 ml pack size, use of a multiple dose syringe is recommended. To refill the syringe, use of a draw off needle is recommended to avoid excessive broaching of the stopper.

4.6 Adverse reactions (frequency and seriousness)

Cattle

Transitory discomfort has been observed in some cattle following subcutaneous administration. A low incidence of soft tissue swelling at the injection site has been observed. These reactions have disappeared without treatment.

Pigs

Mild and transient pain reactions may be seen in some pigs following subcutaneous injection. All these reactions disappeared without treatment.

4.7 Use during pregnancy, lactation or lay

Pregnancy

The product can be administered to beef cows and pigs at any stage of pregnancy.

Lactation

Do not use in dairy cows producing milk for human consumption

Do not use in non-lactating dairy cows within 60 days of calving. The product can be used in sows at any stage of lactation.

Fertility

Fertility is not affected by administration of the product.

4.8 Interaction with other medicinal products and other forms of interactions

The product can be used concurrently without adverse effects with foot and mouth disease vaccine or clostridial vaccine, given at separate injection sites.

4.9 Amounts to be administered and administration route

Each ml contains 10 mg of ivermectin sufficient to treat 50 kg of bodyweight of cattle and 33 kg of bodyweight of pigs. The volume administered per injection site should not exceed 10 ml. The injection may be given with any standard automatic or single-dose or hypodermic syringe. Use of 17 gauge x ½ inch needle is suggested. Replace with a fresh sterile needle after every 10 to 12 animals. Injection of wet or dirty animals is not recommended. If using a single-dose hypodermic syringe, use a separate sterile needle to withdraw the product from the container.

Cattle:

The product should be given only by subcutaneous injection at the recommended dosage level of 200 mcg ivermectin per kg bodyweight under the loose skin in front of, or behind, the shoulder in cattle. This is equivalent to 1 ml per 50 kg bodyweight.

Pigs:

In pigs, the recommended dosage level is 300 mcg ivermectin per kg bodyweight. This is equivalent to 1 ml per 33kg bodyweight. The recommended route of administration is by subcutaneous injection into the neck.

Young Pigs:

In young pigs, especially those below 16kg for which less than 0.5 ml of product is indicated, dosing accuracy is important. The use of a syringe that can accurately deliver as little as 0.1 ml is recommended

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Cattle

Single doses of 4.0 mg ivermectin per kg (20 x the use level) given subcutaneously resulted in ataxia and depression.

Pigs

A dose of 30 mg ivermectin per kg (100 x the recommended dose of 0.3 mg per kg) injected subcutaneously to pigs caused lethargy, ataxia, bilateral mydriasis, intermittent tremors, laboured breathing and lateral recumbency.

4.11 Withdrawal period(s)

Cattle: Must not be treated within 49 days of slaughter for human consumption. Do not use in lactating cows producing milk for human consumption. Do not use in non-lactating dairy cows including pregnant dairy heifers within 60 days of calving.

Pigs: Must not be treated within 28 days of slaughter for human consumption.

5 PHARMACOLOGICAL or IMMUNOLOGICAL PROPERTIES

ATC vet code: QP54AA01
Therapeutic group Endectocide

5.1 Pharmacodynamic properties

Ivermectin is a member of the macrocyclic lactone class of endectocides which have a unique mode of action. Compounds of the class bind selectively and with high affinity to glutamate-gated chloride ion channels which occur in invertebrate nerve and muscle cells. This leads to an increase in the permeability of the cell membrane to chloride ions with hyperpolarization of the nerve or muscle cell, resulting in paralysis and death of the parasite. Compounds of this class may also interact with other ligand-gated chloride channels, such as those gated by the neurotransmitter gamma-aminobutyric (GABA).

The margin of safety for compounds of this class is attributable to the fact that mammals do not have glutamate-gated chloride channels, the macrocyclic lactones have a low affinity for other mammalian ligand-gated chloride channels and they do not readily cross the blood-brain barrier.

5.2 Pharmacokinetic particulars

Cattle

At a dose level of 0.2 mg ivermectin per kg a mean C_{max} of 30.43 ng/ml is reached at a mean T_{max} of 131 hours and the mean half-life in plasma is 142.39 hours. It is also established that ivermectin is carried mainly in the plasma (80 %). This distribution between plasma and blood cells remains relatively constant.

Pigs

In pigs, at a dose level of 0.3 mg ivermectin per kg bodyweight, a mean C_{max} of 6.87 ng/ml was reached at a mean T_{max} of 86.75 hours, and the mean elimination half life was 133.56 hours and the drug persisted in plasma for up to 28 days.

Excretion: length of time and route

Cattle

Only about 1-2 % is excreted in the urine the remainder is excreted in the faeces, approximately 60% of which is excreted as unaltered drug. The remainder is excreted as metabolites or degradation products.

Pigs

Biliary excretion is also the major route of ivermectin excretion in pigs.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Glycerol
Glycerol formal

6.2 Major incompatibilities

None known.

6.3 Shelf-life

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.
Shelf-life after first opening the immediate packaging: 28 days.

6.4 Special precautions for storage

This veterinary medicinal product does not require any special storage precautions.

6.5 Nature and composition of immediate packaging

Multidose polyethylene bottles of 50 ml, 250 ml and 500 ml sealed with bromobutyl seals and aluminium overseals.
Not all pack sizes may be marketed.

6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials derived from the use of such products

Any unused product or waste materials should be disposed of in accordance with national requirements. The product should not enter water courses as this may be dangerous to fish and other aquatic organisms.

7 MARKETING AUTHORISATION HOLDER

Bimeda Animal Health Limited
2, 3 & 4 Airton Close
Airton Road
Tallaght
Dublin 24
Ireland

8 MARKETING AUTHORISATION NUMBER(S)

VPA22033/069/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 18 August 2006
Date of last renewal: 23 May 2007

10 DATE OF REVISION OF THE TEXT

June 2019