

# Summary of Product Characteristics

## 1 NAME OF THE VETERINARY MEDICINAL PRODUCT

Anthelmin Plus Flavour Tablets for dogs

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains:

### Active substances:

Praziquantel	50 mg
Pyrantel embonate	144 mg
Febantel	150 mg

### Excipients:

For the full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Tablets.

Round, yellow, uncoated tablets with visible darker spots and bevelled edges with cross line on one side and plain on other side.

The tablets can be divided into equal halves or equal quarters.

## 4 CLINICAL PARTICULARS

### 4.1 Target Species

Dogs (small and medium size)

### 4.2 Indications for use, specifying the target species

For the treatment of mixed infestations with the following roundworms and tapeworms in adult dogs and puppies:

Nematodes

Ascarids: *Toxocara canis*, *Toxascaris leonina* (late immature forms and mature forms)

Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults)

Cestodes

Tapeworms: *Taenia* spp., *Dipylidium caninum*

### 4.3 Contraindications

Do not use simultaneously with piperazine compounds.

Do not exceed the stated dosage when treating pregnant bitches.

Do not use in animals with a known hypersensitivity to the active substance or to any of excipients.

Not for use in dogs younger than 2 weeks of age and/or weighing less than 2 kg.

### 4.4 Special warnings for each target species

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*. Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken.

Parasite resistance to any particular class of anthelmintic may develop following frequent, repeated use of an anthelmintic of that class.

## 4.5 Special precautions for use

### Special precautions for use in animals

Any part-used tablets should be discarded.

### Special precautions to be taken by the person administering the veterinary medicinal product to animals

In the interests of good hygiene, persons administering the tablet directly to a dog or by adding it to the dog's food, should wash their hands afterwards.

In case of accidental ingestion, seek medical advice and show the package leaflet to the physician.

## 4.6 Adverse reactions (frequency and seriousness)

In very rare cases transient loose faeces, diarrhoea and/or vomiting may occur.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals displaying adverse reactions during the course of one treatment)
- common (more than 1 but less than 10 animals in 100 animals)
- uncommon (more than 1 but less than 10 animals in 1,000 animals)
- rare (more than 1 but less than 10 animals in 10,000 animals)
- very rare (less than 1 animal in 10,000 animals, including isolated reports).

## 4.7 Use during pregnancy, lactation or lay

Consult a veterinary surgeon before treating pregnant animals for roundworms.

The tablets may be used during lactation (see Section 4.3 and 4.9).

Do not use in bitches during the first two-thirds of pregnancy.

## 4.8 Interaction with other medicinal products and other forms of interaction

Do not combine with piperazine as the anthelmintic effects of pyrantel and piperazine (used in many worming products for dogs) may be antagonized.

Concurrent use with other cholinergic compounds can lead to toxicity.

## 4.9 Amounts to be administered and administration route

For oral administration.

### Dosage

The recommended dose rates are: 15 mg/kg bodyweight febantel, 14.4 mg/kg pyrantel and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 10 kg bodyweight.

Tablets may be halved/quartered to allow accuracy of dosing.

### Administration and Duration of Treatment

The tablet(s) can be given directly to the dog or disguised in food. No restriction of access to food is required either before or after administration of the product.

To ensure administration of a correct dose, body weight should be determined as accurately as possible.

Puppies may be wormed with this product from 2 weeks of age and every 2 weeks until 12 weeks of age. Thereafter they should be treated at 3 monthly intervals until 6 months of age.

For the control of *Toxocara*, nursing bitches should be dosed 2 weeks after giving birth and every 2 weeks until weaning.

For adult dogs, a single dose should be used. The advice of a veterinarian should be sought regarding the need for and frequency of repeat treatment.

In the event of a heavy roundworm infestation, a repeat dose should be given after 14 days.

#### 4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Benzimidazoles possess wide safety margin. Pyrantel is not absorbed systematically to any extent. Praziquantel also has a wide safety margin, of up to five times the recommended dose.

#### 4.11 Withdrawal Period(s)

Not applicable.

### 5 PHARMACOLOGICAL or IMMUNOLOGICAL PROPERTIES

Pharmacotherapeutic group: Anthelmintics, Benzimidazoles and related substances

ATCvet code: QP52AC55

#### 5.1 Pharmacodynamic properties

The product contains anthelmintics active against roundworms and tapeworms. The product contains three active substances: febantel, pyrantel embonate (pamoate) and praziquantel, a partially hydrogenated pyrazino-isoquinoline derivative used widely as an anthelmintic for both human and veterinary use. Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis and thereby allow removal from the gastro-intestinal (GI) system by peristalsis.

With the mammalian system febantel undergoes ring closure forming fenbendazole and oxfendazole. It is these chemical entities which exert the anthelmintic effect by inhibition of tubulin polymerization. Formation of microtubules is thereby prevented, resulting in disruption to structures vital to the normal functioning of the helminth. Glucose uptake, in particular, is affected, leading to depletion in cell ATP. The parasite dies upon exhaustion of its energy reserves, which occurs 2 – 3 days later.

Praziquantel is very rapidly absorbed and distributed throughout the parasite. Both *in vitro* and *in vivo* studies have shown that praziquantel causes severe damage to the parasite integument, resulting in contraction and paralysis. There is an almost instantaneous tetanic contraction of the parasite musculature and a rapid vacuolisation of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

In this fixed combination product pyrantel and febantel act synergistically against all relevant nematodes (ascarids and hookworms) in dogs. In particular, the activity spectrum covers *Toxocara canis*, *Toxascaris leonina*, *Uncinaria stenocephala* and *Ancylostoma caninum*. The spectrum of activity of praziquantel covers also cestode species in dogs, in particular all *Taenia* spp. and *Dipylidium caninum*. Praziquantel acts against adult and immature forms of these parasites.

#### 5.2 Pharmacokinetic properties

Perorally administered praziquantel is absorbed almost completely from the intestinal tract. After absorption, the drug is distributed to all organs. Praziquantel is metabolized into inactive forms in the liver and secreted in bile. It is excreted within 24 hours to more than 95 % of the administered dosage. Only traces of non-metabolised praziquantel are excreted.

The pamoate salt of pyrantel has low aqueous solubility, an attribute that reduces absorption from the gut and allows the drug to reach and be effective against parasites in the large intestine. Because of the low systemic absorption of pyrantel pamoate, there is very little danger of adverse reactions/toxicity in the host. Following absorption, pyrantel pamoate is quickly and almost completely metabolized into inactive metabolites that are excreted rapidly in the urine.

Febantel is absorbed relatively rapidly and metabolized to a number of metabolites including fenbendazole and oxfendazole, which have anthelmintic activity.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose Monohydrate  
Maize Starch  
Povidone K-30  
Sodium Lauryl Sulfate  
Microcrystalline Cellulose  
Colloidal Anhydrous Silica  
Magnesium Stearate  
Meat Flavour

### **6.2 Incompatibilities**

None known.

### **6.3 Shelf-life**

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and composition of immediate packaging**

OPA/Al/PVC-Al blister.  
Cardboard box containing 2, 4, 10, 30, 50, 100 or 300 tablets.  
Not all pack sizes may be marketed.

### **6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

## **7 MARKETING AUTHORISATION HOLDER**

KRKA d.d, Novo mesto  
Šmarješka cesta 6  
8501 Novo mesto  
Slovenia

## **8 MARKETING AUTHORISATION NUMBER(S)**

VPA 10774/033/001

## **9 DATE OF THE FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 1<sup>st</sup> May 2015

Date of last renewal: 20<sup>th</sup> May 2016

**10 DATE OF REVISION OF THE TEXT**

July 2016