

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Lamoro 200mg Dispersible Tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 200mg of lamotrigine.

Excipient: each tablet contains 15mg of Aspartame (E951).

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Dispersible tablet

Lamoro 200mg Dispersible Tablets are white to off-white, round tablets plain on both sides (approx 12mm in diameter).

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Epilepsy:

Adults and adolescents aged 13 years and above

- Adjunctive or monotherapy treatment of partial seizures and generalised seizures, including tonic-clonic seizures
- Seizures associated with Lennox-Gastaut Syndrome. Lamoro is given as adjunctive therapy but may be the initial antiepileptic drug (AED) to start with in Lennox-Gastaut syndrome.

Children and adolescents aged 2 to 12 years

- Adjunctive treatment of partial seizures and generalised seizures, including tonic-clonic seizures and the seizures associated with Lennox-Gastaut Syndrome.
- Monotherapy of typical absence seizures.

Bipolar disorder

Adults aged 18 years and above

- Prevention of depressive episodes in patients with bipolar I disorder who experience predominantly depressive episodes (see section 5.1).

Lamoro is not indicated for the acute treatment of manic or depressive episodes.

4.2 Posology and method of administration

LAMORO Dispersible Tablets may be chewed, dispersed in a small volume of water (at least enough to cover the whole tablet) or swallowed whole with a little water.

If a calculated dose of lamotrigine (e.g. for use in children (epilepsy only) or patients with hepatic impairment) cannot be divided into multiple lower strength tablets, the dose to be administered is that equal to the nearest lower strength of whole tablets.

Restarting Therapy

Prescribers should assess the need for escalation to maintenance dose when restarting Lamotrigine in patients who have discontinued Lamotrigine for any reason, since the risk of serious rash is associated with high initial doses and exceeding the recommended dose escalation for Lamotrigine (see section 4.4). The greater the interval of time since the previous dose, the more consideration should be given to escalation to the maintenance dose. When the interval since discontinuing Lamotrigine exceeds five half-lives (see section 5.2), Lamotrigine should generally be escalated to the maintenance dose according to the appropriate schedule.

It is recommended that Lamotrigine not be restarted in patients who have discontinued due to rash associated with prior treatment with Lamotrigine unless the potential benefit clearly outweighs the risk.

Epilepsy

The recommended dose escalation and maintenance doses for adults and adolescents aged 13 years and above (Table 1) and for children and adolescents aged 2 to 12 years (Table 2) are given below. Because of a risk of rash the initial dose and subsequent dose escalation should not be exceeded (see section 4.4).

When concomitant AEDs are withdrawn or other AEDs/medicinal products are added on to treatment regimes containing lamotrigine, consideration should be given to the effect this may have on lamotrigine pharmacokinetics (see section 4.5).

Table 1: Recommended treatment regimen in EPILEPSY for adults and children 13 years of age and older

Treatment regimen	Weeks 1 + 2	Weeks 3 + 4	Maintenance Dose
Monotherapy	25 mg/day (once a day)	50 mg/day (once a day)	100 – 200 mg/day (once a day or two divided doses) To achieve maintenance, doses may be increased by 50 – 100 mg every one to two weeks until optimal response is achieved 500 mg/day has been required by some patients to achieve desired response
Add-on therapy with valproate regardless of any concomitant medications (see section 4.5)	12.5 mg/day (given as 25 mg on alternate days)	25 mg/day (once a day)	100 – 200 mg/day (once a day or two divided doses) To achieve maintenance, doses may be increased by maximum of 25 – 50 mg every one to two weeks until optimal response is achieved

Add-on therapy without valproate and with inducers of lamotrigine glucuronidation	This dosage regimen should be used without valproate but with: phenytoin carbamazepine, phenobarbitone, primidone rifampicin lopinavir/ritonavir	50 mg/day (once a day)	100 mg/day (two divided doses)	200 –400 mg/day (two divided doses) To achieve maintenance, doses may be increased by maximum of 100 mg every one to two weeks until optimal response is achieved 700 mg/day has been required by some patients to achieve desired response
Add-on therapy without valproate and without inducers of lamotrigine glucuronidation	This dosage regimen should be used with other medicinal products that do not significantly inhibit or induce lamotrigine glucuronidation	25 mg/day (once a day)	50 mg/day (once a day)	100 – 200 mg/day (once a day or two divided doses) To achieve maintenance, doses may be increased by maximum of 50 – 100 mg every one to two weeks until optimal response is achieved
In patients taking AED’s where the pharmacokinetic interaction with Lamotrigine is currently not known (<i>see Interactions</i>), the treatment regimen as recommended for Lamotrigine with concurrent valproate should be used.				

Table 2: Recommended treatment regimen in EPILEPSY for children aged 2-12 years (Total daily dose in mg/kg bodyweight/day)

Treatment regimen		Weeks 1 + 2	Weeks 3 + 4	Maintenance Dose
Monotherapy of typical absence seizures		0.3 mg/kg/day (once a day or two divided doses)	0.6 mg/kg/day (once a day or two divided doses)	1 – 15 mg/kg/day (once a day or two divided doses) To achieve maintenance, doses may be increased by maximum of 0.6 mg/kg every one to two weeks until optimal response is achieved, with a maximum maintenance dose of 200 mg/day
Add-on therapy with valproate regardless of any other concomitant medication		0.15 mg/kg/day* (once a day)	0.3 mg/kg/day (once a day)	1 – 5 mg/kg/day (once a day or two divided doses) To achieve maintenance, doses may be increased by maximum of 0.3 mg/kg every one to two weeks until optimal response is achieved, with a maximum maintenance dose of 200 mg/day
Add-on therapy without valproate	With inducers of Lamotrigine glucuronidation e.g. phenytoin, carbamazepine,	0.6 mg/kg/day (two divided doses)	1.2 mg/kg/day (two divided doses)	5 – 15 mg/kg/day (once a day or two divided doses) To achieve maintenance, doses may be increased by maximum

	phenobarbitone, primidone Rifampicin, Lopinavir/ritonavir			of 1.2 mg/kg every one to two weeks until optimal response is achieved, with a maximum maintenance dose of 400 mg/day
	Without inducers or inhibitors of Lamotrigine glucuronidation (see section 4.5)	0.3 mg/kg (one or two divided doses)	0.6 mg/kg (one or two divided doses)	1 – 10 mg/kg (maximum of 200 mg/day) (once a day or two divided doses) To achieve maintenance, doses may be increased by maximum of 0.6 mg/kg every one to two weeks until optimal response is achieved, with a maximum maintenance dose of 200 mg/day
In patients taking AED's where the pharmacokinetic interaction with Lamotrigine is currently not known (<i>see Interactions</i>), the treatment regimen as recommended for Lamotrigine with concurrent valproate should be used.				

* NOTE: - If the calculated daily dose in patients taking valproate is 1 to 2 mg, then 2 mg Lamotrigine may be taken on alternate days for the first two weeks. If the calculated daily dose in patients taking valproate is less than 1 mg, then Lamotrigine should not be administered.

To ensure a therapeutic dose is maintained the weight of the child must be monitored and the dose reviewed as weight changes occur. It is likely that patients aged 2-6 years will require a maintenance dose at the higher end of the recommended range.
If epileptic control is achieved with adjunctive treatment, concomitant AEDs may be withdrawn and patients continued on Lamoro monotherapy.

Children below 2 years

There are limited data on the efficacy and safety of lamotrigine for adjunctive therapy of partial seizures in children aged 1 month to 2 years (see section 4.4). There are no data in children below 1 month of age. Thus Lamoro is not recommended for use in children below 2 years of age. If, based on clinical need, a decision to treat is nevertheless taken, see section 4.4, 5.1 and 5.2.

Bipolar disorder

The recommended dose escalation and maintenance doses for adults of 18 years of age and above are given in the tables below. The transition regimen involves escalating the dose of lamotrigine to a maintenance stabilisation dose over six weeks (Table 3) after which other psychotropic medicinal products and/or AEDs can be withdrawn, if clinically indicated (Table 4). The dose adjustments following addition of other psychotropic medicinal products and/or AEDs are also provided below (Table 5). Because of the risk of rash the initial dose and subsequent dose escalation should not be exceeded (see section 4.4).

Table 3: Recommended dose escalation to the maintenance total daily stabilisation dose for adults (over 18 years of age) treated for Bipolar Disorder

Treatment Regimen	Weeks 1-2	Weeks 3-4	Week 5	Target Stabilisation Dose (Week 6)**
a) Monotherapy with Lamotrigine or Adjunct therapy without Valproate and	25 mg/day (once a day)	50 mg/day (once a day or two divided doses)	100 mg/day (once a day or two divided doses)	200 mg/day – usual target dose for optimal response (once a day or two divided doses) Doses in the range 100 - 400 mg/day used in

without inducers of Lamotrigine glucuronidation (see section 4.5)				clinical trials
b) Adjunct therapy with inhibitors of Lamotrigine glucuronidation (e.g. valproate) (see section 4.5)	12.5 mg/day (given 25 mg alternate days)	25 mg/day (once a day)	50 mg/day (once a day or two divided doses)	100 mg/day – usual target dose for optimal response (once a day or two divided doses) Maximum dose of 200mg/day can be used depending on clinical response
c) Adjunct therapy with inducers of Lamotrigine glucuronidation e.g. phenytoin, carbamazepine, phenobarbitone, primidone Rifampicin Lopinavir/ritonavir) in patients NOT taking inhibitors of Lamotrigine glucuronidation (e.g. valproate)	50 mg/day (once a day or two divided doses)	100 mg/day (two divided doses)	200 mg/day (two divided doses)	300 mg/day in week 6, increasing to 400 mg/day if necessary in week 7 to achieve optimal response (two divided doses)
NOTE: In patients taking psychotropic medicines where the pharmacokinetic interaction with Lamotrigine is currently not known, the dose escalation as recommended for Lamotrigine with concurrent valproate, should be used.				

****The Target stabilisation dose will alter depending on clinical response.**

Table 4: Adults aged 18 years and above – maintenance stabilization total daily dose following withdrawal of concomitant medicinal products in treatment of bipolar disorder
Once the target daily maintenance stabilization dose has been achieved, other medicinal products may be withdrawn as shown below.

Treatment Regimen	Current lamotrigine stabilization dose (prior to withdrawal)	Week 1 (beginning with withdrawal)	Week 2	Week 3 Onwards*
Withdrawal of valproate (inhibitor of lamotrigine glucuronidation – see section 4.5), depending on original dose of lamotrigine:				
When valproate is withdrawn, double the stabilization dose, not exceeding an increase of more than 100mg/week	100mg/day	200mg/day	Maintain this dose (200mg/day) (two divided doses)	
	200mg/day	300mg/day	400mg/day	Maintain this dose 400mg/day
Withdrawal of inducers of lamotrigine glucuronidation (see section 4.5), depending on original dose of lamotrigine:				
This dosage regimen should be used when the following are	400mg/day	400mg/day	300mg/day	200mg/day
	300mg/day	300mg/day	225mg/day	150mg/day

withdrawn:				
phenytoin, carbamazepine, phenobarbitone, primidone, rifampicin, lopinavir/ritonavir	200mg/day	200mg/day	150mg/day	100mg/day
Withdrawal of medicinal products that do NOT significantly inhibit or induce lamotrigine glucuronidation (see section 4.5):				
This dosage regimen should be used when other medicinal products that do not significantly inhibit or induce lamotrigine glucuronidation are withdrawn		Maintain target dose achieved in dose escalation (200mg/day; two divided doses) (dose range 100 – 400mg/day)		
In patients taking medicinal products where the pharmacokinetic interaction with lamotrigine is currently not known (see section 4.5), the treatment regimen recommended for lamotrigine is to initially maintain the current dose and adjust the lamotrigine treatment based on clinical response.				

* Dose may be increased to 400 mg/day as needed.

Table 5: Adults aged 18 years and above – adjustment of lamotrigine daily dosing following the addition of other medicinal products in treatment of bipolar disorder

There is no clinical experience in adjusting the lamotrigine daily dose following the addition of other medicinal products. However, based on interaction studies with other medicinal products, the following recommendations can be made:

Treatment Regimen	Current lamotrigine stabilization dose (prior to addition)	Week 1 (beginning with addition)	Week 2	Week 3 Onwards
Addition of valproate (inhibitor of lamotrigine glucuronidation – see section 4.5), depending on original dose of lamotrigine:				
This dosage regimen should be used when valproate is added regardless of any concomitant medicinal products	200mg/day	100mg/day	Maintain this dose (100mg/day)	
	300mg/day	150mg/day	Maintain this dose (150mg/day)	
	400mg/day	200mg/day	Maintain this dose (200mg/day)	
Addition of inducers of lamotrigine glucuronidation in patients NOT taking valproate (see section 4.5), depending on original dose of lamotrigine:				
This dosage regimen should be used when the following are added without valproate: phenytoin, carbamazepine, phenobarbitone, primidone, rifampicin, lopinavir/ritonavir	200mg/day	200mg/day	300mg/day	400mg/day
	150mg/day	150mg/day	225mg/day	300mg/day
	100mg/day	100mg/day	150mg/day	200mg/day
Addition of medicinal products that do NOT significantly inhibit or induce lamotrigine glucuronidation (see section 4.5):				
This dosage regimen should be used when other medicinal products that do not significantly inhibit or induce lamotrigine glucuronidation are		Maintain target dose achieved in dose escalation (200mg/day; dose range 100 – 400mg/day)		

added	
In patients taking medicinal products where the pharmacokinetic interaction with lamotrigine is currently not known (see section 4.5), the treatment regimen as recommended for lamotrigine with concurrent valproate, should be used.	

Discontinuation of Lamotrigine in patients with bipolar disorder:

In clinical trials, there was no increase in the incidence, severity or type of adverse experiences following abrupt termination of Lamotrigine versus placebo. Therefore, patients may terminate Lamotrigine without a step-wise reduction of dose.

Children and adolescents below 18 years

Lamoro is not recommended for use in children below 18 years of age due to a lack of data on safety and efficacy (see section 4.4).

GENERAL DOSING RECOMMENDATIONS FOR LAMOTRIGINE IN SPECIAL PATIENT POPULATIONS

Women taking hormonal contraceptives

The use of an ethinyloestradiol/levonorgestrel (30 µg/150 µg) combination increases the clearance of lamotrigine by approximately two-fold, resulting in decreased lamotrigine levels. Following titration, higher maintenance doses of lamotrigine (by as much as two-fold) may be needed to attain a maximal therapeutic response. During the pill-free week, a two-fold increase in lamotrigine levels has been observed. Dose-related adverse events cannot be excluded. Therefore, consideration should be given to using contraception without a pill-free week, as first-line therapy (for example, continuous hormonal contraceptives or non-hormonal methods; see sections 4.4 and 4.5).

Starting hormonal contraceptives in patients already taking maintenance doses of lamotrigine and NOT taking inducers of lamotrigine glucuronidation

The maintenance dose of lamotrigine will in most cases need to be increased by as much as two-fold (see sections 4.4 and 4.5). It is recommended that from the time that the hormonal contraceptive is started, the lamotrigine dose is increased by 50 to 100 mg/day every week, according to the individual clinical response. Dose increases should not exceed this rate, unless the clinical response supports larger increases. Measurement of serum lamotrigine concentrations before and after starting hormonal contraceptives may be considered, as confirmation that the baseline concentration of lamotrigine is being maintained. If necessary, the dose should be adapted. In women taking a hormonal contraceptive that includes one week of inactive treatment ("pill-free week"), serum lamotrigine level monitoring should be conducted during week 3 of active treatment, i.e. on days 15 to 21 of the pill cycle. Therefore, consideration should be given to using contraception without a pill-free week, as first-line therapy (for example, continuous hormonal contraceptives or non-hormonal methods; see sections 4.4 and 4.5).

Stopping hormonal contraceptives in patients already taking maintenance doses of lamotrigine and NOT taking inducers of lamotrigine glucuronidation

The maintenance dose of lamotrigine will in most cases need to be decreased by as much as 50% (see sections 4.4 and 4.5). It is recommended to gradually decrease the daily dose of lamotrigine by 50-100 mg each week (at a rate not exceeding 25% of the total daily dose per week) over a period of 3 weeks, unless the clinical response indicates otherwise. Measurement of serum lamotrigine concentrations before and after stopping hormonal contraceptives may be considered, as confirmation that the baseline concentration of lamotrigine is being maintained. In women who wish to stop taking a hormonal contraceptive that includes one week of inactive treatment ("pill-free week"), serum lamotrigine level monitoring should be conducted during week 3 of active treatment, i.e. on days 15 to 21 of the pill cycle. Samples for assessment of lamotrigine levels after permanently stopping the contraceptive pill should not be collected during the first week after stopping the pill.

Starting lamotrigine in patients already taking hormonal contraceptives

Dose escalation should follow the normal dose recommendation described in the tables.

Starting and stopping hormonal contraceptives in patients already taking maintenance doses of lamotrigine and TAKING inducers of lamotrigine glucuronidation

Adjustment to the recommended maintenance dose of lamotrigine may not be required.

Use with atazanavir/ritonavir

No adjustments to the recommended dose escalation of lamotrigine should be necessary when lamotrigine is added to the existing atazanavir/ritonavir therapy.

In patients already taking maintenance doses of lamotrigine and not taking glucuronidation inducers, the lamotrigine dose may need to be increased if atazanavir/ritonavir is added, or decreased if atazanavir/ritonavir is discontinued.

Plasma lamotrigine monitoring should be conducted before and during 2 weeks after starting or stopping atazanavir/ritonavir, in order to see if lamotrigine dose adjustment is needed (see section 4.5).

Use with lopinavir/ritonavir

No adjustments to the recommended dose escalation of lamotrigine should be necessary when lamotrigine is added to the existing lopinavir/ritonavir therapy.

In patients already taking maintenance doses of lamotrigine and not taking glucuronidation inducers, the lamotrigine dose may need to be increased if lopinavir/ritonavir is added, or decreased if lopinavir/ritonavir is discontinued. Plasma lamotrigine monitoring should be conducted before and during 2 weeks after starting or stopping lopinavir/ritonavir, in order to see if lamotrigine dose adjustment is needed (see section 4.5).

Elderly (over 65 years of age):-

No dosage adjustment from recommended schedule is required. The pharmacokinetics of Lamotrigine in this age group does not differ significantly from a non-elderly adult population.

Hepatic impairment:-

Initial, escalation and maintenance doses should generally be reduced by approximately 50% in patients with moderate (Child-Pugh grade B) and 75% in severe (Child-Pugh grade C) hepatic impairment. Escalation and maintenance doses should be adjusted according to clinical response (see section 5.2).

Renal impairment

Caution should be exercised when administering Lamotrigine to patients with renal failure. For patients with end-stage renal failure, initial doses of Lamotrigine should be based on patients' concomitant medications; reduced maintenance doses may be effective for patients with significant renal functional impairment. (See 4.4). For more detailed pharmacokinetic information see 5.2..

4.3 Contraindications

Hypersensitivity to the active substance or any of the excipients in section 6.1.

4.4 Special warnings and precautions for useSkin Rash

There have been reports of adverse skin reactions, which have generally occurred within the first eight weeks after initiation of lamotrigine treatment. The majority of rashes are mild and self-limiting, however, serious rashes requiring hospitalization and discontinuation of lamotrigine have also been reported. These have included potentially life-threatening rashes such as Stevens-Johnson syndrome (SJS), and Toxic Epidermal Necrolysis (TEN) and Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) also known as hypersensitivity syndrome (HSS) (see section 4.8).

In adults enrolled in studies utilising the current lamotrigine dosing recommendations, the incidence of serious skin rashes is approximately 1 in 500 in epilepsy patients. Approximately half of these cases have been reported as SJS (1 in 1000). In clinical trials in patients with bipolar disorder, the incidence of serious rash is approximately 1 in 1000.

The risk of serious skin rashes in children is higher than in adults. Available data from a number of studies suggest the incidence of rashes associated with hospitalisation in epileptic children is from 1 in 300 to 1 in 100.

In children, the initial presentation of a rash can be mistaken for an infection, physicians should consider the possibility of a drug reaction in children that develop symptoms of rash and fever during the first eight weeks of therapy.

Additionally, the overall risk of rash appears to be strongly associated with:-

- High initial doses of lamotrigine and exceeding the recommended dose escalation of lamotrigine therapy (see section 4.2).
- Concomitant use of valproate, (see section 4.2).

Caution is also required when treating patients with a history of allergy or rash to other AEDs as the frequency of non-serious rash after treatment with lamotrigine was approximately three times higher in these patients than in those without such history.

All patients (adults and children) who develop a rash should be promptly evaluated and lamotrigine withdrawn immediately unless the rash is clearly not related to lamotrigine treatment. It is recommended that Lamoro not be restarted in patients who have discontinued due to rash associated with prior treatment with lamotrigine unless the potential benefit clearly outweighs the risk. If the patient has developed SJS, TEN or DRESS with the use of lamotrigine, treatment with lamotrigine must not be re-started in this patient at any time.

Rash has also been reported as part of a hypersensitivity syndrome associated with a variable pattern of systemic symptoms including fever, lymphadenopathy, facial oedema and abnormalities of the blood and liver and aseptic meningitis (see section 4.8). The syndrome shows a wide spectrum of clinical severity and may, rarely, lead to disseminated intravascular coagulation and multiorgan failure. It is important to note that early manifestations of hypersensitivity (e.g., fever, lymphadenopathy) may be present even though rash is not evident. If such signs and symptoms are present, the patient should be evaluated immediately and lamotrigine discontinued if an alternative aetiology cannot be established.

Aseptic meningitis was reversible on withdrawal of the drug in most cases, but recurred in a number of cases on re-exposure to lamotrigine. Re-exposure resulted in a rapid return of symptoms that were frequently more severe. Lamotrigine should not be restarted in patients who have discontinued due to aseptic meningitis associated with prior treatment of lamotrigine.

Clinical Worsening and Suicide Risk:

Suicidal ideation and behaviour have been reported in patients treated with antiepileptic agents in several indications. A meta-analysis of randomised placebo controlled trials of anti epileptic drugs has also shown a small increased risk of suicidal ideation and behaviour. The mechanism of this risk is not known and the available data do not exclude the possibility of an increased risk for lamotrigine.

Therefore patients should be monitored for signs of suicidal ideation and behaviours and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of suicidal ideation or behaviour emerge.

In patients with bipolar disorder, worsening of depressive symptoms and/or the emergence of suicidality may occur whether or not they are taking medications for bipolar disorder, including Lamoro. Therefore patients receiving Lamoro for bipolar disorder should be closely monitored for clinical worsening (including development of new symptoms) and suicidality, especially at the beginning of a course of treatment, or at the time of dose changes. Certain patients, such as those with a history of suicidal behaviour or thoughts, young adults, and those patients exhibiting a significant degree of suicidal ideation prior to commencement of treatment, may be at a greater risk of suicidal thoughts or suicide attempts, and should receive careful monitoring during treatment.

Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients who experience clinical worsening (including development of new symptoms) and/or the emergence of suicidal ideation/behaviour, especially if these symptoms are severe, abrupt in onset, or were not part of the patients presenting symptoms.

Hormonal contraceptivesEffects of hormonal contraceptives on Lamotrigine efficacy:

The use of an ethinyloestradiol/levonorgestrel (30 µg / 150 µg) combination increases the clearance of Lamotrigine by approximately two-fold resulting in decreased Lamotrigine levels (*see section 4.5*). A decrease in lamotrigine levels has been associated with loss of seizure control. Following titration, higher maintenance doses of Lamotrigine (by as much as two fold) may be needed to attain a maximal therapeutic response. When stopping hormonal contraceptives, the clearance of lamotrigine may be halved. Increases in lamotrigine concentrations may be associated with dose-related adverse events. Patients should be monitored with respect to this.

In women not already taking an inducer of Lamotrigine glucuronidation and taking a hormonal contraceptive that includes one week of inactive medication (e.g. "pill-free week"), gradual transient increases in Lamotrigine levels will occur during the week of inactive medication. Variations in lamotrigine levels of this order may be associated with adverse effects. Therefore consideration should be given to using contraception without a pill-free week, as first line therapy (for example, continuous hormonal contraceptives or non-hormonal methods).

The interaction between other oral contraceptive or HRT treatments and lamotrigine have not been studied, though they may similarly affect Lamotrigine pharmacokinetic parameters (*see section 4.2*).

Effects of Lamotrigine on hormonal contraceptive efficacy:

An interaction study in 16 healthy volunteers has shown that when Lamotrigine and a hormonal contraceptive (ethinyloestradiol/levonorgestrel combination) are administered in combination, there is a modest increase in levonorgestrel clearance and changes in serum FSH and LH (*see section 4.5*). The impact of these changes on ovarian ovulatory activity is unknown. However, the possibility of these changes resulting in decreased contraceptive efficacy in some patients taking hormonal preparations with Lamotrigine cannot be excluded. Therefore patients should be instructed to promptly report changes in their menstrual pattern, i.e., breakthrough bleeding.

Dihydrofolate Reductase

Lamotrigine is a weak inhibitor of dihydrofolate reductase, hence there is a possibility of interference with folate metabolism during long-term therapy. However, during prolonged human dosing, Lamotrigine did not induce significant changes in the haemoglobin concentration, mean corpuscular volume, or serum or red blood cell folate concentrations up to 1 year or red blood cell folate concentrations for up to 5 years.

Renal Failure

In single dose studies in subjects with end stage renal failure, plasma concentrations of Lamotrigine were not significantly altered. However, accumulation of the glucuronide metabolite is to be expected; caution should therefore be exercised in treating patients with renal failure.

Patients taking other preparations containing Lamotrigine

Lamoro should not be administered to patients currently being treated with any other preparation containing Lamotrigine without consulting a doctor.

Development in children

There are no data on the effect of lamotrigine on growth, sexual maturation and cognitive, emotional and behavioural developments in children.

Precautions relating to epilepsy:

As with other AEDs, abrupt withdrawal of Lamotrigine may provoke rebound seizures. Unless safety concerns (for example rash) require an abrupt withdrawal, the dose of Lamoro should be gradually decreased over a period of 2 weeks.

There are reports in the literature that severe convulsive seizures including status epilepticus may lead to rhabdomyolysis, multiorgan dysfunction and disseminated intravascular coagulation, sometimes with fatal outcome. Similar cases have occurred in association with the use of Lamotrigine.

A clinically significant worsening of seizure frequency instead of an improvement may be observed. In patients with more than one seizure type, the observed benefit of control for one seizure type should be weighed against any

observed worsening in another seizure type.

Myoclonic seizures may be worsened by lamotrigine.

There is a suggestion in the data that responses in combination with enzyme inducers is less than in combination with non-enzyme inducing antiepileptic agents. The reason is unclear.

In children taking lamotrigine for the treatment of typical absence seizures, efficacy may not be maintained in all patients.

Precautions relating to Bipolar Disorder:

Children and adolescents below 18 years

Treatment with antidepressants is associated with an increased risk of suicidal thinking and behavior in children and adolescents with major depressive disorder and other psychiatric disorders.

4.5 Interaction with other medicinal products and other forms of interaction

Interaction studies have only been performed in adults.

UDP-glucuronyl transferases have been identified as the enzymes responsible for metabolism of Lamotrigine. There is no evidence that Lamotrigine causes clinically significant induction or inhibition of hepatic oxidative drug-metabolising enzymes, and interactions between Lamotrigine and drugs metabolised by cytochrome P450 enzymes are unlikely to occur. Lamotrigine may induce its own metabolism but the effect is modest and unlikely to have significant clinical consequences.

Table 6: Effects of other drugs on glucuronidation of Lamotrigine

Medicinal products that significantly inhibit glucuronidation of Lamotrigine	Medicinal products that significantly induce glucuronidation of Lamotrigine	Medicinal products that do not significantly inhibit or induce glucuronidation of Lamotrigine
Valproate	Carbamazepine Phenytoin Primidone Phenobarbitone Rifampicin Lopinavir/ritonavir Atazanavir/ritonavir* Ethinylloestradiol/ levonorgestrel combination**	Lithium Bupropion Olanzapine Oxcarbazepine Felbamate Gabapentin Levetiracetam Pregabalin Topiramate Zonisamide Aripiprazole

*For dosing guidance (see section 4.2).

** Other oral contraceptive and HRT treatments have not been studied, though they may similarly affect Lamotrigine pharmacokinetic parameters (see sections 4.2 and 4.4).

Interactions involving antiepileptic drugs

Valproate, which inhibits the glucuronidation of Lamotrigine, reduces the metabolism of Lamotrigine and increases the mean half life of Lamotrigine nearly two fold. In patients receiving concomitant therapy with valproate, the appropriate treatment regimen should be used (see section 4.2).

Certain antiepileptic agents (such as phenytoin, carbamazepine, phenobarbitone and primidone) which induce hepatic drug-metabolising enzymes induce the glucuronidation of Lamotrigine and enhance the metabolism of Lamotrigine. In patients receiving concomitant therapy with phenytoin, carbamazepine, pheonbarbitone or primidone, the appropriate treatment regimen should be used (see section 4.2).

There have been reports of central nervous system events including dizziness, ataxia, diplopia, blurred vision and

nausea in patients taking carbamazepine following the introduction of Lamotrigine.

These events usually resolve when the dose of carbamazepine is reduced. A similar effect was seen during a study of Lamotrigine and oxcarbazepine in healthy study volunteers, but dose reduction was not investigated.

There are reports in the literature of decreased lamotrigine levels when lamotrigine was given in combination with oxcarbazepine. However in a prospective study in healthy adult volunteers using doses of 200mg lamotrigine and 1200mg oxcarbazepine, oxcarbazepine did not alter the metabolism of lamotrigine and lamotrigine did not alter the metabolism of oxcarbazepine. Therefore in patients receiving concomitant therapy with oxcarbazepine, the treatment regimen for lamotrigine adjunctive therapy without valproate and without inducers of lamotrigine glucuronidation should be used (see section 4.2).

In a study of healthy volunteers, co-administration of felbamate (1200mg twice daily) with lamotrigine (100mg twice daily for 10 days) appeared to have no clinically relevant effects on the pharmacokinetics of lamotrigine.

Based on a retrospective analysis of plasma levels in patients who received lamotrigine both with and without gabapentin, gabapentin does not appear to change the apparent clearance of lamotrigine.

Potential interactions between levetiracetam and lamotrigine were assessed by evaluating serum concentrations of both agents during placebo-controlled clinical trials. These data indicate that lamotrigine does not influence the pharmacokinetics of levetiracetam and that levetiracetam does not influence the pharmacokinetics of lamotrigine.

Steady-state trough plasma concentrations of lamotrigine were not affected by concomitant pregabalin (200mg, 3 times daily) administration. There are no pharmacokinetic interactions between lamotrigine and pregabalin.

Topiramate resulted in no change in plasma concentrations of lamotrigine. Administration of lamotrigine resulted in a 15% increase in topiramate concentrations.

In a study of patients with epilepsy, coadministration of zonisamide (200mg to 400mg/day) with lamotrigine (150 to 500mg/day) for 35 days had no significant effect on the pharmacokinetics of lamotrigine.

Although changes in the plasma concentrations of other anti-epileptic drugs have been reported, controlled studies have shown no evidence that Lamotrigine affects the plasma concentrations of concomitant anti-epileptic drugs. Evidence from *in vitro* studies indicates that Lamotrigine does not displace other anti-epileptic drugs from protein binding sites.

Interactions involving other psychoactive agents

The pharmacokinetics of lithium after 2g of anhydrous lithium gluconate given twice daily for six days to 20 healthy subjects were not altered by co-administration of 100 mg/day Lamotrigine.

Multiple oral doses of bupropion had no statistically significant effects on the single dose pharmacokinetics of Lamotrigine in 12 subjects and had only a slight increase in the AUC of Lamotrigine glucuronide.

In a study in healthy adult volunteers, 15 mg olanzapine reduced the AUC and C_{\max} of Lamotrigine by an average of 24% and 20%, respectively. An effect of this magnitude is not generally expected to be clinically relevant. Lamotrigine at 200 mg did not affect the pharmacokinetics of olanzapine.

Multiple oral doses of lamotrigine 400mg daily had no clinically significant effect on the single dose pharmacokinetics of 2mg risperidone in 14 healthy adult volunteers. Following the co-administration of risperidone 2mg with lamotrigine, 12 out of the 14 volunteers reported somnolence compared to 1 out of 20 when risperidone was given alone, and none when lamotrigine was administered alone.

In a study of 18 adult patients with bipolar I disorder, receiving an established regimen of lamotrigine (100-400 mg/day), doses of aripiprazole were increased from 10 mg/day to a target of 30 mg/day over a 7 day period and continued once daily for a further 7 days. An average reduction of approximately 10% in C_{\max} and AUC of lamotrigine was observed. An effect of this magnitude is not expected to be of clinical consequence.

In vitro inhibition experiments indicated that the formation of Lamotrigine's primary metabolite, the 2-N-glucuronide, was minimally affected by co-incubation with amitriptyline, bupropion, clonazepam, haloperidol, or lorazepam.

These experiments also suggested that metabolism of lamotrigine was unlikely to be inhibited by clozapine, fluoxetine, phenelzine, risperidone, sertraline or trazodone. In addition, a study of bufuralol metabolism using human liver microsome preparations suggested that lamotrigine would not reduce the clearance of medicinal products metabolized predominantly by CYP2D6.

Interactions involving hormonal contraceptives

Effect of hormonal contraceptives on Lamotrigine pharmacokinetics

In a study of 16 female volunteers, dosing with 30 µg ethinylloestradiol/150 µg levonorgestrel in a combined oral contraceptive pill caused an approximately two-fold increase in Lamotrigine oral clearance, resulting in an average 52% and 39% reduction in Lamotrigine AUC and C_{\max} , respectively. Serum Lamotrigine concentrations gradually increased during the course of the week of inactive medication (e.g. "pill-free" week), with pre-dose concentrations at the end of the week of inactive medication being, on average, approximately two-fold higher than during co-therapy (see section 4.4). No adjustments to the recommended dose escalation guidelines for lamotrigine should be necessary solely based on the use of hormonal contraceptives, but the maintenance dose of lamotrigine will need to be increased or decreased in most cases when starting or stopping hormonal contraceptives (see section 4.2).

Effect of Lamotrigine on hormonal contraceptive pharmacokinetics

In a study of 16 female volunteers, a steady state dose of 300 mg Lamotrigine had no effect on the pharmacokinetics of the ethinylloestradiol component of a combined oral contraceptive pill. A modest increase in oral clearance of the levonorgestrel component was observed, resulting in an average 19% and 12% reduction in levonorgestrel AUC and C_{\max} , respectively. Measurement of serum FSH, LH and oestradiol during the study indicated some loss of suppression of ovarian hormonal activity in some women, although measurement of serum progesterone indicated that there was no hormonal evidence of ovulation in any of the 16 subjects. The impact of the modest increase in levonorgestrel clearance, and the changes in serum FSH and LH, on ovarian ovulatory activity is unknown (*see 4.4 Warnings and Precautions*). The effects of doses of Lamotrigine other than 300 mg/day have not been studied and studies with other female hormonal preparations have not been conducted.

Interactions involving other medicinal products

In a study in 10 male volunteers, rifampicin increased lamotrigine clearance and decreased lamotrigine half-life due to induction of the hepatic enzymes responsible for glucuronidation. In patients receiving concomitant therapy with rifampicin, the appropriate treatment regimen should be used (see section 4.2).

In a study in healthy volunteers, lopinavir/ritonavir approximately halved the plasma concentrations of lamotrigine, probably by induction of glucuronidation. In patients receiving concomitant therapy with lopinavir/ritonavir, the appropriate treatment regimen should be used (see section 4.2).

In a study in healthy adult volunteers, atazanavir/ritonavir (300mg/100mg) administered for 9 days reduced the plasma AUC and C_{\max} of lamotrigine (single 100mg dose) by an average of 32% and 6%, respectively. In patients receiving concomitant therapy with atazanavir/ritonavir, the appropriate treatment regimen should be used (see section 4.2). Data from *in vitro* assessment demonstrate that lamotrigine, but not the N(2)-glucuronide metabolite, is an inhibitor of Organic Transporter 2 (OCT 2) at potentially clinically relevant concentrations. These data demonstrate that lamotrigine is an inhibitor of OCT 2, with IC_{50} value of 53.8 µM. Co-administration of lamotrigine with renally excreted medicinal products which are substrates of OCT2 (e.g. metformin, gabapentin and varenicline) may result in increased plasma levels of these drugs.

The clinical significance of this has not been clearly defined, however care should be taken in patients co-administered with these medicinal products.

4.6 Fertility, pregnancy and lactation

Risk related to antiepileptic drugs in general

Specialist advice should be given to women who are of childbearing potential. The need for treatment with AEDs should be reviewed when a woman is planning to become pregnant. In women being treated for epilepsy, sudden discontinuation of AED therapy should be avoided as this may lead to breakthrough seizures that could have serious consequences for the woman and the unborn child.

Monotherapy should be preferred whenever possible because therapy with multiple AEDs could be associated with a higher risk of congenital malformations than monotherapy, depending on the associated antiepileptics.

Risk related to lamotrigine

Pregnancy:

A large amount of data on pregnant women exposed to Lamotrigine monotherapy during the first trimester of pregnancy (more than 8700) do not suggest a substantial increase in the risk for major congenital malformations including oral clefts. Animal studies have shown developmental toxicity (see section 5.3).
If therapy with lamoro is considered necessary during pregnancy, the lowest possible therapeutic dose is recommended.

Lamotrigine has a slight inhibitory effect on dihydrofolic acid reductase and could therefore theoretically lead to an increased risk of embryofoetal damage by reducing folic acid levels (see section 4.4). Intake of folic acid when planning pregnancy and during early pregnancy may be considered.

Physiological changes during pregnancy may affect Lamotrigine levels and/or therapeutic effect. There have been reports of decreased lamotrigine plasma levels during pregnancy with a potential risk of loss of seizure control. After birth lamotrigine levels may increase rapidly with a risk of dose-related adverse events. Therefore lamotrigine serum concentrations should be monitored before, during and after pregnancy, as well as shortly after birth. If necessary, the dose should be adapted to maintain the lamotrigine serum concentration at the same level as before pregnancy, or adapted according to clinical response. In addition, dose-related undesirable effects should be monitored after birth.

Lactation:

Lamotrigine has been reported to pass into breast milk in highly variable concentrations, resulting in total lamotrigine levels in infants of up to approximately 50% of the mothers. Therefore in some breast-fed infants, serum concentrations of lamotrigine may reach levels at which pharmacological effects occur. Among a limited group of exposed infants, no adverse effects were observed.
The potential benefits of breast-feeding should be weighed against the risk of adverse effects occurring in the infant. Individual risk/benefit assessment is important, and close monitoring of both mother and infant is advisable, especially during the first 3 weeks postpartum.

Fertility:

Animal experiments did not reveal impairment of fertility by lamotrigine (see section 5.3).

4.7 Effects on ability to drive and use machines

As there is individual variation in response to all anti-epileptic drug therapy, patients taking Lamoro to treat epilepsy should consult their physician on the specific issues of driving and epilepsy.

No studies on the effects on the ability to drive and use machines have been performed.
Two volunteer studies have demonstrated that the effect of Lamotrigine on fine visual motor co-ordination, eye movements, body sway and subjective sedative effects did not differ from placebo. In clinical trials with Lamotrigine adverse events of a neurological character such as dizziness and diplopia have been reported. Therefore, patients should see how Lamotrigine therapy affects them before driving or operating machinery.

4.8 Undesirable effects

The undesirable effects for epilepsy and bipolar disorder indications are based on the data currently available. The table below should be consulted when considering the overall safety profile of lamotrigine.

Adverse reactions identified from epilepsy monotherapy clinical trials (identified by a dagger †), from bipolar disorder clinical trials (identified by a section mark §) and during other clinical experience are listed in the table below by their incidence in clinical trials.

The following convention has been utilised for the classification of undesirable effects:- Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data).

System Organ Class	Adverse Event	Frequency
Blood and lymphatic system disorders	Haematological abnormalities ¹ including neutropenia, leucopenia, anaemia, thrombocytopenia, pancytopenia, aplastic anaemia, agranulocytosis Lymphadenopathy ¹	Very rare Not known
Immune System Disorders	Hypersensitivity syndrome ² (including such symptoms as, fever, lymphadenopathy, facial oedema, abnormalities of the blood and liver, disseminated intravascular coagulation, multi organ failure).	Very rare
Psychiatric Disorders	Aggression, irritability Confusion, hallucinations, tics Nightmares	Common Very rare Not known
Nervous System Disorders	Headache ^{†§} Somnolence ^{†§} , dizziness ^{†§} , tremor [†] , insomnia [†] , agitation [§] Ataxia [†] Nystagmus [†] Unsteadiness, movement disorders, worsening of Parkinson's disease ³ , extrapyramidal effects, choreoathetosis [†] , increase in seizure frequency Aseptic meningitis (see section 4.4)	Very common Common Uncommon Rare Very rare Not known
Eye disorders	Diplopia [†] , blurred vision [†] Conjunctivitis	Uncommon Rare
Gastrointestinal disorders	Nausea [†] , vomiting [†] , diarrhoea [†] , dry mouth [§]	Common
Hepatobiliary disorders	Hepatic failure, hepatic dysfunction ⁴ , increased liver function tests	Very rare
Skin and subcutaneous tissue disorders	Skin rash ^{5§} Alopecia Stevens–Johnson Syndrome § Toxic epidermal necrolysis DRESS	Very common Uncommon Rare Very rare Not known
Musculoskeletal and connective tissue disorders	Arthralgia [§] Lupus-like reactions	Common Very rare
General disorders and administration site conditions	Tiredness, pain [§] , back pain [§]	Common

Description of selected adverse reactions

1. Haematological abnormalities and lymphadenopathy may or may not be associated with the hypersensitivity syndrome (see Immune system disorders²).
2. Rash has also been reported as part of a hypersensitivity syndrome associated with a variable pattern of systemic symptoms including fever, lymphadenopathy, facial oedema and abnormalities of the blood and liver. The syndrome shows a wide spectrum of clinical severity and may, rarely, lead to disseminated intravascular coagulation and multiorgan failure. It is important to note that early manifestations of hypersensitivity (for example fever, lymphadenopathy) may be present even though rash is not evident. If such signs and symptoms are present, the patient should be evaluated immediately and Lamoro discontinued if an alternative aetiology cannot be established.
3. These effects have been reported during other clinical experience. There have been reports that lamotrigine may worsen parkinsonian symptoms in patients with pre-existing Parkinson's disease, and isolated reports of extrapyramidal effects and choreoathetosis in patients without this underlying condition.
4. Hepatic dysfunction usually occurs in association with hypersensitivity reactions but isolated cases have been reported without overt signs of hypersensitivity.
5. In clinical trials in adults, skin rashes occurred in up to 8-12% of patients taking lamotrigine and in 5-6% of patients taking placebo. The skin rashes led to the withdrawal of lamotrigine treatment in 2% of patients. The rash, usually maculopapular in appearance, generally appears within eight weeks of starting treatment and resolves on withdrawal of Lamoro (see section 4.4). Serious potentially life-threatening skin rashes, including Stevens–Johnson syndrome and toxic epidermal necrolysis (Lyell's Syndrome) have been reported. Although the majority recover on withdrawal of lamotrigine treatment, some patients experience irreversible scarring and there have been rare cases of associated death (see section 4.4).

The overall risk of rash, appears to be strongly associated with:

- high initial doses of lamotrigine and exceeding the recommended dose escalation of lamotrigine therapy (see section 4.2)
- concomitant use of valproate (see section 4.2).

Rash has also been reported as part of a hypersensitivity syndrome associated with a variable pattern of systemic symptoms (see Immune system disorders).

There have been reports of decreased bone mineral density, osteopenia, osteoporosis and fractures in patients on longterm therapy with lamotrigine. The mechanism by which lamotrigine affects bone metabolism has not been identified.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, Earlsfort Terrace, IRL - Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; e-mail: medsafety@hpra.ie.

4.9 Overdose*Symptoms and signs:-*

Acute ingestion of doses in excess of 10-20 times the maximum therapeutic dose has been reported. Overdose has resulted in symptoms including nystagmus, ataxia, impaired consciousness and coma. QRS broadening (intraventricular conduction delay) has also been observed in overdose patients. Broadening of QRS duration to more than 100 msec may be associated with more severe toxicity.

Treatment:-

In the event of overdosage, the patient should be admitted to hospital and given appropriate supportive therapy. Therapy aimed at decreasing absorption (activated charcoal) should be performed if indicated. Further management should be as clinically indicated. There is no experience with haemodialysis as treatment of overdose. In six volunteers with kidney failure, 20% of the lamotrigine was removed from the body during a 4-hour haemodialysis session (see

section 5.2).

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

ATC Code: N03 AX09

Mechanism of action:-

The results of pharmacological studies suggest that Lamotrigine is a use- and voltage-dependent blocker of voltage gated sodium channels. It inhibits sustained repetitive firing of neurons and inhibits release of glutamate (the neurotransmitter which plays a key role in the generation of epileptic seizures). These effects are likely to contribute to the anticonvulsant properties of lamotrigine.

In contrast, the mechanisms by which lamotrigine exerts its therapeutic action in bipolar disorder have not been established, although interaction with voltage gated sodium channels is likely to be important.

Pharmacodynamic effects:

In tests designed to evaluate the central nervous system effects of medicinal products, the results obtained using doses of 240 mg Lamotrigine administered to healthy volunteers did not differ from placebo, whereas both 1000 mg phenytoin and 10 mg diazepam each significantly impaired fine visual motor co-ordination and eye movements, increased body sway and produced subjective sedative effects.

In another study, single oral doses of 600 mg carbamazepine significantly impaired fine visual motor co-ordination and eye movements, while increasing both body sway and heart rate, whereas results with Lamotrigine at doses of 150 mg and 300 mg did not differ from placebo.

Clinical efficacy and safety in children aged 1 to 24 months

The efficacy and safety of adjunctive therapy in partial seizures in patients aged 1 to 24 months has been evaluated in a small double-blind placebo-controlled withdrawal study. Treatment was initiated in 177 subjects, with a dose titration schedule similar to that of children aged 2 to 12 years. Lamotrigine 2 mg tablets are the lowest strength available, therefore the standard dosing schedule was adapted in some cases during the titration phase (for example, by administering a 2 mg tablet on alternate days when the calculated dose was less than 2 mg). Serum levels were measured at the end of week 2 of titration and the subsequent dose either reduced or not increased if the concentration exceeded 0.41 µg/mL, the expected concentration in adults at this time point. Dose reductions of up to 90% were required in some patients at the end of week 2. Thirty-eight responders (> 40% decrease in seizure frequency) were randomised to placebo or continuation of lamotrigine. The proportion of subjects with treatment failure was 84% (16/19 subjects) in the placebo arm and 58% (11/19 subjects) in the lamotrigine arm. The difference was not statistically significant: 26.3%, CI95% -2.6% < 50.2%, p=0.07.

A total of 256 subjects between 1 to 24 months of age have been exposed to lamotrigine in the dose range 1 to 15 mg/kg/day for up to 72 weeks. The safety profile of lamotrigine in children aged 1 month to 2 years was similar to that in older children except that clinically significant worsening of seizures (>=50%) was reported more often in children under 2 years of age (26%) as compared to older children (14%).

Clinical efficacy and safety in Lennox-Gastaut syndrome

There are no data for monotherapy in seizures associated with Lennox-Gastaut syndrome.

Clinical efficacy in the prevention of mood episodes in patients with bipolar disorder

The efficacy of lamotrigine in the prevention of mood episodes in patients with bipolar I disorder has been evaluated in two studies.

Study SCAB2003 was a multicentre, double-blind, double dummy, placebo and lithium-controlled, randomised fixed dose evaluation of the long-term prevention of relapse and recurrence of depression and/or mania in patients with bipolar I disorder who had recently or were currently experiencing a major depressive episode. Once stabilised using Lamotrigine monotherapy or adjunctive therapy, patients were randomly assigned into one of five treatment groups:

Lamotrigine (50, 200, 400 mg/day), lithium (serum levels of 0.8 to 1.1 mMol/L) or placebo for a maximum of 76 weeks (18 months). The primary endpoint was "Time to Intervention for a Mood Episode (TIME)", where the interventions were additional pharmacotherapy or electroconvulsive therapy (ECT). Study SCAB2006 had a similar design as study SCAB2003, but differed from study SCAB2003 in evaluating a flexible dose of lamotrigine (100 to 400 mg/day) and including patients with bipolar I disorder who had recently or were currently experiencing a manic episode. The results are shown in Table 7.

Table 7: Summary of results from studies investigating the efficacy of lamotrigine in the prevention of mood episodes in patients with bipolar I disorder

‘Proportion’ of patients being event free at week 76						
	Study SCAB2003 Bipolar I			Study SCAB2006 Bipolar I		
Inclusion criterion	Major depressive episode			Major manic episode		
	Lamotrigine	Lithium	Placebo	Lamotrigine	Lithium	Placebo
Intervention free	0.22	0.21	0.12	0.17	0.24	0.04
p-value Log rank test	0.004	0.006		0.023	0.006	
Depression free	0.51	0.46	0.41	0.82	0.71	0.40
p-value Log rank test	0.047	0.209		0.015	0.167	
Free of mania	0.70	0.86	0.67	0.53	0.64	0.37
p-value Log rank test	0.339	0.026		0.280	0.006	

In supportive analyses of time to first depressive episode and time to first manic/hypomanic or mixed episode, the Lamotrigine-treated patients had significantly longer times to first depressive episode than placebo patients and the treatment difference with respect to time to manic/hypomanic or mixed episodes was not statistically significant.

The efficacy of lamotrigine in combination with mood stabilizers has not been adequately studied.

Children (10-12 years of age) and Adolescents (13-17 years of age)

A multicentre, parallel group, placebo-controlled, double-blind, randomised withdrawal study, evaluated the efficacy and safety of lamotrigine IR as add-on maintenance therapy to delay mood episodes in male and female children and adolescents (age 10-17 years) who had been diagnosed with bipolar I disorder and who had remitted or improved from a bipolar episode while treated with lamotrigine in combinations with concomitant antipsychotic or other mood-stabilising drugs. The result of the primary efficacy analysis (time to occurrence of a bipolar event – TOBE) did not reach statistical significance (p=0.0717), thus efficacy was not shown. In addition, safety results showed increased reporting of suicidal behaviours in lamotrigine treated patients: 5% (4 patients) in the lamotrigine arm compared to 0 in placebo (see section 4.2).

Study of the effect of lamotrigine on cardiac conduction

A study in healthy adult volunteers evaluated the effect of repeat doses of lamotrigine (up to 400 mg/day) on cardiac conduction, as assessed by 12-lead ECG. There was no clinically significant effect of lamotrigine on QT interval compared to placebo.

5.2 Pharmacokinetic properties

Absorption:-

Lamotrigine is rapidly and completely absorbed from the gut with no significant first pass metabolism. Peak plasma concentrations occur approximately 2.5 hours after oral drug administration. Time to maximum concentration is slightly delayed after food but the extent of absorption is unaffected. The pharmacokinetics are linear up to 450 mg, the highest single dose tested. There is considerable inter-individual variation in steady state maximum concentrations but within an individual concentrations rarely vary.

Distribution:-

Binding to plasma proteins is about 55%; it is very unlikely that displacement from plasma proteins would result in toxicity.

The volume of distribution is 0.92 to 1.22 L/kg.

Metabolism:-

UDP-glucuronyl transferases have been identified as the enzymes responsible for metabolism of Lamotrigine. Lamotrigine induces its own metabolism to a modest extent depending on dose. However, there is no evidence that Lamotrigine affects the pharmacokinetics of other AEDs and data suggest that interactions between Lamotrigine and medicinal products metabolised by cytochrome P₄₅₀ enzymes are unlikely to occur.

Elimination:-

The apparent plasma clearance in healthy subjects is approximately 30 mL/min. Clearance of Lamotrigine is primarily metabolic with subsequent elimination of glucuronide-conjugated material in urine. Less than 10% is excreted unchanged in the urine. Only about 2% of lamotrigine-related material is excreted in faeces. Clearance and half-life are independent of dose. The apparent plasma half-life in healthy subjects is estimated to be approximately 33 hours (range 14 to 103 hours). In a study of subjects with Gilbert's Syndrome, mean apparent clearance was reduced by 32% compared with normal controls but the values are within the range for the general population.

The half-life of Lamotrigine is greatly affected by concomitant medicinal products.

Mean half-life is reduced to approximately 14 hours when given with **glucuronidation** inducing medicinal products such as carbamazepine and phenytoin and is increased to a mean of approximately 70 hours when co-administered with valproate alone (see 4.2 *Posology and Method of Administration and Interactions*).

Linearity:-

The pharmacokinetics of Lamotrigine are linear up to 450 mg, the highest single dose tested.

Children:-

Clearance adjusted for bodyweight is higher in children than in adults with the highest values in children under five years. The half-life of Lamotrigine is generally shorter in children than in adults with a mean value of approximately 7 hours when given with enzyme-inducing medicinal products such as carbamazepine and phenytoin and increasing to mean values of 45 to 50 hours when co-administered with valproate alone (see 4.2 *Posology and Method of Administration*).

Infants aged 2 to 26 months

In 143 paediatric patients aged 2 to 26 months, weighing 3 to 16 kg, clearance was reduced compared to older children with the same body weight, receiving similar oral doses per kg body weight as children older than 2 years. The mean half-life was estimated at 23 hours in infants younger than 26 months on enzyme-inducing therapy, 136 hours when co-administered with valproate and 38 hours in subjects treated without enzyme inducers/inhibitors. The inter-individual variability for oral clearance was high in the group of paediatric patients of 2 to 26 months (47%). The predicted serum concentration levels in children of 2 to 26 months were in general in the same range as those in older children, though higher C_{max} levels are likely to be observed in some children with a body weight below 10 kg.

Elderly:-

Results of a population pharmacokinetic analysis including both young and elderly patients with epilepsy, enrolled in the same trials, indicated that the clearance of Lamotrigine did not change to a clinically relevant extent. After single doses apparent clearance decreased by 12% from 35mL/min at age 20 to 31mL/min at 70 years.

The decrease after 48 weeks of treatment was 10% from 41 to 37 mL/min between the young and elderly groups. In addition, pharmacokinetics of Lamotrigine was studied in 12 healthy elderly subjects following a 150 mg single dose. The mean clearance in the elderly (0.39 mL/min/kg) lies within the range of the mean clearance values (0.31 to 0.65 mL/min/kg) obtained in 9 studies with non-elderly adults after single doses of 30 to 450 mg.

Patients with renal impairment:-

Twelve volunteers with chronic renal failure, and another six individuals undergoing hemodialysis were each given a single 100 mg dose of lamotrigine. Mean clearances were 0.42 mL/min/kg (chronic renal failure), 0.33 mL/min/kg (between hemodialysis) and 1.57 mL/min/kg (during hemodialysis), compared with 0.58 mL/min/kg in healthy volunteers. Mean plasma half-lives were 42.9 hours (chronic renal failure), 57.4 hours (between hemodialysis) and 13.0 hours (during hemodialysis), compared with 26.2 hours in healthy volunteers. On average, approximately 20% (range = 5.6 to 35.1) of the amount of lamotrigine present in the body was eliminated during a 4-hour hemodialysis session. For this patient population, initial doses of lamotrigine should be based on the patient's concomitant medicinal products; reduced maintenance doses may be effective for patients with significant renal functional impairment (*see sections 4.2 and 4.4*).

Patients with hepatic impairment:-

A single-dose pharmacokinetic study was performed in 24 subjects with various degrees of hepatic impairment and 12 healthy subjects as controls. The median apparent clearance of Lamotrigine was 0.31, 0.24 or 0.10 mL/min/kg in patients with Grade A, B, or C (Child - Pugh Classification) hepatic impairment, respectively, compared to 0.34 mL/min/kg in the healthy controls. Initial, escalation, and maintenance doses should generally be reduced in patients with moderate or severe hepatic impairment (*see section 4.2*).

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on studies of safety pharmacology, repeated dose toxicity, genotoxicity and carcinogenic potential.

In reproductive and developmental toxicity studies in rodents and rabbits, no teratogenic effects but reduced foetal weight and retarded skeletal ossification were observed, at exposure levels below or similar to the expected clinical exposure. Since higher exposure levels could not be tested in animals due to the severity of maternal toxicity, the teratogenic potential of lamotrigine has not been characterised above clinical exposure.

In rats, enhanced foetal as well as post-natal mortality was observed when lamotrigine was administered during late gestation and through the early post-natal period. These effects were observed at the expected clinical exposure.

In juvenile rats, an effect on learning in the Biel maze test, a slight delay in balanopreputial separation and vaginal patency and a decreased postnatal body weight gain in F1 animals were observed at exposures approximately two-times higher than the therapeutic exposures in human adults.

Animal experiments did not reveal impairment of fertility by lamotrigine. Lamotrigine reduced foetal folic acid levels in rats. Folic acid deficiency is assumed to be associated with an enhanced risk of congenital malformations in animals as well as in humans.

Lamotrigine caused a dose-related inhibition of the hERG channel tail current in human embryonic kidney cells. The IC₅₀ was approximately nine-times above the maximum therapeutic free concentration. Lamotrigine did not cause QT prolongation in animals at exposures up to approximately two-times the maximum therapeutic free concentration. In a clinical study, there was no clinically significant effect of lamotrigine on QT interval in healthy adult volunteers (*see section 5.1*).

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Crospovidone
Povidone K-30
Mannitol (E421)
Microcrystalline cellulose, silicified
Aspartame (E951)
Croscarmellose sodium
Magnesium stearate
Blackcurrant flavour

6.2 Incompatibilities

Not applicable

6.3 Shelf life

30 months

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

Blister strips made of clear PVC/ PVdC/ Aluminium foil

Pack sizes: 7, 14, 28, 30, 56 & 60 tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

No special requirements.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7 MARKETING AUTHORISATION HOLDER

Pinewood Laboratories Limited
Trading as Pinewood Healthcare
Ballymacarbry
Clonmel
Co. Tipperary

8 MARKETING AUTHORISATION NUMBER

PA0281/128/005

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 16th February 2007

Date of last renewal: 16th February 2012

10 DATE OF REVISION OF THE TEXT

January 2017