

Summary of Product Characteristics

1 NAME OF THE MEDICINAL PRODUCT

Fydrane 0.2 mg/ml + 3.1 mg/ml + 10 mg/ml, solution for injection

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

1 ml of solution for injection contains 0.2 mg of tropicamide, 3.1 mg of phenylephrine hydrochloride and 10 mg of lidocaine hydrochloride.

One dose of 0.2 ml solution contains 0.04 mg of tropicamide, 0.62 mg of phenylephrine hydrochloride and 2 mg of lidocaine hydrochloride.

Excipient with a known effect: sodium (0.59 mg per dose; see section 4.4).

For the full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Solution for injection.

Clear and slightly brownish-yellow solution, practically free from visible particles.

pH: 6.9 - 7.5

Osmolality: 290 – 350 mosmol/kg

4 CLINICAL PARTICULARS

4.1 Therapeutic Indications

Fydrane is indicated for cataract surgery to obtain mydriasis and intraocular anaesthesia, during the surgical procedure.

Fydrane is indicated in adults only.

4.2 Posology and method of administration

Intracameral use. One ampoule for single eye use.

Fydrane must be administered by an ophthalmic surgeon.

Posology

Fydrane should only be used in patients who have demonstrated, at a previous visit, a satisfactory pupil dilation with topical mydriatic therapy.

Adults:

Slowly inject, by intracameral route, 0.2 ml of Fydrane in only one injection, at the start of the surgical procedure.

Special population

Elderly:

No dose adjustment is necessary.

Paediatric population:

The safety and efficacy of Fydrane in children aged 0 to 18 years have not been established.

Patients with renal impairment:

Considering the low dose and the very low systemic exposure (see section 5.2), no dose adjustment is necessary (see section 4.4).

Patients with hepatic impairment:

Considering the low dose and the very low systemic exposure (see section 5.2), no dose adjustment is necessary.

Method of administration

Intracameral use.

The following procedure should be followed:

1. Five minutes before performing the preoperative antiseptic procedure and the first incision, one to two drops of anaesthetic eye drops should be instilled in the eye.
2. At the beginning of surgery, 0.2 ml of Fydrane is slowly injected in only one injection by an ophthalmic surgeon, via intracameral route, through the side port or principal port.

For instructions on handling the medicinal product before administration, see section 6.6.

4.3 Contraindications

- Hypersensitivity to the active substances (tropicamide, phenylephrine hydrochloride and lidocaine hydrochloride) or to any of the excipients listed in section 6.1.
- Known hypersensitivity to anaesthetics of the amide type.
- Known hypersensitivity to atropine derivatives.

4.4 Special warnings and precautions for use

The recommended dose is 0.2 ml of Fydrane; no additional dose should be injected as no significant add-on effect has been demonstrated, and as increased endothelial cell loss was observed.

Corneal endothelial toxicity has not been reported at the recommended dose of Fydrane; nevertheless, due to limited data, this risk cannot be excluded.

There is no clinical experience with Fydrane in:

- insulin-dependent or uncontrolled diabetic patients,
- patients with corneal disease, especially those with any coexisting endothelial cell impairment,
- patients with history of uveitis,
- patients with pupillary abnormalities or presenting an ocular traumatism,
- patients with very dark irides,
- cataract surgery when combined with corneal transplantation.

There is no experience in patients at risk of floppy iris syndrome with Fydrane. Such patients should benefit of a step-by-step pupil dilation strategy starting with the administration of mydriatic eye drops.

There is no clinical experience during cataract surgery with Fydrane in patients treated with topical mydriatics and for whom pupil constriction (or even miosis) occurs during surgery.

Fydrane is not recommended to be used in cataract surgery when combined with vitrectomy, due to the vasoconstricting effects of phenylephrine.

Fydrane is not recommended in subjects with a shallow anterior chamber or a history of acute narrow angle glaucoma.

Special precautions for use:

Fydrane was shown to produce undetectable or very low systemic concentrations of active substances (see section 5.2). Since systemic effects of phenylephrine and lidocaine are dose dependent, it is unlikely that these effects occur with Fydrane. However, as the risk cannot be excluded, it is reminded that:

- Phenylephrine has sympathomimetic activity that might affect patients in the event of hypertension, cardiac disorders, hyperthyroidism, atherosclerosis or prostate disorders and all subjects presenting with a contraindication to the systemic use of pressor amines;
- Lidocaine should be used with caution in patients with epilepsy, myasthenia gravis, cardiac conduction disturbances, congestive heart failure, bradycardia, severe shock, impaired respiratory function or impaired renal function with a creatinine clearance of less than 10mL/minute.

This medicinal product contains less than 1 mmol sodium (23 mg) per dose, i.e. essentially “sodium-free”.

4.5 Interaction with other medicinal products and other forms of interaction

No interaction studies have been performed with Fydrane.

Since the systemic exposure is expected to be very low (see section 5.2), systemic interactions are unlikely.

4.6 Fertility, pregnancy and lactation**Pregnancy**

There are no adequate data from the use of phenylephrine and tropicamide in pregnant women. Animal studies are insufficient with respect to effects on pregnancy, embryonic/foetal development, parturition and postnatal development.

Although animal studies have revealed no evidence of harm to the foetus, lidocaine crosses the placenta and should not be administered during pregnancy.

Even though a negligible systemic uptake is expected, a low systemic exposure cannot be excluded. Therefore, Fydrane should not be used during pregnancy.

Breastfeeding

No data are available concerning the secretion of phenylephrine or tropicamide into breast milk. However, phenylephrine is poorly absorbed orally, implying that absorption by the infant would be negligible. On the other hand, infants may be very sensitive to anticholinergics, and despite the expected negligible systemic exposure, tropicamide is therefore not recommended during breast feeding.

Small amounts of lidocaine are secreted into breast milk and there is a possibility of an allergic reaction in the infant.

Therefore, Fydrane should not be used during breast feeding.

Fertility

There is no information on whether Fydrane may affect fertility in human males or females.

4.7 Effects on ability to drive and use machines

Fydrane has a moderate influence on the ability to drive and use machines, due to its mydriatic effect. Consequently, after cataract surgery with one Fydrane injection, the patient should be advised not to drive and/or use machines while the visual disturbances persist.

4.8 Undesirable effects

Adverse reactions were reported with Fydrane during clinical trials (see section 5.1). Most were ocular and of mild to

moderate intensity.

Summary of the safety profile:

Posterior capsule rupture and cystoid macular oedema are well known complications occurring during or after cataract surgery. They may occur uncommonly (less than 1 case per 100 patients).

Tabulated list of adverse reactions:

Adverse events are categorised by frequency as follows: Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); not known (frequency cannot be estimated from available data).

Adverse reactions, reported during clinical trials, are presented according to System Organ Class in the table below in order of decreased seriousness within each frequency grouping:

| System Organ class | Frequency | Adverse reaction |
|---------------------------------|-----------|---|
| <i>Nervous system disorders</i> | uncommon | Headache |
| <i>Eye disorders</i> | uncommon | Keratitis, Cystoid macular oedema, Intraocular pressure increased, Posterior capsule rupture, Ocular hyperaemia |
| <i>Vascular disorders</i> | uncommon | Hypertension |

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, Earlsfort Terrace, IRL – Dublin 2; Tel: +353 1 6764971; Fax: +353 1 6762517. Website: www.hpra.ie; E-mail: medsafety@hpra.ie.

4.9 Overdose

Due to single administration and low expected systemic passage of Fydrane, overdose is not expected; however a risk of overdose cannot be excluded.

One ampoule of 0.6 ml solution contains 0.12 mg of tropicamide, 1.86 mg of phenylephrine hydrochloride and 6 mg of lidocaine hydrochloride.

The symptoms of phenylephrine ophthalmic overdose are likely to be effects resulting from systemic absorption, including extreme tiredness, sweating, dizziness, a slow heartbeat, and coma. Because severe toxic reaction to phenylephrine is of rapid onset and short duration, treatment is primarily supportive. Prompt injection of a rapidly acting alpha-adrenergic blocking agent such as phentolamine (dose 2 to 5 mg in intravenous use) has been recommended.

The symptoms of tropicamide ophthalmic overdose include headache, fast heartbeat, dry mouth and skin, unusual drowsiness, and flushing. Systemic effects from tropicamide are not expected. Should an overdose occur causing local effects, e.g. sustained mydriasis, pilocarpine or 0.25% w/v physostigmine should be applied.

In the event of excessive absorption of lidocaine into the bloodstream, symptoms may include CNS effects (such as convulsions, unconsciousness and possibly respiratory arrest) and cardiovascular reactions (such as hypotension, myocardial depression, bradycardia and possibly cardiac arrest). Treatment of a patient suffering from systemic toxicity of lidocaine consists of arresting the convulsions and ensuring adequate ventilation with oxygen, if necessary by assisted or controlled ventilation (respiration).

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: MYDRIATICS and CYCLOPLEGICS, Tropicamide combinations, ATC code: S01FA56.

Fydrane is a solution for intracameral injection which combines two synthetic mydriatic agents (tropicamide – anticholinergic, and phenylephrine - alpha sympathomimetic) and one local anaesthetic (lidocaine hydrochloride).

Mechanism of action:

Phenylephrine is a direct acting sympathomimetic agent. It causes mydriasis via the stimulation of alpha-adrenergic receptors of the pupillary dilator (the resulting contraction of the pupillary dilator causes pupil dilation). There is almost no cycloplegic effect.

Tropicamide is a parasympatholytic agent, which acts by binding to and blocking the M4 muscarinic receptors of the eye muscles. It prevents the iris sphincter muscle and ciliary body muscle from responding to cholinergic stimulation, producing dilation of the pupil and paralysis of the ciliary muscle (cyclopegia).

Lidocaine is a local anaesthetic of the amide type. It acts by inhibiting the ionic refluxes required for the initiation and conduction of impulses, thereby stabilising the neuronal membrane.

Pharmacodynamic effects

Although tropicamide as a monotherapy produces both mydriasis and cycloplegia, additional mydriasis occurs if sympathomimetic agents such as phenylephrine are used simultaneously. Such synergistic combinations are commonly prescribed to achieve maximal dilation of the pupil for cataract extraction.

As an average, 95% of the dilation measured before the viscoelastic injection was obtained within 30 seconds after a single 200-µL intracameral injection of Fydrane during phase II clinical study. Pupil sizes observed during phase II and III clinical trials are presented in the table below (patients who received a single 200-µL intracameral injection of Fydrane):

| | Phase II study, n=24 | | Phase III study, n=181 | |
|------------------------|---|--|--|---------------------------|
| | Within 30 seconds after Fydrane injection | After injection of Fydrane, and then of viscoelastic | After injection of Fydrane, and then of viscoelastic | Just before IOL injection |
| Pupil size (mm) | | | | |
| Mean (SD) | 6.7 (0.7) | 7.7 (0.7) | 7.8 (0.8) | 7.9 (0.9) |
| Median | 6.7 | 7.7 | 7.8 | 7.9 |

In phase III study, after a single 200-µL injection of Fydrane and injection of viscoelastic (just before capsulorhexis), the pupil size was at least 7 mm for 86.7% of the patients. In these clinical phase II and III studies, mydriasis with Fydrane was demonstrated to be stable until the end of the surgery.

Return to normal pupil size is known to be obtained after 5-7 hours.

Clinical efficacy and safety

Clinical efficacy:

The mydriatic and anaesthetic effects of Fydrane were evaluated in a phase III, multicentre, randomised, open study in comparison with a standard topical treatment (phenylephrine and tropicamide) in 555 patients undergoing cataract surgery with a pupil diameter ≥ 7 mm following topical mydriatic application. Tetracaine 1% eye drops was instilled 5 minutes and 1 minute before surgery in both groups.

Mydriasis:

Non-inferiority of Fydrane *versus* the Reference treatment (tropicamide 0.5% eye drops and phenylephrine 10% eye drops, application of one drop of each repeated 3 times prior a surgery) was demonstrated for the primary and co-primary efficacy criteria in the mITT Population (see Table below):

| mITT Population | Fydrane | Reference Treatment | Difference (%) between groups (Fydrane - Reference) [95% CI] |
|---|--------------------------------------|--------------------------------------|--|
| Primary efficacy criterion Number (%) of responders* 95% CI | N=268 265 (98.9) [96.8 ; 99.8] | N=281 266 (94.7) [91.3 ; 97.0] | 4.2 [-4.2 ; 12.6] |
| Co-primary efficacy criterion Number (%) of responders** 95% CI | N=250 246 (98.4) [96.0 ; 99.6] | N=261 246 (94.3) [90.7 ; 96.7] | 4.1 [-4.5 ; 12.8] |
| * A responder was defined as a patient for whom the capsulorhexis was performed without use of any additive mydriatic treatment | | | |
| ** A responder was defined as a patient for whom the capsulorhexis was performed without use of any additive mydriatic treatment and for whom the pupil size just before capsulorhexis was ≥ 5.5 mm. | | | |

During the phase III study, in the Fydrane group (N=268), 197 patients received a single 200-µL intracameral injection and 71 received an additional 100-µL intracameral injection which has not demonstrated a significant add-on effect and for which increased endothelial cell loss was observed.

The data analysis on the patients with a single 200-µL intracameral injection, for whom the capsulorhexis was performed without use of any additive mydriatic treatment and for whom the pupil size just before capsulorhexis was > 6 mm, is presented in the table below.

| | Fydrane 200-µL | Reference Treatment | Difference (%) between groups (Fydrane 200-µL - Reference) [95% CI] |
|--|--|---|---|
| N Number (%) of patients with no additive mydriatic treatment and with the pupil size just before capsulorhexis > 6 mm 95% CI | N=181 180 (99.4) [97.0; 100.0] | N=261 246 (94.3) [90.7; 96.7] | 5.2 [-4.3; 14.6] |

Anaesthesia:
Before intraocular lens injection, the patients' comfort was statistically significantly better with Fydrane (p=0.034), and no statistically significant difference between groups was seen at the other time points of the surgery (before viscoelastic injection, capsulorhexis and cefuroxime injection).

5.2 Pharmacokinetic properties

No ocular pharmacokinetic data are available for Fydrane.

Following intracameral injection of Fydrane in 15 patients undergoing cataract surgery, the concentrations of the active ingredients assayed in plasma 2, 12 and 30 min post-injection were compared to a standard topical treatment (phenylephrine 10% eye drops and tropicamide 0.5% eye drops). Regarding tropicamide, all patients in Fydrane group were below the limit of quantification (< 0.1 ng/mL) whereas all patients in the Reference group had a level above this

limit. Level of phenylephrine (quantification limit < 0.1 ng/mL) was not detectible in all patients of the Fydrane group with exception of 2 patients (maximum 0.59 ng/mL) versus all patients of the Reference group with a level above limit of quantitation (maximum 1.42 ng/mL). The plasma lidocaine concentration was measured in all Fydrane -treated patients with a highest concentration of 1.45 ng/mL (well below the values causing some systemic effects: between 1,500 and 5,000 µg/mL).

5.3 Preclinical safety data

In rabbits, the ocular tolerance after single intracameral administration of 200µL of Fydrane with or without rinsing (slit-lamp, aqueous flare, corneal thickness and cellular density of the endothelium, electroretinography and histology) was very good in the seven days post-dosing period.

Signs of ocular intolerance were only observed for formulations with higher concentrations of the three active substances (at or above 5 times the concentrations in Fydrane). The highest tested concentration (10 fold) showed increases in the thickness of the cornea, and severe ocular changes resulted in one animal being sacrificed on Day 3.

Systemic toxicity of the fixed combination of phenylephrine, tropicamide and lidocaine has not been investigated. Nevertheless, since the ophthalmological safety of the three individual substances is considered established and Fydrane is only administered by single intracameral injection, no particular risk is expected for the combination.

Likewise, the safety pharmacology, genotoxicity and reproduction toxicity of the individual substances of the fixed combination has not been evaluated. In rats, administration of phenylephrine (12.5 mg/kg, s.c.) resulted in reduced uterine blood flow (86.8% reduction in about 15 minutes), thereby exhibiting foetotoxic and co-teratogenic properties. For lidocaine, no teratogenic effects were observed in studies of embryonic/foetal development in rats and rabbits. Embryotoxicity and a reduction in postnatal survival were only observed at maternally toxic doses. Lidocaine was also not genotoxic.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium chloride
Disodium phosphate dodecahydrate
Disodium phosphate dihydrate
Disodium edetate
Water for injections

6.2 Incompatibilities

No incompatibility with most commonly used products in cataract surgery was reported in literature with the active ingredients, and during clinical trials. For usual viscoelastics, this was also confirmed by pharmaceutical interaction test.

6.3 Shelf life

3 years

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

One 1 ml sterile brown glass (type I) ampoule filled with 0.6 ml of solution for injection, per paper/PVC blister. Box of 1, 20 and 100 ampoules together with respectively 1, 20 and 100, 5-micron sterile filter needles.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product

For single eye use only.

Use immediately after first opening of the ampoule.

Warning: Do not use if blister or peelable backing is damaged or broken. Open under aseptic conditions only. The content of the blister are guaranteed as sterile.

The solution should be visually inspected and should only be used if it is a clear, slightly brownish-yellow and practically free from visible particles solution.

Fydrane must be administered by intracameral injection, by an ophthalmic surgeon in the recommended aseptic conditions of cataract surgery.

To prepare the product for intracameral injection, please adhere to the following instructions:

1. Inspect unopened blister to ensure that it is intact. Peel open blister.
2. Break open the ampoule containing the drug product. The One Point Cut (OPC) ampoule must be opened as follows: Hold the bottom part of the ampoule with the thumb pointing to the coloured point. Grasp the top of the ampoule with the other hand, positioning the thumb at the coloured point and press back to break at the existing cut under the point.
3. Assemble the 5-micron filter sterile needle (provided) onto a sterile syringe. Remove the 5-micron filter sterile needle protector and withdraw at least 0.2 ml of the solution for injection from the ampoule into the syringe.
4. Disconnect the needle from the syringe and assemble the syringe with an appropriate anterior chamber cannula.
5. Carefully expel the air from the syringe. Adjust to 0.2 ml. The syringe is ready for injection.
6. Slowly inject the 0.2 ml syringe volume into the anterior chamber of the eye, as only one injection, through the side port or principal port.
7. After use, discard the remaining solution appropriately. Do not keep it for subsequent use.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements. Discard used needles in a sharps container.

7 MARKETING AUTHORISATION HOLDER

Laboratoires Thea
12, rue Louis Blériot
63017 Clermont-Ferrand Cedex 2
France

8 MARKETING AUTHORISATION NUMBER

PA1107/012/001

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 28th August 2015

10 DATE OF REVISION OF THE TEXT